

World Journal of *Clinical Cases*

World J Clin Cases 2024 December 16; 12(35): 6754-6863



EDITORIAL

- 6754** Update on the aetiopathogenesis of obstructive sleep apnea: Role of inflammatory and immune mediated mechanisms
Nag DS, Varghese K, Swain A, Patel R, Sahu S, Sam M
- 6760** Intensive care unit-acquired weakness: Unveiling significant risk factors and preemptive strategies through machine learning
He XY, Zhao YH, Wan QW, Tang FS
- 6764** Advancing oral cancer care: Insights from Tongluo Jiedu prescription
Cheng CY, Hao WR, Liu JC, Cheng TH
- 6770** Effects of atrial septal defects on the cardiac conduction system
Kang JH, Wu HY, Long WJ
- 6775** Periodontitis and chronic kidney disease: A bidirectional relationship based on inflammation and oxidative stress
Martínez Nieto M, De Leon Rodríguez ML, Anaya Macías RDC, Lomeli Martínez SM
- 6782** Cytokine release syndrome induced by anti-programmed death-1 treatment in a psoriasis patient: A dark side of immune checkpoint inhibitors
Maldonado-García JL, Fragozo A, Pavón L

REVIEW

- 6791** Acellular dermal matrices in reconstructive surgery; history, current implications and future perspectives for surgeons
Dilek ÖF, Sevim KZ, Dilek ON

ORIGINAL ARTICLE**Retrospective Study**

- 6808** Comprehensive epidemiological assessment of trauma incidents at a level I trauma center
Su ZY, Wei H, Wang WN, Lin YF, He YL, Liu Y, Lin RB, Liu YT, Michael N

SYSTEMATIC REVIEWS

- 6815** Gut microbiota changes associated with frailty in older adults: A systematic review of observational studies
Wen NN, Sun LW, Geng Q, Zheng GH

CASE REPORT

- 6826** *psk1* virulence gene-induced pulmonary and systemic tuberculosis in a young woman with normal immune function: A case report
Wu F, Yang B, Xiao Y, Ren LL, Chen HY, Hu XL, Pan YY, Chen YS, Li HR
- 6834** Rare primary gastric peripheral T-cell lymphoma not otherwise specified: A case report
Jang HR, Lee K, Lim KH
- 6840** Cat scratch disease in children with nocturnal fever: A case report
Yin QL, Liu YQ, Zhang HM, Zhang YL, Qi SM, Wen JQ, Zhang WH

LETTER TO THE EDITOR

- 6848** Understanding network meta-analysis
Au SCL
- 6851** Effects of foot reflexology on disease
He MY, Ud Din MJ, Xu HF, Wang SY, Ying GH, Qian H, Wu B, Qi HD, Wang X, Zhang G
- 6855** Clinical landscape and treatment of acute non-variceal upper gastrointestinal bleeding: Insights from a high-volume center in Shaanxi, China
Improta L
- 6859** Role of high-dose amoxicillin dual therapy for *Helicobacter pylori* eradication in an Irish cohort: A prospective study
Palmirotta R, Cafiero C, Colella M

ABOUT COVER

Peer Reviewer of *World Journal of Clinical Cases*, Muhamad Zakaria Brimo Alsaman, MD, Associate Chief Physician, Department of Vascular Surgery, Al-Razi Hospital, Aleppo, Syria; Faculty of Medicine, University of Aleppo, Aleppo 12212, Syria. dr.zakariabrimo@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now abstracted and indexed in PubMed, PubMed Central, *Reference Citation Analysis*, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for *WJCC* as 1.0; JIF without journal self cites: 0.9; 5-year JIF: 1.1; JIF Rank: 168/325 in medicine, general and internal; JIF Quartile: Q3; and 5-year JIF Quartile: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Lei Zhang*, Production Department Director: *Si Zhao*, Cover Editor: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

December 16, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Intensive care unit-acquired weakness: Unveiling significant risk factors and preemptive strategies through machine learning

Xiao-Yu He, Yi-Huan Zhao, Qian-Wen Wan, Fu-Shan Tang

Specialty type: Computer science, artificial intelligence

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade B

Novelty: Grade B

Creativity or Innovation: Grade B

Scientific Significance: Grade B

P-Reviewer: Nagamine T

Received: March 18, 2024

Revised: August 22, 2024

Accepted: September 4, 2024

Published online: December 16, 2024

Processing time: 219 Days and 20.4 Hours



Xiao-Yu He, Yi-Huan Zhao, Qian-Wen Wan, Fu-Shan Tang, Department of Clinical Pharmacy, Key Laboratory of Basic Pharmacology of Guizhou Province and School of Pharmacy, Zunyi Medical University, Zunyi 563006, Guizhou Province, China

Corresponding author: Fu-Shan Tang, PhD, Professor, Department of Clinical Pharmacy, Key Laboratory of Basic Pharmacology of Guizhou Province and School of Pharmacy, Zunyi Medical University, No. 6 Xuefu West Road, Xinpu New District, Zunyi 563006, Guizhou Province, China. fstang@vip.163.com

Abstract

This editorial discusses an article recently published in the *World Journal of Clinical Cases*, focusing on risk factors associated with intensive care unit-acquired weakness (ICU-AW). ICU-AW is a serious neuromuscular complication seen in critically ill patients, characterized by muscle dysfunction, weakness, and sensory impairments. Post-discharge, patients may encounter various obstacles impacting their quality of life. The pathogenesis involves intricate changes in muscle and nerve function, potentially leading to significant disabilities. Given its global significance, ICU-AW has become a key research area. The study identified critical risk factors using a multilayer perceptron neural network model, highlighting the impact of intensive care unit stay duration and mechanical ventilation duration on ICU-AW. Recommendations were provided for preventing ICU-AW, emphasizing comprehensive interventions and risk factor mitigation. This editorial stresses the importance of external validation, cross-validation, and model transparency to enhance model reliability. Moreover, the application of machine learning in clinical medicine has demonstrated clear benefits in improving disease understanding and treatment decisions. While machine learning presents opportunities, challenges such as model reliability and data management necessitate thorough validation and ethical considerations. In conclusion, integrating machine learning into healthcare offers significant potential and challenges. Enhancing data management, validating models, and upholding ethical standards are crucial for maximizing the benefits of machine learning in clinical practice.

Key Words: Intensive care unit-acquired weakness; Risk factors; Machine learning; Clinical medicine; Treatment decision

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This editorial emphasizes the importance of recognizing the risk factors linked to intensive care unit-acquired weakness and highlights the vital role of machine learning in identifying and managing these factors to improve patient outcomes and enhance the quality of care in clinical settings.

Citation: He XY, Zhao YH, Wan QW, Tang FS. Intensive care unit-acquired weakness: Unveiling significant risk factors and preemptive strategies through machine learning. *World J Clin Cases* 2024; 12(35): 6760-6763

URL: <https://www.wjgnet.com/2307-8960/full/v12/i35/6760.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v12.i35.6760>

INTRODUCTION

Intensive care unit-acquired weakness (ICU-AW) is a debilitating neuromuscular complication that occurs in patients undergoing intensive care treatment in the intensive care unit (ICU)[1]. It is characterized by skeletal muscle dysfunction, leading to various clinical manifestations such as limb muscle weakness, atrophy, diminished deep tendon reflexes, and sensory impairments[2]. Importantly, patients with ICU-AW may also experience a range of challenges after being discharged from the hospital, including physical dysfunction, cognitive impairment, depression, and anxiety disorders [3]. The pathogenesis of ICU-AW is complex and involves intricate functional and structural changes in both muscles and nerves. This condition not only has the potential to result in severe outcomes like tetraplegia or paraplegia but can also lead to long-term disability, significantly impacting the quality of life for patients post-discharge. Given its profound impact on ICU patients, ICU-AW has become a prominent research topic of great interest to scholars worldwide, attracting extensive attention both domestically and internationally[4].

DISCUSSION

An article published by Wang and Long[5] offers a comprehensive exploration of the risk factors associated with ICU-AW. Through the use of a multilayer perceptron neural network model, the study successfully identified critical risk factors for ICU-AW, emphasizing the substantial impact of ICU stay duration and mechanical ventilation duration on its development. These findings provide actionable recommendations for managing these factors to mitigate the risk of ICU-AW within clinical settings. However, it is essential to recognize that the occurrence of ICU-AW is influenced by various other factors, necessitating comprehensive interventions to reduce its incidence. The study's insights serve as valuable reference information for clinicians, aiding in informed decision-making for the prevention and treatment of ICU-AW. It is recommended to further explain the differences between these risk factors identified by machine learning and actual medical practices and to process the data through an iteratively updated machine-learning approach.

While the article demonstrates significant progress in researching ICU-AW risk factors, there are opportunities for improvement in model construction and validation. To enhance the model's robustness, future studies should incorporate external validation[6] to ensure generalizability and employ cross-validation techniques to improve stability and reliability. To avoid the bias that older data may have on machine learning models, it is recommended to group the data by year or time period. This helps to identify the impact of data over different time periods on the model results and to assess the possibility of time bias. Additionally, comparing the multilayer perceptron neural network approach with other common machine learning algorithms would offer valuable insights into its advantages. It is also recommended that researchers share optimal model parameters, make data and model code publicly available, and encourage reproducibility of the study's results.

We propose the following specific recommendations for preventive strategies and interventions regarding ICU-AW. First, the article suggests that reducing the length of ICU stay and duration of mechanical ventilation on ICU-AW effectively lowers the probability of ICU-AW. This finding highlights the significance of early rehabilitation and ventilator withdrawal, offering a new perspective for clinical practice. Second, nutritional interventions and exercise interventions emerge as key strategies for preventing ICU-AW[7]. Nutritional support helps maintain muscle tissue metabolic balance and reduce muscle breakdown, while exercise interventions preserve muscle mass and function through muscle contraction and strength training. The simultaneous implementation of both approaches is expected to yield superior preventive outcomes. Lastly, the application of an early multidisciplinary collaboration in critically ill patient rehabilitation has demonstrated significant achievements. This early multidisciplinary collaboration facilitates close interdisciplinary and cross-disciplinary cooperation, delivering comprehensive and systematic rehabilitation services. Not only does this approach effectively prevent complications like ICU-AW, but it also substantially shortens patients' rehabilitation duration and enhances their daily functioning, muscle strength, and overall, quality of life[8]. This successful practice offers a novel and effective intervention strategy for rehabilitating critically ill patients. Future research is required to determine the extent to which efforts to prevent muscle weakness in the ICU are effective in terms of actual patient prognosis and activities of daily living. The effect of specific strategies such as shortening the duration of mechanical ventilation, early rehabilitation and intervention during deventilation, active nutrition intervention, and even exercise intervention during mechanical ventilation on the prevention of muscle decline in the model should be estimated by machine learning.

All in all, while the article provides valuable insights, it is crucial to enhance the model's stability and reliability through improvements in out-of-sample validation, cross-validation, algorithm comparison, and ensuring transparency in data and code. Moving forward, concrete preventive strategies and interventions for ICU-AW, including reducing ICU stay and mechanical ventilation duration, implementing nutritional and exercise interventions, and promoting early multidisciplinary collaboration in critically ill patient rehabilitation, are recommended. These suggestions provide essential guidance for future clinical practice and research, with the aim of effectively preventing ICU-AW, improving patient recovery outcomes, and enhancing overall quality of life.

CLINICAL APPLICATIONS AND CHALLENGES OF MACHINE LEARNING

In recent years, the utilization of machine learning in clinical medicine has expanded, showcasing its unique advantages. Machine learning provides healthcare professionals with a deeper understanding of disease causation and influencing factors, offering a scientific basis and decision support for clinical treatment. For instance, in colorectal cancer diagnosis, traditional methods face limitations that can be overcome by integrating machine learning with medical imaging, which enables more effective early screening[9]. Machine learning also aids in monitoring aggressive lymphomas and provides critical evidence for cases requiring early intervention[10]. Furthermore, in predicting macrosomia during pregnancy, machine learning empowers healthcare professionals to make more precise risk assessments, facilitating timely interventions for the well-being of both mothers and children[11].

In the realm of clinical pharmacy, machine learning demonstrates significant potential across drug development and clinical implementation. By leveraging drug-target knowledge to train predictive models capable of assessing interactions between new drugs and targets, machine learning offers valuable insights for drug development[12]. Moreover, when constructing predictive models for adverse drug reactions, integrating data from various sources helps reduce the risks associated with such reactions for patients and the healthcare system[13]. Additionally, machine learning-based Clinical Decision Support Systems (CDSSs) play a pivotal role in enhancing patient safety by providing clinicians with clinically valid alerts, improving decision-making processes, and minimizing medication errors[14]. The implementation of CDSS not only ensures patient safety but also enhances the efficiency and reliability of clinical decision-making within healthcare systems.

Despite the widespread application of machine learning, we must address potential challenges. Ensuring the reliability and stability of constructed and validated machine learning models is crucial. Rigorous data set divisions, cross-validation techniques, and other tools are essential to thoroughly evaluate model performance[15]. Efficient collection, processing, and utilization of the increasing volume of clinical data pose another challenge, necessitating the development of robust data processing methods, enhanced data quality management, and reinforced data security measures. Furthermore, legal and ethical issues in the clinical application of machine learning should be addressed through regulatory frameworks, legislative measures, and strengthening the ethical review process[16].

CONCLUSION

In conclusion, while machine learning offers immense potential in clinical medicine and pharmacy, addressing associated challenges is paramount. Through continuous technological advancements, improved data management practices, and ethical reviews, machine learning is poised to play an increasingly vital role in the pharmaceutical sector, contributing significantly to human health. Anticipating further progress in machine learning applications across diverse fields, we look forward to the continued evolution and impact of this technology.

FOOTNOTES

Author contributions: He XY and Zhao YH contributed equally to this work; He XY and Zhao YH contributed to the manuscript outline and composed the initial draft; He XY and Wan QW were responsible for sourcing and organizing the relevant literature; Tang FS and Zhao YH originated the concept for this manuscript; Tang FS provided supervision, reviewed the paper, and finalized the manuscript; all authors have read and approved the final manuscript.

Conflict-of-interest statement: All the authors have nothing to disclose for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: China

ORCID number: Xiao-Yu He 0009-0009-5698-5789; Yi-Huan Zhao 0000-0002-9870-246X; Qian-Wen Wan 0009-0005-3398-0898; Fu-Shan Tang 0000-0001-8779-1041.

S-Editor: Liu JH

L-Editor: Wang TQ

P-Editor: Wang WB

REFERENCES

- 1 **Wang W**, Xu C, Ma X, Zhang X, Xie P. Intensive Care Unit-Acquired Weakness: A Review of Recent Progress With a Look Toward the Future. *Front Med (Lausanne)* 2020; **7**: 559789 [PMID: 33330523 DOI: 10.3389/fmed.2020.559789]
- 2 **Li Z**, Cai Y, Zhang Q, Zhang P, Sun R, Jiang H, Wan J, Wu F, Wang X, Tao X. Intensive care unit acquired weakness: A protocol for an overview of systematic reviews and meta-analysis. *Medicine (Baltimore)* 2020; **99**: e21926 [PMID: 32846861 DOI: 10.1097/MD.00000000000021926]
- 3 **Zhang W**, Wu J, Gu Q, Gu Y, Zhao Y, Ge X, Sun X, Lian J, Zeng Q. Changes in muscle ultrasound for the diagnosis of intensive care unit acquired weakness in critically ill patients. *Sci Rep* 2021; **11**: 18280 [PMID: 34521934 DOI: 10.1038/s41598-021-97680-y]
- 4 **Yang Z**, Wang X, Wang F, Peng Z, Fan Y. A systematic review and meta-analysis of risk factors for intensive care unit acquired weakness. *Medicine (Baltimore)* 2022; **101**: e31405 [PMID: 36316900 DOI: 10.1097/MD.00000000000031405]
- 5 **Wang L**, Long DY. Significant risk factors for intensive care unit-acquired weakness: A processing strategy based on repeated machine learning. *World J Clin Cases* 2024; **12**: 1235-1242 [PMID: 38524515 DOI: 10.12998/wjcc.v12.i7.1235]
- 6 **Zhang W**, Tang Y, Liu H, Yuan LP, Wang CC, Chen SF, Huang J, Xiao XY. Risk prediction models for intensive care unit-acquired weakness in intensive care unit patients: A systematic review. *PLoS One* 2021; **16**: e0257768 [PMID: 34559850 DOI: 10.1371/journal.pone.0257768]
- 7 **Heyland DK**, Stapleton RD, Mourtzakis M, Hough CL, Morris P, Deutz NE, Colantuoni E, Day A, Prado CM, Needham DM. Combining nutrition and exercise to optimize survival and recovery from critical illness: Conceptual and methodological issues. *Clin Nutr* 2016; **35**: 1196-1206 [PMID: 26212171 DOI: 10.1016/j.clnu.2015.07.003]
- 8 **Wang B**, He X, Tian S, Feng C, Feng W, Song L. Influence of Early Multidisciplinary Collaboration on Prevention of ICU-Acquired Weakness in Critically Ill Patients. *Dis Markers* 2022; **2022**: 3823368 [PMID: 35942131 DOI: 10.1155/2022/3823368]
- 9 **Peng Y**, Deng H. Medical image fusion based on machine learning for health diagnosis and monitoring of colorectal cancer. *BMC Med Imaging* 2024; **24**: 24 [PMID: 38267874 DOI: 10.1186/s12880-024-01207-6]
- 10 **Bai A**, Si M, Xue P, Qu Y, Jiang Y. Artificial intelligence performance in detecting lymphoma from medical imaging: a systematic review and meta-analysis. *BMC Med Inform Decis Mak* 2024; **24**: 13 [PMID: 38191361 DOI: 10.1186/s12911-023-02397-9]
- 11 **Shamshuzzoha M**, Islam MM. Early Prediction Model of Macrosomia Using Machine Learning for Clinical Decision Support. *Diagnostics (Basel)* 2023; **13** [PMID: 37685292 DOI: 10.3390/diagnostics13172754]
- 12 **Xu L**, Ru X, Song R. Application of Machine Learning for Drug-Target Interaction Prediction. *Front Genet* 2021; **12**: 680117 [PMID: 34234813 DOI: 10.3389/fgene.2021.680117]
- 13 **Nguyen DA**, Nguyen CH, Mamitsuka H. A survey on adverse drug reaction studies: data, tasks and machine learning methods. *Brief Bioinform* 2021; **22**: 164-177 [PMID: 31838499 DOI: 10.1093/bib/bbz140]
- 14 **Chen CY**, Chen YL, Scholl J, Yang HC, Li YJ. Ability of machine-learning based clinical decision support system to reduce alert fatigue, wrong-drug errors, and alert users about look alike, sound alike medication. *Comput Methods Programs Biomed* 2024; **243**: 107869 [PMID: 37924770 DOI: 10.1016/j.cmpb.2023.107869]
- 15 **Aittokallio T**. What are the current challenges for machine learning in drug discovery and repurposing? *Expert Opin Drug Discov* 2022; **17**: 423-425 [PMID: 35255749 DOI: 10.1080/17460441.2022.2050694]
- 16 **Vayena E**, Blasimme A, Cohen IG. Machine learning in medicine: Addressing ethical challenges. *PLoS Med* 2018; **15**: e1002689 [PMID: 30399149 DOI: 10.1371/journal.pmed.1002689]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

