

Supplementary Table 1 South West Oncology Group (SWOG) toxic grading score

	0	1	2	3	4
Fever	None	37.1-38.0 °C	38.1-40.0 °C	> 40.0 °C	> 40.0 °C for more than 24 hours or accompanied by hypotension
Nausea	None	Able to eat reasonable intake	Intake significantly decreased, but can eat	No significant intake	-
Vomiting	None	1 episode in 24 hours	2-5 episodes in 24 hours	6-10 episodes in 24 hours	> 10 episodes in 24 hours or requiring parenteral support
Abdominal pain	None	Requiring non- narcotics	Requiring oral narcotics	Requiring parenteral narcotics	Uncontrollable

Supplementary Table 2 Common Toxicity Criteria Adverse Event (CTCAE version 5)

	Grade 1	Grade 2	Grade 3	Grade 4
Fever	38.0-39.0 °C	> 39.0-40.0 °C	> 40.0 °C for < 24 h	
Nausea	Loss of appetite without alteration in eating habits	Oral intake decreased without significant weight loss, dehydration or malnutrition	Inadequate oral caloric or fluid intake; tube feeding, TPN, or hospitalization indicated	
Vomiting	1-2 episodes (separated by 5 min) in 24 h. Intervention not indicated	3-5 episodes (separated by 5 min) in 24 h. Medical intervention indicated	> 6 episodes (separated by 5 min) in 24 h; tube feeding, TPN or hospitalization indicated	Life-threatening consequences; urgent intervention indicated
Abdominal pain	Mild pain; limiting instrumental ADL	Moderate pain	Severe pain; limiting self-care ADL	-
AST	> ULN-3.0 × ULN	> 3.0-5.0 × ULN	> 5.0-20.0 × ULN	> 20.0 × ULN
ALT	> ULN-3.0 × ULN	> 3.0-5.0 × ULN	> 5.0-20.0 × ULN	> 20.0 × ULN
ALP	> ULN-2.5 × ULN if baseline was normal	> 2.5-5.0 × ULN if baseline was normal	> 5.0-20.0 × ULN if baseline was normal	> 20.0 × ULN if baseline was normal
	2.0-2.5 × baseline if baseline was abnormal	> 2.5-5.0 × baseline if baseline was abnormal	> 5.0-20.0 × baseline if baseline was abnormal	> 20.0 × baseline if baseline was abnormal

Total bilirubin	> ULN-1.5 × ULN if baseline was normal	> 1.5-3.0 × ULN if baseline was normal	> 3.0-10.0 × ULN if baseline was normal	> 10.0 × ULN if baseline was normal
	> 1.5-2.5 × ULN if baseline was abnormal	> 2.5-3.0 × ULN if baseline was abnormal	> 3.0-10.0 × ULN if baseline was abnormal	> 10.0 × ULN if baseline was abnormal
Hyperglycemia	FPG > ULN-160 mg/dL	FPG > 160-250 mg/dL	FPG > 250-500 mg/dL	FPG > 500 mg/dL
	Abnormal glucose above baseline with no medical intervention	Change in daily management from BL for a diabetic; oral Anti-glycemic agent initiated	Insulin initiated; therapy hospitalization indicated	Life-threatening consequences; urgent intervention indicated

ADL: Activities of daily living; ALP: Alkaline phosphatase; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; FPG : fasting plasma glucose; ULN: Upper limit of normal

