

7

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 39864

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Abnormal expression of HMGB-3 significantly associated with malignant transformation of hepatocytes

Zheng WJ *et al.* HMGB-3 in HCC

Wen-Jie Zheng, Min Yao, Miao Fang, Li Wang, Zhi-Zhen Dong, Deng-Fu Yao

Abstract

AIM

To explore the relationship between dynamic expression of high mobility group box-3 (HMGB-3) and malignant transformation of hepatocytes.

Match Overview

1	Crossref 113 words Wang, Li, Min Yao, Zhizhen Dong, Yun Zhang, and Deng fu Yao. "Circulating specific biomarkers in diagnosis of ..."	3%
2	Internet 94 words crawled on 14-Jan-2018 www.wjgnet.com	2%
3	Internet 24 words crawled on 28-Aug-2016 www.semanticscholar.org	1%
4	Internet 21 words crawled on 27-Jul-2016 diagnosticpathology.biomedcentral.com	1%
5	Internet 19 words crawled on 11-Jun-2018 www.oncotarget.com	<1%
6	Internet 18 words crawled on 22-Oct-2017 www.impactjournals.com	<1%
7	Internet 16 words crawled on 05-Oct-2017 real.mtak.hu	<1%
8	Crossref 14 words R. Ji. "Human type II pneumocyte chemotactic responses to CXCR3 activation are mediated by splice variant A", A	<1%
9	Crossref 12 words Wenjie Zheng, Min Yao, Qi Qian, Wenli Sai, Liwei Qiu, Junling Yang, Wei Wu, Zhizhen Dong, Dengfu Yao. "Oncog	<1%

找到约 20,800 条结果 (用时 0.48 秒)

Emerging Role of High-Mobility Group Box 1 (HMGB1) in Liver Diseases

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3883963/>  [翻译此页](#)

作者: R Chen - 2013 - 被引用次数: 56 - 相关文章

2013年11月7日 - Damage-associated molecular pattern (DAMP) molecules are essential for the ... by hepatocytes in I/R. TLR4 plays a central role in liver I/R (Figure 3). ... Interestingly, pretreatment of mice with HMGB1 protein significantly ... (b) the level of HMGB1 expression and release in hepatocytes correlated with peak ...

Abstract · HMGB1 RECEPTORS · HMGB1 AND LIVER I/R · HMGB1 AND VIRAL ...

Gankyrin drives malignant transformation of chronic liver damage ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4669699/>  [翻译此页](#)

作者: X Zhao - 2015 - 被引用次数: 15 - 相关文章

2015年5月7日 - Gankyrin in hepatocytes enhances CCl4-mediated liver fibrosis ... Consistently, real-time PCR analysis also confirmed higher expression of liver fibrosis-related genes TGF- β 1, ... As expected, morphologic analysis showed more abnormal ... Accordingly, microarray assay revealed significant upregulation or ...

Nonalcoholic Lipid Accumulation and Hepatocyte Malignant ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4913080/>  [翻译此页](#)

作者: J Gu - 2016 - 被引用次数: 8 - 相关文章

2016年6月15日 - Hepatocellular carcinoma (HCC) is the 5th most common malignant ... lead to malignant transformation of infected liver cells remain largely unknown. ... 3. CPT-II gene mutation and hepatic lipid accumulation. A. CPT-II gene exons 1-5. ... level of leptin expression is significantly higher in NAFLD patients, ...

缺少字词: hmgb

Oncogenic role of the Notch pathway in primary liver cancer (Review ...

<https://www.spandidos-publications.com/10.3892/ol.2016.4609> - [翻译此页](#)

作者: J Lu - 2016 - 被引用次数: 9 - 相关文章

2016年5月18日 - The Notch signaling pathway is significantly associated with liver disease (23). ... Thus, Notch receptor expression is abnormal in HCC (27). ... The receptors are transmembrane

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 20,800 条结果 (用时 0.48 秒)

Emerging Role of High-Mobility Group Box 1 (HMGB1) in Liver Diseases

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3883963/> [▼ 翻译此页](#)

作者: R Chen - 2013 - 被引用次数: 56 - 相关文章

2013年11月7日 - Damage-associated molecular pattern (DAMP) molecules are essential for the ... by hepatocytes in I/R. TLR4 plays a central role in liver I/R (Figure 3). ... Interestingly, pretreatment of mice with HMGB1 protein significantly ... (b) the level of HMGB1 expression and release in hepatocytes correlated with peak ...

[Abstract](#) · [HMGB1 RECEPTORS](#) · [HMGB1 AND LIVER I/R](#) · [HMGB1 AND VIRAL ...](#)

Gankyrin drives malignant transformation of chronic liver damage ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4669699/> [▼ 翻译此页](#)

作者: X Zhao - 2015 - 被引用次数: 15 - 相关文章

2015年5月7日 - Gankyrin in hepatocytes enhances CCl4-mediated liver fibrosis ... Consistently, real-time PCR analysis also confirmed higher expression of liver fibrosis-related genes TGF- β 1, ... As expected, morphologic analysis showed more abnormal ... Accordingly, microarray assay revealed significant upregulation or ...

Nonalcoholic Lipid Accumulation and Hepatocyte Malignant ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4913080/> [▼ 翻译此页](#)

作者: J Gu - 2016 - 被引用次数: 8 - 相关文章

2016年6月15日 - Hepatocellular carcinoma (HCC) is the 5th most common malignant ... lead to malignant transformation of infected liver cells remain largely unknown. ... 3. CPT-II gene mutation and hepatic lipid accumulation. A. CPT-II gene exons 1-5. ... level of leptin expression is significantly higher in NAFLD patients, ...

缺少字词: hmgb

Oncogenic role of the Notch pathway in primary liver cancer (Review ...

<https://www.spandidos-publications.com/10.3892/ol.2016.4609> - [翻译此页](#)

作者: J Lu - 2016 - 被引用次数: 9 - 相关文章

2016年5月18日 - The Notch signaling pathway is significantly associated with liver disease (23). ... Thus, Notch receptor expression is abnormal in HCC (27). ... The receptors are transmembrane