

World Journal of *Diabetes*

World J Diabetes 2024 November 15; 15(11): 2157-2271



EDITORIAL

- 2157 Diabetes and obesity: A debate on bariatric interventions and its implications
Tatmatsu-Rocha JC, Lima da Silva MR
- 2162 Diabetes mellitus and comorbidities in elderly people from the Lugu community: A critical-reflective analysis
Tatmatsu-Rocha JC, Gomes-Pinto JC
- 2167 Glucagon-like peptide-1 agonists: Role of the gut in hypoglycemia unawareness, and the rationale in type 1 diabetes
Mirghani HO
- 2173 Targeting neuronal PAS domain protein 2 and KN motif/ankyrin repeat domains 1: Advances in type 2 diabetes therapy
Cheng CH, Hao WR, Cheng TH
- 2177 Macrophages: Key players in diabetic wound healing
Zhou X, Guo YL, Xu C, Wang J

EXPERT CONSENSUS

- 2182 Guidelines and consensus: Jejunoileostomy for diabetes mellitus-surgical norms and expert consensus (2023 version)
Shen JW, Ji CY, Fang XD, Yang B, Zhang T, Li ZC, Li HZ, Liu ZY, Tang J, Liao CW, Lu JZ, Yang X, Zhang XG

REVIEW

- 2189 Autophagy-dependent ferroptosis may play a critical role in early stages of diabetic retinopathy
Sun WJ, An XD, Zhang YH, Tang SS, Sun YT, Kang XM, Jiang LL, Zhao XF, Gao Q, Ji HY, Lian FM

ORIGINAL ARTICLE**Basic Study**

- 2203 Effect of exercise during pregnancy on offspring development through ameliorating high glucose and hypoxia in gestational diabetes mellitus
Tang YB, Wang LS, Wu YH, Zhang LX, Hu LY, Wu Q, Zhou ML, Liang ZX
- 2220 Tiliroside protects against diabetic nephropathy in streptozotocin-induced diabetes rats by attenuating oxidative stress and inflammation
Shang Y, Yan CY, Li H, Liu N, Zhang HF

LETTER TO THE EDITOR

- 2237** Role of intestinal glucagon-like peptide-1 in hypoglycemia response impairment in type 1 diabetes
Cheng CH, Hao WR, Cheng TH
- 2242** Atrial fibrillation and prediabetes: Interplay between left atrium and systemic diseases
Hung MJ
- 2245** Glymphatic system function in diverse glucose metabolism states
Byeon H
- 2251** Relationship between age and subfoveal choroidal thickness and its clinical implications
Sinha S, Nishant P, Morya AK, Singh A
- 2255** Bariatric and endo-bariatric interventions for diabetes: What is the current evidence?
Mondal S, Ambrose Fistus V, Pappachan JM
- 2264** Stem cell exosomes: New hope for recovery from diabetic brain hemorrhage
Cheng CH, Hao WR, Cheng TH

ABOUT COVER

Peer Review of *World Journal of Diabetes*, Dimiter Avtanski, PhD, Director, Endocrine Research Laboratory, Friedman Diabetes Institute, Lenox Hill Hospital, New York, NY 10022, United States. davtanski@northwell.edu

AIMS AND SCOPE

The primary aim of *World Journal of Diabetes* (*WJD*, *World J Diabetes*) is to provide scholars and readers from various fields of diabetes with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJD mainly publishes articles reporting research results and findings obtained in the field of diabetes and covering a wide range of topics including risk factors for diabetes, diabetes complications, experimental diabetes mellitus, type 1 diabetes mellitus, type 2 diabetes mellitus, gestational diabetes, diabetic angiopathies, diabetic cardiomyopathies, diabetic coma, diabetic ketoacidosis, diabetic nephropathies, diabetic neuropathies, Donohue syndrome, fetal macrosomia, and prediabetic state.

INDEXING/ABSTRACTING

The *WJD* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for *WJD* as 4.2; JIF without journal self cites: 4.1; 5-year JIF: 4.2; JIF Rank: 40/186 in endocrinology and metabolism; JIF Quartile: Q1; and 5-year JIF Quartile: Q2.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Yu-Xi Chen*; Production Department Director: *Xu Guo*; Cover Editor: *Jia-Ru Fan*.

NAME OF JOURNAL

World Journal of Diabetes

ISSN

ISSN 1948-9358 (online)

LAUNCH DATE

June 15, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Lu Cai, Md. Shahidul Islam, Michael Horowitz

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9358/editorialboard.htm>

PUBLICATION DATE

November 15, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Relationship between age and subfoveal choroidal thickness and its clinical implications

Sony Sinha, Prateek Nishant, Arvind K Morya, Arshi Singh

Specialty type: Endocrinology and metabolism

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification
Scientific Quality: Grade A, Grade B

Novelty: Grade B, Grade C

Creativity or Innovation: Grade B, Grade C

Scientific Significance: Grade B, Grade B

P-Reviewer: Li YG; Xue F

Received: July 31, 2024

Revised: September 17, 2024

Accepted: September 23, 2024

Published online: November 15, 2024

Processing time: 78 Days and 19.6 Hours



Sony Sinha, Department of Ophthalmology-Vitreo-Retina, Neuro-Ophthalmology and Oculoplasty, All India Institute of Medical Sciences, Patna 801507, Bihar, India

Prateek Nishant, Department of Ophthalmology, ESIC Medical College, Patna 801103, Bihar, India

Arvind K Morya, Department of Ophthalmology, All India Institute of Medical Sciences, Hyderabad 508126, Telangana, India

Arshi Singh, Department of Ophthalmology, Guru Nanak Eye Centre, New Delhi 110001, New Delhi, India

Corresponding author: Arvind K Morya, Doctor, Academic Editor, Department of Ophthalmology, All India Institute of Medical Sciences, Bibi Nagar, Hyderabad 508126, Telangana, India. bulbul.morya@gmail.com

Abstract

The retrospective study by Lei *et al* is an investigation of the relationship between age and subfoveal choroidal thickness (SFCT) in Chinese patients with proliferative diabetic retinopathy. Elements of the study design prevent the generalizability of the study findings, limiting their clinical implications. We recommend consideration of stricter eligibility criteria, other variables like duration of diabetes, interpretation of gender-differences in SFCT, longitudinal follow-up, use of newer choroidal flow indices, comparison of values with normal controls, subgroup analysis to determine the effect of prior treatment, as well as consideration of various real-world scenarios in future studies.

Key Words: Diabetic retinopathy; Optical coherence tomography; Phenotypic sex; Research design; Visual acuity

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The retrospective study by Lei *et al* is an investigation of the relationship between age and subfoveal choroidal thickness (SFCT) in Chinese patients with proliferative diabetic retinopathy. Elements of the study design prevent the generalizability of the study findings, limiting their clinical implications. We recommend consideration of stricter eligibility criteria, other variables like duration of diabetes, interpretation of gender-differences in SFCT, longitudinal follow-up, use of newer choroidal flow indices, comparison of values with normal controls, subgroup analysis to determine the effect of prior treatment, as well as consideration of various real-world scenarios in future studies.

Citation: Sinha S, Nishant P, Morya AK, Singh A. Relationship between age and subfoveal choroidal thickness and its clinical implications. *World J Diabetes* 2024; 15(11): 2251-2254

URL: <https://www.wjgnet.com/1948-9358/full/v15/i11/2251.htm>

DOI: <https://dx.doi.org/10.4239/wjd.v15.i11.2251>

TO THE EDITOR

We read with interest the retrospective study on the non-linear relationship between age and subfoveal choroidal thickness (SFCT) in Chinese patients with proliferative diabetic retinopathy (PDR) by Lei *et al*[1] recently published in the *World Journal of Diabetes*. We commend the authors on their attempt to elucidate the changes in blood dynamics of the choroidal circulation occurring in the clinical setting of PDR[1].

This cross-sectional retrospective study has been conducted in the Chinese population among hospitalized individuals with type 2 diabetes who underwent vitrectomy for PDR-and the contralateral eyes of these individuals were included for the study[1]. We commend the authors for applying appropriate statistical tests such as measures of descriptive data, multivariate linear regression model with spline smoothing for the non-linear relationship thus discovered, and the recursive algorithm to calculate the inflection point. The authors developed a multivariate linear regression model to describe the relationship, which is a technically correct approach to minimize residual confounders. The authors have also indicated that the trend is consistent in all strata except in patients of anemia, proving robustness of their model.

Previously, Xu *et al*[2] have investigated the SFCT in diabetic patients with and without diabetic retinopathy (DR) in the population-based Beijing Eye Study in 2011, including nearly 3500 individuals aged 50-93 years. They concluded that patients with diabetes mellitus (DM) had statistically significantly more SFCT, whereas the presence and stage of DR were not associated additionally with abnormal SFCT[2]. As the present study was undertaken including a comparatively younger patient cohort, its contrasting findings were expected to have important implications for the Chinese population by serving to add on to the findings of the Beijing eye study. However, we believe that there are elements of the study design which prevent the generalizability of the study findings and make them exclusive to the present study, limiting their clinical implications. We would like to highlight these factors which should be taken into account while interpreting the study and utilizing it in clinical practice.

In the present study, the sample included patients who underwent unilateral vitrectomy, and the SFCT was measured in contralateral eyes using extended depth imaging (EDI) on optical coherence tomography (OCT). Considering that the severity of DR is generally symmetrical between the two eyes of the patient, and given that vitrectomy is primarily recommended for severe PDR or its complications like high-risk PDR, non-resolving vitreous haemorrhage or vitreomacular interface (VMI) abnormalities, it is reasonable to argue if the findings of the study align with such advanced PDR cases only[3]. The decrease of SFCT has been found to be significant after the onset of severe DR, and is proportionate to the severity of DR[4]. The authors have not commented if any high-risk characteristics or VMI abnormalities were observed in any subset of the eyes included, and if so, what were the changes in SFCT in these eyes.

It is commendable that the authors took into account and tried to eliminate the possibility of diurnal variation influencing choroidal thickness by obtaining images in the afternoon. It would have been interesting to know the reason for the same, given that the peaks for SFCT and various related indices of choroidal circulation occur between 2:00 and 6:00[5]. Also, it is not known whether these indices remain well-correlated with SFCT in eyes with DR. In line with the authors' recommendations for future studies, OCT angiography could be used to simultaneously measure these choroidal indices to reveal the true changes in choroidal dynamics in PDR in this population[6].

In all the three adjusted models, gender is a common parameter. This has to be viewed in the light of the correlation of SFCT with gender elucidated in available literature, that SFCT is thicker in men than women[7]. The authors found significant difference in the results of stratified analysis between age and SFCT in males but not in females. This finding has not been discussed by the authors and thus, the clinical implications of this finding are presently unclear.

In the present study, yearly reduction in SFCT in PDR patients was found to be 1.68 microns, based on the measurements of SFCT in patients of varying age. It is well-known that SFCT decreases with increasing duration of diabetes. However, the inference that the rate is similar to the average rate of loss of 1.40 to 4.80 microns/year in healthy participants, appears erroneous. Choroidal thickness has been found to decrease in diabetic patients even without evidence of DR[4]. One then wonders why the rate of loss is the same as that in healthy subjects instead of being significantly greater? There is need to longitudinally follow-up individual participants and perform serial EDI-OCT measurements of SFCT to derive the actual rate of SFCT reduction in the subjects. It should also take into account the duration of diabetes in these subjects. This approach is supported by the arguments provided by the authors, on the mechanisms of how hyperglycaemia causes choroidopathy and hence influences choroidal thickness.

The authors concluded that age is a significant determinant in the SFCT, which is in concurrence with previous studies showing significant decrease in SFCT after the age of 60 years[8]. However, caution should be exercised in interpreting the finding that there occurs an inflection point at 54 years regarding trends in SFCT with age. The authors remark that identifying the turning point of age 54 and SFCT in PDR increased statistical power and clinical significance of the study but fail to elaborate as to how they reached this conclusion. It would have been pertinent to perform subgroup analysis amongst treatment naïve patients to elucidate the relationship in question more accurately[9]. It would also be pertinent to compare age related changes in choroidal indices, notably choroidal vascularity index (CVI) and SFCT amongst normal individuals of the same area and ethnicity with those having DM without PDR, treated PDR with and without treated diabetic macular edema (DME), and treatment naïve PDR[10].

Finally, considering that contradictory results observed in studies for SFCT in PDR patients the world over, it is difficult to understand the clinical implications of the findings of the present study. Presently, the evidence does not adequately support the assumption that thinner SFCT-for example in myopia-is associated with any visual acuity changes in otherwise normal patients[11]. However, as *per* recent research, reduced SFCT is associated with poorer gain in visual acuity after vitreoretinal intervention, such as that for epiretinal membrane, and this may be attributable to changes in ocular blood flow[12]. Whether the reduced SFCT results in similarly poor visual acuity gains after intervention for PDR is not fully understood. Dourandeesh *et al*[13] recently concluded an evaluation of changes in SFCT in cases of tractional DME after pars plana vitrectomy, wherein they observed that SFCT decreased and visual acuity increased significantly following the intervention[13]. However, there was no subgroup analysis comparing subjects operated early and those who presented late in their disease course. In addition, there is paucity of evidence regarding the longitudinal change of SFCT after vitreoretinal surgery for PDR[14]. Thus, whether progressive reduction of SFCT is altered by early vitreoretinal surgery for PDR is another future avenue to be explored. Recently, chorioretinal vascularity is being increasingly accepted as a biomarker of predictiveness in vitreomacular surgery as well as anti-vascular endothelial growth factor (VEGF) treatment[15,16]. Whether the newer anti-VEGF agents have some beneficial effects in preventing or reversing alterations of choroidal circulation in PDR in the same way as they do in neovascular age-related macular degeneration, are some of the real-world questions which require further studies.

In conclusion, the present study adds to current literature by providing information on the SFCT in advanced PDR subjects. The results have limited generalizability and unclear clinical implications. Future studies are recommended, involving newer diagnostic tools like OCT angiography, newer indices like CVI, longitudinal comparative study designs, subgroup analyses and comparisons between normal, treated and treatment naïve eyes, as well as SFCT measurements in real-world scenarios.

ACKNOWLEDGEMENTS

We thank the reviewers for their comments and suggestions.

FOOTNOTES

Author contributions: Sinha S and Morya AK designed the research; Nishant P and Singh A conducted literature search; Nishant P drafted the letter; Sinha S and Nishant P drafted the letter; All authors reviewed the final submitted version of the manuscript and agree to be held accountable for all aspects of the manuscript.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: India

ORCID number: Sony Sinha 0000-0002-6133-5977; Arvind K Morya 0000-0003-0462-119X.

S-Editor: Fan M

L-Editor: A

P-Editor: Chen YX

REFERENCES

- 1 Lei CY, Xie JY, Ran QB, Zhang MX. Non-linear relationship between age and subfoveal choroidal thickness in Chinese patients with proliferative diabetic retinopathy. *World J Diabetes* 2024; **15**: 1903-1915 [PMID: 39280183 DOI: 10.4239/wjd.v15.i9.1903]
- 2 Xu J, Xu L, Du KF, Shao L, Chen CX, Zhou JQ, Wang YX, You QS, Jonas JB, Wei WB. Subfoveal choroidal thickness in diabetes and diabetic retinopathy. *Ophthalmology* 2013; **120**: 2023-2028 [PMID: 23697958 DOI: 10.1016/j.ophtha.2013.03.009]

- 3 **Chen SN**, Chen SJ, Wu TT, Wu WC, Yang CH, Yang CM. Refining vitrectomy for proliferative diabetic retinopathy. *Graefes Arch Clin Exp Ophthalmol* 2023; **261**: 3659-3670 [PMID: 37314522 DOI: 10.1007/s00417-023-06134-w]
- 4 **Ambiya V**, Kumar A, Baranwal VK, Kapoor G, Arora A, Kalra N, Sharma J. Change in subfoveal choroidal thickness in diabetes and in various grades of diabetic retinopathy. *Int J Retina Vitreous* 2018; **4**: 34 [PMID: 30214825 DOI: 10.1186/s40942-018-0136-9]
- 5 **Lal B**, Alonso-Caneiro D, Read SA, Carkeet A. Diurnal changes in choroidal optical coherence tomography angiography indices over 24 hours in healthy young adults. *Sci Rep* 2023; **13**: 3551 [PMID: 36864086 DOI: 10.1038/s41598-023-30433-1]
- 6 **Nouri H**, Abtahi SH, Mazloumi M, Samadikhadem S, Arevalo JF, Ahmadi H. Optical coherence tomography angiography in diabetic retinopathy: A major review. *Surv Ophthalmol* 2024; **69**: 558-574 [PMID: 38521424 DOI: 10.1016/j.survophthal.2024.03.004]
- 7 **Wang W**, He M, Zhong X. Sex-Dependent Choroidal Thickness Differences in Healthy Adults: A Study Based on Original and Synthesized Data. *Curr Eye Res* 2018; **43**: 796-803 [PMID: 29451995 DOI: 10.1080/02713683.2018.1428995]
- 8 **Ding X**, Li J, Zeng J, Ma W, Liu R, Li T, Yu S, Tang S. Choroidal thickness in healthy Chinese subjects. *Invest Ophthalmol Vis Sci* 2011; **52**: 9555-9560 [PMID: 22058342 DOI: 10.1167/iovs.11-8076]
- 9 **Wang H**, Tao Y. Choroidal structural changes correlate with severity of diabetic retinopathy in diabetes mellitus. *BMC Ophthalmol* 2019; **19**: 186 [PMID: 31419954 DOI: 10.1186/s12886-019-1189-8]
- 10 **Gupta C**, Tan R, Mishra C, Khandelwal N, Raman R, Kim R, Agrawal R, Sen P. Choroidal structural analysis in eyes with diabetic retinopathy and diabetic macular edema-A novel OCT based imaging biomarker. *PLoS One* 2018; **13**: e0207435 [PMID: 30533048 DOI: 10.1371/journal.pone.0207435]
- 11 **Mahmud M**, Ahem A, Bastion MC, Omar R, Nawi AM, Razak N, Satali AM, Khialdin SM, Din NM. The Relationship Between Choroidal and Photoreceptor Layer Thickness With Visual Acuity in Highly Myopic Eyes. *Front Cell Neurosci* 2022; **16**: 800065 [PMID: 35185474 DOI: 10.3389/fncel.2022.800065]
- 12 **Rommel F**, Brinkmann MP, Sochurek JAM, Prasuhn M, Grisanti S, Ranjbar M. Ocular Blood Flow Changes Impact Visual Acuity Gain after Surgical Treatment for Idiopathic Epiretinal Membrane. *J Clin Med* 2020; **9** [PMID: 32517301 DOI: 10.3390/jcm9061768]
- 13 **Dourandeesh M**, Moeini M, Shaye ZA, Shoeibi N, Hosseini SM, Banaee T. Subfoveal choroidal thickness following pars plana vitrectomy in tractional diabetic macular edema. *Eur J Ophthalmol* 2023; **33**: 1405-1411 [PMID: 36476066 DOI: 10.1177/11206721221144137]
- 14 **Kim DI**, Bae KW, Hwang DD. Longitudinal Change of Choroidal Thickness after Pars Plana Vitrectomy for Idiopathic Epiretinal Membrane. *J Clin Med* 2022; **11** [PMID: 36498525 DOI: 10.3390/jcm11236950]
- 15 **D'Aloisio R**, Ruggeri ML, Porreca A, Di Nicola M, Aharrh-Gnama A, Quarta A, Gironi M, Toto L, Mastropasqua R. Choroidal Vascularity Index Fluctuations in Epiretinal Membranes in Vitreoretinal Surgery: Comparison Between Idiopathic and Diabetic Ones. *Transl Vis Sci Technol* 2023; **12**: 9 [PMID: 38060233 DOI: 10.1167/tvst.12.12.9]
- 16 **Minnella AM**, Centini C, Gambini G, Savastano MC, Pagliei V, Falsini B, Rizzo S, Ciasca G, Maceroni M. Choroidal Thickness Changes After Intravitreal Aflibercept Injections in Treatment-Naïve Neovascular AMD. *Adv Ther* 2022; **39**: 3248-3261 [PMID: 35597837 DOI: 10.1007/s12325-022-02129-x]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

