

World Journal of *Clinical Cases*

World J Clin Cases 2019 July 6; 7(13): 1535-1731



REVIEW

- 1535 Intracranial pressure monitoring: Gold standard and recent innovations
Nag DS, Sahu S, Swain A, Kant S
- 1554 Role of the brain-gut axis in gastrointestinal cancer
Di YZ, Han BS, Di JM, Liu WY, Tang Q

MINIREVIEWS

- 1571 Cholestatic liver diseases: An era of emerging therapies
Samant H, Manatsathit W, Dies D, Shokouh-Amiri H, Zibari G, Boktor M, Alexander JS

ORIGINAL ARTICLE**Basic Study**

- 1582 Neural metabolic activity in idiopathic tinnitus patients after repetitive transcranial magnetic stimulation
Kan Y, Wang W, Zhang SX, Ma H, Wang ZC, Yang JG

Retrospective Study

- 1591 Neuroendoscopic and microscopic transsphenoidal approach for resection of nonfunctional pituitary adenomas
Ding ZQ, Zhang SF, Wang QH
- 1599 Safety and efficacy of transjugular intrahepatic portosystemic shunt combined with palliative treatment in patients with hepatocellular carcinoma
Luo SH, Chu JG, Huang H, Yao KC
- 1611 Leveraging machine learning techniques for predicting pancreatic neuroendocrine tumor grades using biochemical and tumor markers
Zhou RQ, Ji HC, Liu Q, Zhu CY, Liu R

Clinical Trials Study

- 1623 Development and validation of a model to determine risk of refractory benign esophageal strictures
Lu Q, Lei TT, Wang YL, Yan HL, Lin B, Zhu LL, Ma HS, Yang JL

SYSTEMATIC REVIEWS

- 1634 Hydatid cyst of the colon: A systematic review of the literature
Lataú-Córdoba MÁ, Ruiz-Blanco S, Sanchez M, Santiago-Boyero C, Soto-García P, Sun W, Ramia JM

CASE REPORT

- 1643 Colon cancer arising from colonic diverticulum: A case report
Kayano H, Ueda Y, Machida T, Hiraiwa S, Zakoji H, Tajiri T, Mukai M, Nomura E
- 1652 Endoscopic submucosal dissection as excisional biopsy for anorectal malignant melanoma: A case report
Manabe S, Boku Y, Takeda M, Usui F, Hirata I, Takahashi S
- 1660 Gastrointestinal infection-related disseminated intravascular coagulation mimicking Shiga toxin-mediated hemolytic uremic syndrome - implications of classical clinical indexes in making the diagnosis: A case report and literature review
Li XY, Mai YF, Huang J, Pai P
- 1671 A complicated case of innominate and right common arterial aneurysms due to Takayasu's arteritis
Wang WD, Sun R, Zhou MX, Liu XR, Zheng YH, Chen YX
- 1677 Multimodality-imaging manifestations of primary renal-allograft synovial sarcoma: First case report and literature review
Xu RF, He EH, Yi ZX, Lin J, Zhang YN, Qian LX
- 1686 Chronic progression of recurrent orthokeratinized odontogenic cyst into squamous cell carcinoma: A case report
Wu RY, Shao Z, Wu TF
- 1696 Gastric adenocarcinoma of fundic gland type after *Helicobacter pylori* eradication: A case report
Yu YN, Yin XY, Sun Q, Liu H, Zhang Q, Chen YQ, Zhao QX, Tian ZB
- 1703 Synchronous multiple primary gastrointestinal cancers with *CDH1* mutations: A case report
Hu MN, Lv W, Hu RY, Si YF, Lu XW, Deng YJ, Deng H
- 1711 Primary hepatoid adenocarcinoma of the lung in Yungui Plateau, China: A case report
Shi YF, Lu JG, Yang QM, Duan J, Lei YM, Zhao W, Liu YQ
- 1717 Liver failure associated with benzbromarone: A case report and review of the literature
Zhang MY, Niu JQ, Wen XY, Jin QL
- 1726 Giant low-grade appendiceal mucinous neoplasm: A case report
Yang JM, Zhang WH, Yang DD, Jiang H, Yu L, Gao F

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Hitoshi Hirose, MD, PhD, Associate Professor, Department of Cardiothoracic Surgery, Thomas Jefferson University, Philadelphia, PA 19107, United States

AIMS AND SCOPE

World Journal of Clinical Cases (*World J Clin Cases*, *WJCC*, online ISSN 2307-8960, DOI: 10.12998) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

The primary task of *WJCC* is to rapidly publish high-quality Case Report, Clinical Management, Editorial, Field of Vision, Frontier, Medical Ethics, Original Articles, Meta-Analysis, Minireviews, and Review, in the fields of allergy, anesthesiology, cardiac medicine, clinical genetics, clinical neurology, critical care, dentistry, dermatology, emergency medicine, endocrinology, family medicine, gastroenterology and hepatology, etc.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in PubMed, PubMed Central, Science Citation Index Expanded (also known as SciSearch®), and Journal Citation Reports/Science Edition. The 2019 Edition of Journal Citation Reports cites the 2018 impact factor for *WJCC* as 1.153 (5-year impact factor: N/A), ranking *WJCC* as 99 among 160 journals in Medicine, General and Internal (quartile in category Q3).

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: *Yan-Xia Xing*
 Proofing Production Department Director: *Yun-Xiaojuan Wu*

NAME OF JOURNAL
World Journal of Clinical Cases

ISSN
 ISSN 2307-8960 (online)

LAUNCH DATE
 April 16, 2013

FREQUENCY
 Semimonthly

EDITORS-IN-CHIEF
 Dennis A Bloomfield, Sandro Vento

EDITORIAL BOARD MEMBERS
<https://www.wjnet.com/2307-8960/editorialboard.htm>

EDITORIAL OFFICE
 Jin-Lei Wang, Director

PUBLICATION DATE
 July 6, 2019

COPYRIGHT
 © 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS
<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS
<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT
<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE
<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS
<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION
<https://www.f6publishing.com>

Clinical Trials Study

Development and validation of a model to determine risk of refractory benign esophageal strictures

Qing Lu, Tian-Tian Lei, Yi-Lan Wang, Hai-Lin Yan, Bo Lin, Lin-Lin Zhu, Hong-Sheng Ma, Jin-Lin Yang

ORCID number: Qing Lu (0000-0001-9107-1311); Tian-Tian Lei (0000-0001-9884-185X); Yi-Lan Wang (0000-0002-9689-4883); Hai-Lin Yan (0000-0002-1997-2454); Bo Lin (0000-0003-0696-7717); Lin-Lin Zhu (0000-0002-4740-2838); Hong-Sheng Ma (0000-0002-2292-6387); Jin-Lin Yang (0000-0001-8726-7258).

Author contributions: Lu Q and Lei TT contributed equally to this work; Yang JL and Ma HS designed the research; Lu Q, Yan HL, Wang YL, Lei TT, and Zhu LL performed the research; Lu Q, Wang YL, and Lei TT performed the follow-ups; Lu Q analyzed the data and wrote the paper; all authors read and approved the final manuscript.

Institutional review board

statement: This study was reviewed and approved by the Institutional Review Board of the West China Hospital of Sichuan University.

Informed consent statement:

Patients were not required to give informed consent to the study because the analysis used anonymous clinical data that were obtained after each patient agreed to treatment by written consent.

Conflict-of-interest statement: All authors declare no conflicts of interest.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non

Qing Lu, Yi-Lan Wang, Hai-Lin Yan, Bo Lin, Lin-Lin Zhu, Jin-Lin Yang, Department of Gastroenterology, West China Hospital of Sichuan University, Chengdu 610041, Sichuan Province, China

Tian-Tian Lei, Hong-Sheng Ma, Department of Day Surgery, West China Hospital of Sichuan University, Chengdu 610041, Sichuan Province, China

Corresponding author: Jin-Lin Yang, MD, MHSc, PhD, Chief Doctor, Doctor, Lecturer, MHSc, Professor, Department of Gastroenterology, West China Hospital of Sichuan University, No. 37, Guoxue Road, Chengdu 610000, Sichuan Province, China.
mouse-577@163.com

Telephone: +86-18-980602058

Fax: +86-28-85423387

Abstract**BACKGROUND**

Current research has identified several risk factors for refractory benign esophageal strictures (RBES), but research is scarce on the prediction of RBES in benign esophageal strictures patients. Meanwhile, the long-term outcomes of RBES remain unclear. The aim of this study was to develop and validate a model to determine the progression of RBES in patients with benign esophageal strictures. And we also explored the long-term outcomes and safety in patients with RBES.

AIM

To develop and validate a model to determine the progression of RBES in patients with benign esophageal strictures, based on the demographic data and endoscopic findings.

METHODS

A total of 507 benign esophageal stricture patients treated by dilation alone or in combination with stenting were retrospectively enrolled between January 2009 and February 2018. The primary outcome was to establish a risk-scoring model predicting RBES in benign esophageal strictures. The secondary outcome was to explore the clinical effectiveness and adverse events in patients with RBES.

RESULTS

In the study, age, etiology, and number and length of strictures were the independent risk factors for the refractory performance of benign esophageal strictures. According to risk factors of benign esophageal strictures, a risk-scoring

Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: February 27, 2019

Peer-review started: February 27, 2019

First decision: March 27, 2019

Revised: April 17, 2019

Accepted: May 2, 2019

Article in press: May 3, 2019

Published online: July 6, 2019

P-Reviewer: Jarcuska P, Konishi H, Lima M

S-Editor: Ji FF

L-Editor: Wang TQ

E-Editor: Xing YX



model for predicting RBES in benign esophageal strictures was established: The risk score ranged from 0 to 8 points, and the risk scores were divided into low risk (0-2 points), intermediate risk (3-5 points), and high risk (6-8 points). The proportions of RBES in the corresponding risk categories were 1.0%, 12.2%, and 76.0%, respectively. Among 507 patients, 57 had RBES (39 males; median age, 60 years). The success rate of dilation treatment (51.2%, 21/41) was higher than that of stent placement (37.5%, 6/16).

CONCLUSION

In this study, 11.3% (57/507) patients had RBES at our hospital. The risk-scoring model predicting RBES in benign esophageal strictures could predict the long-term outcome of patients with strictures ahead.

Key words: Refractory benign esophageal strictures; Esophageal dilation; Esophageal stents; Long-term outcomes; Retrospective analysis

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: We developed and validated a model to determine the progression of refractory benign esophageal strictures in patients with benign esophageal strictures. The risk-scoring model could predict the long-term outcome of patients with strictures ahead. The endoscopic therapy was the standard treatment for all benign stenosis patients. Our findings may have an impact on daily practice. The presence of a bad score may mean surgical indication and refractoriness to endoscopic therapy.

Citation: Lu Q, Lei TT, Wang YL, Yan HL, Lin B, Zhu LL, Ma HS, Yang JL. Development and validation of a model to determine risk of refractory benign esophageal strictures. *World J Clin Cases* 2019; 7(13): 1623-1633

URL: <https://www.wjgnet.com/2307-8960/full/v7/i13/1623.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v7.i13.1623>

INTRODUCTION

Benign strictures of the esophagus include peptic esophageal strictures, caustic esophageal strictures, post-surgical esophageal strictures, post-endoscopic submucosal dissection (ESD) esophageal strictures, esophageal strictures caused by local radiotherapy and tuberculosis, and others^[1,2]. The goals of therapy for benign esophageal strictures are relieving dysphagia and the prevention of stricture recurrence^[3]. Dilation with bougie or balloon is the classic treatment. In the majority of patients, this can be accomplished with esophageal dilation, but in cases of refractoriness to dilation, additional therapy may be required during long-term follow-up^[4,5]. Kochman *et al*^[6] defined refractory benign esophageal strictures (RBES) as more than 3-5 dilations having been performed without clinical response or when it is impossible to achieve a 14 mm lumen over 3-5 dilation sessions^[6,7]. Current research has identified several risk factors for RBES, but research is scarce on the prediction of RBES in benign esophageal strictures patients. Meanwhile, the long-term outcomes of RBES remain unclear. The aim of this study was to develop and validate a model to determine the progression of RBES in patients with benign esophageal strictures, based on the demographic data and endoscopic findings. And we also explored the long-term outcomes and safety in patients with RBES.

MATERIALS AND METHODS

Patients

This is a single-center retrospective study. A total of 1146 patients with esophageal strictures were treated by dilation alone or in combination with stenting between January 2009 and February 2018 at our hospital. All medical records were retrospectively reviewed. Overall, 507 patients were included in the study. The inclusion criteria were: (1) They had been diagnosed with esophageal benign stricture according to the clinical manifestation, gastroscopy, and pathology; (2) They had complete

medical records and follow-up information, including gender, age at first therapy, etiology, number, location, length and diameter of structures, endoscopic therapy, and adverse events; and (3) They were more than 16 years old. Patients were excluded if they: (1) Had been diagnosed with congenital esophageal strictures, malignant esophageal strictures, or esophageal fistula; (2) Were diagnosed with recurrence of esophageal cancer; (3) Were lost to follow-up, or had incomplete data; (4) Were less than 16 years old; and (5) Had a history of endoscopic incision after stenosis. The inclusion criteria for RBES were: (1) Satisfying the inclusion criteria for benign esophageal stenosis; and (2) Esophageal endoscopic treatment (including esophageal dilation treatment and esophageal stent placement) performed five times or more, and benign esophageal strictures remained with dysphagia or it did not achieve a 14 mm lumen (Figure 1). This study was approved by the Ethics Committee of West China Hospital of Sichuan University.

Equipment and procedure

Gastrosopes (JIF - Q260 or JIF - Q260J, Olympus Medical Systems, Tokyo, Japan) were used according to the stricture characteristics and type of treatment. Each physician had over 10 years of experience in endoscopy. Written consent for endoscopic dilation and stenting was obtained before the operations. All patients received the standard dilation, with balloon and bougie or fully covered self-expandable metal stents (SEMSs) (MTN-SE C-membrane; 60-140 mm; 20 mm; MicroTech, Nanjing, China). The dilation procedures were performed with a through-the-scope balloon (Type 5842; 3 ATM-12 mm, 4.5 ATM-13.5 mm, 8 ATM-15 mm; Boston Scientific Corp; Marlborough, United States) and Savary bougie system (Savary-Gilliard Dilator set, type SGD-70-1; length: 70 cm, diameters: 5, 7, 9, 11, 12.8, 14, and 15 mm; Wilson-Cook Medical, United States). Dilation was the standard treatment for all patients, with other alternatives, such as stents placed on a case-by-case basis at the discretion of the endoscopist performing the procedure, after an appropriate discussion with surgeons and a dietician, as needed. In our study, the stent removal was performed 6-8 wk after insertion to allow the remodeling of the scar tissue.

Endpoints

The primary outcome was to establish a risk-scoring model predicting RBES in patients having benign esophageal strictures. The secondary outcome was to explore the endoscopic therapy effectiveness, ineffectiveness, and adverse events in patients with RBES. Endoscopic therapy success was defined as no need for any other interventions, lasting at least 6 mo^[8]. Treatment failure was defined as the need for any endoscopic intervention, including surgery^[9], during the follow-up period. Adverse events included esophageal fistula, bleeding, stent dysfunction (stent overgrowth, stent migration, and food impaction), and intolerable chest pain. Early and late adverse events were defined as those occurring within 7 d and later than 7 d after stenting, respectively. All patients were followed via telephone contact, from the first therapy session until study termination or patient death to assess symptom resolution.

Statistical analysis

Descriptive statistics are reported as the median or mean and range for continuous variables. The χ^2 test and the Fisher's exact test were used for categorical variables, and the *t*-test was used for continuous variables. To create and validate a risk model for the progression of RBES in benign esophageal strictures patients, we used a split-sample strategy. Using a random number from the uniform (0, 1) distribution, we labeled those with values < 0.3 as a validation cohort and > 0.3 as a development cohort, and this gave us a 70/30 split. This strategy has been used in multiple models^[10,11]. In the development cohort (70%), we tested candidate variables and retained the statistical significance in multivariate binary logistic regression analysis. In addition, multivariate binary logistic regression analysis, with the corresponding odds ratios (OR) and 95% confidence intervals (CI) reported, was performed to identify the independent risk factors for RBES. A scoring model was established to calculate the probabilities of RBES in benign esophageal strictures patients on the basis of the retaining candidate variables. The independent risk factors were chosen as scoring items. Meanwhile, each risk factor score was weighted based on the beta coefficient obtained from the logistic regression analysis. In order to keep the scoring model simple, the respective beta coefficient was rounded to the nearest whole number. The total score represented the sum of scores for each risk factor. Using these scores, we derived three groups of risk: Low (points 0-2), intermediate (3-5), and high (6-8). Probabilities of RBES with 95% CI in each group, stratified by the risk factor, were calculated. In the validation cohort (30%), the model discrimination was tested




Figure 1 Flowchart of selection of patients with benign esophageal strictures.

with the receiver operating characteristic (ROC) curve and area under the ROC curve (AUROC). An AUROC of 1.0 indicated perfect concordance, whereas an AUROC of 0.5 indicated no relationship. External validation of the scoring model was performed by measuring the discriminatory ability with AUROC. To test calibration, we used a Hosmer-Lemeshow (H-L) test along with reporting the calibration slope and intercept. To evaluate the relationship between various clinical characteristics in RBES patients, univariate analyses were performed using the χ^2 test and *t*-test. Statistical analyses were performed using SPSS Statistics version 22.0 (IBM Corp, Armonk, NY, United States). *P*-values less than 0.05 were considered statistically significant.

RESULTS

Baseline characteristics

Patient characteristics are shown in [Table 1](#). In total, 507 patients with benign esophageal strictures were included (178 males; median age, 58 years). Among these patients, esophageal strictures were caused by esophagectomy in 246 (48.5%) patients, ESD in 78 (15.4%), achalasia in 148 (29.2%), caustic injury in 13 (2.6%), mixed etiology in 2 (0.4%), and others in 20 (3.9%). Among the 507 cases, esophageal strictures were located in the cervical esophagus in 40 (7.9%), thoracic esophagus in 315 (62.1%), and ventral esophagus in 152 (30.0%). A single stricture was observed in 477 (94.1%) patients, 24 (4.7%) patients had two strictures, and 6 (1.2%) patients had three or more strictures. The median stricture length was 0.5 cm (range, 0.5-10.0 cm). All 507 patients received endoscopic dilations. A total of 299 (59.0%) patients underwent one dilation session, 85 (16.8%) underwent 2, 42 (8.3%) underwent 3, 24 (4.7%) underwent 4, and 57 (11.2%) had RBES. The median number of dilation session was 5 (range, 3-21) during a mean follow-up period of 31 months (range, 6-106 mo). The SEMSs were placed in 55 (10.8%) patients. A mean of 1.9 (range, 1-6) SEMSs were placed in 20 (26.7%) patients. The mean time of stent placement was 29 d (range, 7-67 d).

Univariate and multivariate analyses of RBES

The clinical features of the 507 benign esophageal strictures are summarized in [Table 2](#). The incidence of RBES was 11.2% (57/507) in benign esophageal strictures. Significant associations between the incidence of RBES and etiology, location, number and length of strictures, and endoscopic therapy were revealed in the univariate analyses ([Table 2](#)). There was a significant association between the incidence of RBES and age, etiology, number and length of strictures, and endoscopic therapy in the multivariate analyses ([Table 2](#)).

Development and validation of a risk-scoring model to predict RBES

We initially included 7 candidate predictors ([Tables 2 and 3](#)) in our analysis.

Table 1 Characteristics of 507 patients with benign esophageal strictures

Category		N (%) / M (range) n = 507
Sex	Female	178 (35.1)
	Male	329 (64.9)
Age (yr)	-	58 (16-88)
Etiology ¹	Post-ESD/ESTD	78 (15.4)
	Achalasia	148 (29.2)
	Caustic	13 (2.6)
	Postoperative	246 (48.5)
	Peptic	3 (0.6)
	Trauma	2 (0.4)
	Mixed	2 (0.4)
	Post-radiation	1 (0.2)
	Iatrogenic	14 (2.8)
	Location of strictures ¹	Cervical
Thoracic		315 (62.1)
Ventral		152 (30.0)
Number of strictures	One	477 (94.1)
	Two	24 (4.7)
	Three or more	6 (1.2)
Length of strictures (cm)	-	0.5 (0.5, 10)
Times	One	299 (59.0)
	Two	85 (16.8)
	Three	42 (8.3)
	Four	24 (4.7)
	Five or more	57 (11.2)
Endoscopic therapy	Stenting + dilation	55 (10.8)
	Dilation only	452 (89.2)

¹Fisher's exact test. The mixed etiology was the patients with refractory benign esophageal strictures caused by caustic injury and esophageal anastomosis following colon surgery. ESD: Endoscopic submucosal dissection; ESTD: Endoscopic submucosal tunnel dissection.

Multivariate binary logistic regression analysis revealed that age, etiology, number, length of strictures, and endoscopic therapy were the independent risk factors for RBES. In reality, stenting was one of the treatment approaches for strictures after multiple dilations with bougie or balloon. We noticed that the stent placement was mostly chosen in patients with RBES, so we excluded endoscopic therapy from candidate predictors. Ultimately, the final model contained only four variables: Age, etiology, and number and length of strictures. Beta coefficients of the four independent risk factors identified in the multivariate analysis are shown in Table 4, and the scores of selected predictors were assigned an integer score based on the Beta coefficient. Based on the scores in each predictor for the development dataset of 345 cases (age, etiology, and number and length of strictures), an 8-point risk-scoring model was built to predict the stratified RBES risk (Table 5). With the scoring model, the percentage of RBES ranged from 0.0 to 100% in the order of calculated risk scores in the development cohort (Table 5). The risk trend of low to high scores presented a significant linear increase ($P < 0.001$, Cochran-Armitage trend test). A ROC curve of the risk-scoring model for RBES was plotted using the development cohort, and the AUROC of the model for the prediction of RBES was 0.858 (Figure 2). The percentage of RBES for each calculated score and the AUROC were also examined in the validation dataset of 162 cases (Table 5 and Figure 2). The percentage increased from 0.0% to 25% as the score increased, and the AUROC was 0.768. H-L test showed that the P -value was 0.813 in the validation dataset. We divided the cohorts into three risk groups: low, intermediate, and high-risk groups based on point scoring system and the probability of progression to RBES (Table 5). The total risk of progression, at the end of study period, of RBES in each of the three risk categories is provided in Table 5: Low risk: 1.0%, intermediate risk: 12.2%, and high risk: 76.0%.

Table 2 Univariate and multivariate analyses of refractory benign esophageal strictures

Category		Non-RBES n (%) / M (range) n = 450	RBES n (%) / M (range) n = 57	P ¹	P ²	OR (95%CI)
Sex	Female	160 (89.9)	18 (10.1)	0.553	0.665	0.916 (0.617-1.361)
	Male	290 (88.1)	39 (11.9)			
Age (yr)	-	58 (16, 88)	60 (16, 75)	0.548	0.034	0.968 (0.939-0.998)
Etiology	Post-ESD/ESTD	55 (70.5)	23 (29.5)	< 0.001	0.007	5.874 (1.623-21.255)
	Postoperative	223 (90.7)	23 (9.3)			
	Achalasia	147 (99.3)	1 (0.3)			
	Caustic	5 (38.5)	8 (61.5)			
	Other	20 (90.9)	2 (9.1)			
Location of strictures	Cervical	30 (75.0)	10 (25.0)	< 0.001	0.603	1.250 (0.539-2.901)
	Thoracic	271 (86.0)	44 (14.0)			
	Ventral	149 (98.0)	3 (2.0)			
Length of strictures (cm)		0.5 (0.5, 8)	2 (0.5, 10)	< 0.001	0.001	1.344 (1.135-1.591)
Number of strictures	One	436 (91.4)	41 (8.6)	< 0.001	0.049	0.615 (0.370-1.023)
	Two or more	14 (46.7)	16 (53.3)			
Endoscopic therapy	Dilation only	411 (90.9)	41 (9.1)	< 0.001	0.009	0.582 (0.388-0.874)
	Stenting + dilation	39 (70.9)	16 (29.1)			

RBES: Refractory benign esophageal strictures; ESD: Endoscopic submucosal dissection; ESTD: Endoscopic submucosal tunnel dissection.

Clinical outcomes and adverse events in 57 patients with RBES

Endoscopic therapy success was achieved in 27 (47.4%) patients. The success rate was higher in patients treated by dilation (77.8%, 21/27) than in those treated with stents (22.2%, 6/27). Endoscopic treatment failure was observed in 30 (52.6%) patients. One patient was treated by surgery, and another patient was treated with endoscopic gastrostomy feeding tubes. Fourteen patients had adverse events after endoscopic therapy. Six patients experienced esophageal fistula. One patient experienced intolerable chest pain, and one patient experienced bleeding. Ten patients experienced stent dysfunction, including stent overgrowth in 2, stent migration in 5, and food impaction in 3 (Table 6). There was a significant relationship between endoscopic treatment success and etiology ($P = 0.003$) and endoscopic therapy ($P = 0.043$) in univariate analysis (Table 7). Comparing bougie to balloon dilation showed no difference in terms of clinical resolution ($P = 0.949$).

DISCUSSION

A total of 507 patients with benign esophageal strictures met the inclusion criteria. Surgery was the main cause of stenosis, accounting for 48.5% (246/507). Multivariate analysis suggested that post-ESD/endoscopic submucosal tunnel dissection strictures, post-surgery strictures, multiple esophageal strictures, and long segment strictures may lead to RBES. Our results are consistent with the previous research results. Lew *et al* considered that the short, straight, and not angulated strictures could positively respond to endoscopic dilation as simple strictures. And those complex strictures that are angulated, long (>2 cm), and irregular were often refractory to dilation. Another study showed that longer strictures had a high risk of being refractory to treatment^[12]. Patients undergoing esophageal surgery, especially esophagectomies, may develop stenosis, due to developing local ischemia. Patients usually require three to five sessions for the resolution of dysphagia. Esophageal strictures are one of the common complications of post-ESD strictures. The majority of the patients needed to experience several dilations to achieve the relief of symptoms, especially those refractory to dilation, and could accept dilation every around 15 d in our study.

Patients with benign esophageal strictures were stratified according to the risk factors. In our study, we considered that caustic strictures were the high-risk predictive factor, and the finding is consistent with the previous research results. Caustic strictures frequently are a difficult type of strictures associated with a greater




Figure 2 The receiver operating characteristic curve of the risk-scoring model for refractory benign esophageal strictures. ROC: Receiver operating characteristic.

need for subsequent dilatations, due to the corrosion resulting in widespread esophageal mucosa damage, and a long-lasting local inflammatory process^[13]. Because the mucosal injury often involves an extended segment of the esophagus, the destructed structure frequently is complex, with angulations, narrow lumens, greater length, and severe fibrosis than simple strictures^[14]. Caustic strictures often require multiple dilatations and are treated for prolonged periods of time. The caustic strictures can be extremely refractory despite repeated dilatations, and an esophagectomy may be required to alleviate dysphagia. But the alternatives of surgical reconstruction or feeding tube frequently are unsatisfactory. In our analysis, most refractory caustic strictures had a clinical resolution.

We built the RBES risk assessment model, and the total risk score ranged from 0 to 8 points, based on which the risk was divided into low (0-2), medium (3-5 points), and high (6-8). The refractory benign esophageal stenosis rates in the low, medium, and categories were 1.0%, 12.2%, and 76.0%, respectively, and the low-risk to high-risk population trend increased linearly and significantly. According to the risk assessment model, we could judge the long-term results of stenosis patients in advance. The study examined a large number of subjects and revealed detailed data on RBES risk stratified by risk factors. RBES risk differed considerably according to the status of the risk factors. These results could be useful in determining the management of patients with benign esophageal strictures. For the high risk RBES, it is desirable to perform gastrectomy and choose other interventions, such as surgery.

Our study patients were long-term followed and evaluated after at least 5 therapy sessions, to assess the long-term outcomes, safety, and therapeutic success of endoscopic therapy in patients with RBES. We defined the endoscopic therapy success as no need for any other interventions, lasting at least 6 months. The endoscopic therapy success had an inconsistent standard. So, in our study, the dysphagia-free period was a proxy for survival quality. Once over 6 months, it indicated good quality of life.

In most studies, apparently, the most promising alternative endoscopic stenting can affect the long-term endoscopic therapy outcome. In fact, in our study, the rate of successful outcome appeared to be worse in the subgroup with stents as compared with those without. So, we observed that the stent therapy did not reduce the frequency of dilation. Immediate and postponed complications after dilation occur in 0.5% to 1.2%^[15], and excess bleeding may occur in less than 0.5% of the cases^[15].

This study has some limitations. First, our study had a potential selection bias, which inherently decreased the statistical power of the study. Second, our study was a retrospective single-center design. A large proportion of our patients were referred, so the incidence of RBES may be overestimated. To best identify the factors that predict RBES and endoscopic therapy efficacy, adverse events, and dysphagia-free period, a large, prospective, and randomized controlled study is needed. Third, although external validation of the scoring model was performed in this study by examining more benign esophageal stricture cases at the same institution with the development set, more external validation in other institutions is still necessary for the future studies.

Overall, 11.3% (57/507) of patients were with RBES in our hospital. The 8-point risk-scoring model predicted RBES in benign esophageal strictures based on age,

Table 3 Characteristics for included patients in development and validation cohorts

Category	Development cohort (n = 345)	Validation cohort (n = 162)	All (n = 507)
Sex			
Female	119	59	178
Male	226	103	329
Age (yr)			
50 ≤ n < 70	197	83	280
<50	103	55	158
≥70	45	24	69
Etiology			
Achalasia	103	45	148
Post-ESD/ESTD	58	20	78
Postoperative	170	76	246
Caustic	6	7	13
Other	8	14	22
Location of strictures			
Cervical	26	14	40
Thoracic	208	107	315
Ventral	111	41	152
Length of strictures (cm)			
<2 cm	282	135	417
≥2 cm	63	27	90
Number of strictures			
One	325	152	477
Two or more	20	10	30

ESD: Endoscopic submucosal dissection; ESTD: Endoscopic submucosal tunnel dissection.

etiology, and number and length of strictures. The risk scores were divided into low (0-2 points), intermediate (3-5 points), and high risk (6-8 points). And the proportions of RBES in the corresponding risk categories were 1.0%, 12.2%, and 76.0%, respectively. Meanwhile, it could predict the long-term outcome of patients with strictures ahead and might enable more individualized care for patients with benign esophageal strictures.

Table 4 Development of a 9-point risk-scoring model to predict refractory benign esophageal strictures

Factor	P	OR	Beta coefficient	Score
Age (yr)				
50 ≤ n < 70	0.066	1 (Ref.)	0	0
<50	0.021	1.2-8.6	1.2	1
≥70	0.776	0.4-3.2	0.1	0
Etiology				
Achalasia	0.037	1 (Ref.)	0	0
Post-ESD/ESTD	0.004	25.8 (2.9-230.5)	3.3	3
Caustic	0.009	52.3 (2.7-1027.8)	4.0	4
Postoperative	0.006	19.2 (2.3-161.2)	3.0	3
Other	0.216	6.7 (0.3-136.6)	1.9	2
Number of strictures				
One	0.042	1 (Ref.)	0	0
Two or more	0.042	3.5 (1.1-11.5)	1.2	1
Length of strictures (cm)				
< 2 cm	< 0.001	1 (Ref.)	0	0
≥ 2 cm	< 0.001	6.4 (2.7-15.2)	1.9	2

RBES: Refractory benign esophageal strictures; ESD: Endoscopic submucosal dissection; ESTD: Endoscopic submucosal tunnel dissection.

Table 5 Risk score and proportion of refractory benign esophageal strictures in esophageal strictures

Risk score	Rate of RBES (n = 345)	Rate of RBES (n = 162)	Risk	Point	Development cohort Rate of RBES	Validation cohort Rate of RBES
0	2.6% (1/38)	0% (0/11)	Low	0-2	1.0% (1/107)	0% (0/43)
1	0% (0/67)	0% (0/28)	Intermediate	3-5	12.2% (26/213)	7.3% (8/110)
2	0% (0/2)	0% (0/4)	High	6-8	76.0% (19/25)	33.3% (3/9)
3	6.7% (10/149)	6% (5/84)				
4	14.3% (4/28)	0% (0/8)				
5	33.3% (12/36)	16.7% (3/18)				
6	73.7% (14/19)	14.3% (1/7)				
7	100% (3/3)	100% (1/1)				
8	66.7% (2/3)	100% (1/1)				

RBES: Refractory benign esophageal strictures.

Table 6 Adverse events in patients with refractory benign esophageal strictures, n (%)

	Early, n = 3	Late, n = 11
Stent overgrowth	0	2
Stent migration	0	5
Food impaction	0	3
Fistula	1	5
Bleeding	1	0
Intolerable chest pain	1	0

Table 7 Characteristics of 57 patients with refractory benign esophageal strictures

Category	N (%) / M (range)	Success (n = 27)	Failure (n = 30)	P
Sex				
Female	18 (31.5%)	9	9	0.787
Male	39 (68.5%)	18	21	
Age (yr)	60 (16-75)	58 (16-75)	60.5 (16-74)	0.472
Etiology				
Post-ESD/ESTD	23 (40.4%)	11	12	0.003
Postoperative	23 (40.4%)	7	16	
Caustic	8 (14%)	8	0	
Other*	3 (5.2%)	1	2	
Location of strictures				
Cervical	10 (17.5%)	5	5	0.135
Thoracic	44 (77.2%)	22	22	
Ventral	3 (5.3%)	0	3	
Number of strictures				
1	41 (71.9%)	16	25	0.351
2	11 (19.3%)	7	4	
3 and more	5 (8.8%)	4	1	
Length of strictures (cm)	2 (0.5-10)	3 (0.5-10)	2 (0.5-10)	0.105
Diameter of strictures (mm)	4 (1-8.5)	4 (1-8.5)	4 (0.5-8.5)	0.240
Times	7 (5-28)	6 (5-19)	7.5 (5-28)	0.129
Endoscopic therapy				
Stenting + dilation	16 (28%)	6	10	0.043
Dilation only	41 (72%)	21	20	

RBES: Refractory benign esophageal strictures; ESD: Endoscopic submucosal dissection; ESTD: Endoscopic submucosal tunnel dissection.

ARTICLE HIGHLIGHTS

Research background

Current research has identified several risk factors for refractory benign esophageal strictures (RBES), but research is scarce on the prediction of RBES in benign esophageal strictures patients. Meanwhile, the long-term outcomes of RBES remain unclear. The aim of this study was to develop and validate a model to determine the progression of RBES in patients with benign esophageal strictures. And we also explored the long-term outcomes and safety in patients with RBES.

Research motivation

To our knowledge, the present study is the largest sample research of benign esophageal strictures, and our observation indicators are complete, the follow-up period is long, the results are credible, and there is strong reference significance in clinical work.

Research objectives

The aim of this study was to develop and validate a model to determine the progression of RBES in patients with benign esophageal strictures.

Research methods

A total of 507 benign esophageal stricture patients treated by dilation alone or in combination with stenting were retrospectively enrolled.

Research results

Among 507 patients, 57 were with RBES (39 males; median age, 60 years). The age, etiology, and number and length of strictures were the independent risk factors for the refractory performance of benign esophageal strictures. According to risk factors of benign esophageal strictures, a risk-scoring model for predicting RBES in benign esophageal strictures was established; the risk score ranged from 0 to 8 points, and the risk scores were divided into low (0-2 points), intermediate (3-5 points), and high risk (6-8 points). The proportions of RBES in the corresponding risk categories were 1.0%, 12.2%, and 76.0%, respectively.

Research conclusions

The risk-scoring model predicting RBES in benign esophageal strictures could predict the long-

term outcome of patients with strictures ahead.

Research perspectives

The present study is a retrospective study to develop and validate a model to determine the progression of RBES in patients with benign esophageal strictures. Future case-control studies and prospective studies are required to confirm our findings.

REFERENCES

- 1 **Spechler SJ.** American gastroenterological association medical position statement on treatment of patients with dysphagia caused by benign disorders of the distal esophagus. *Gastroenterology* 1999; **117**: 229-233 [PMID: 10381932 DOI: 10.1016/S0016-5085(99)70572-X]
- 2 **Kobayashi S, Kanai N, Ohki T, Takagi R, Yamaguchi N, Isomoto H, Kasai Y, Hosoi T, Nakao K, Eguchi S, Yamamoto M, Yamato M, Okano T.** Prevention of esophageal strictures after endoscopic submucosal dissection. *World J Gastroenterol* 2014; **20**: 15098-15109 [PMID: 25386058 DOI: 10.3748/wjg.v20.i41.15098]
- 3 **Standards of Practice Committee; Egan JV, Baron TH, Adler DG, Davila R, Faigel DO, Gan SL, Hirota WK, Leighton JA, Lichtenstein D, Qureshi WA, Rajan E, Shen B, Zuckerman MJ, VanGuilder T, Fanelli RD.** Esophageal dilation. *Gastrointest Endosc* 2006; **63**: 755-760 [PMID: 16650533 DOI: 10.1016/j.gie.2006.02.031]
- 4 **Patterson DJ, Graham DY, Smith JL, Schwartz JT, Alpert E, Lanza FL, Cain GD.** Natural history of benign esophageal stricture treated by dilatation. *Gastroenterology* 1983; **85**: 346-350 [PMID: 6862159]
- 5 **de Wijkerslooth LR, Vlegaar FP, Siersema PD.** Endoscopic management of difficult or recurrent esophageal strictures. *Am J Gastroenterol* 2011; **106**: 2080-91; quiz 2092 [PMID: 22008891 DOI: 10.1038/ajg.2011.348]
- 6 **Kochman ML, McClave SA, Boyce HW.** The refractory and the recurrent esophageal stricture: a definition. *Gastrointest Endosc* 2005; **62**: 474-475 [PMID: 16111985 DOI: 10.1016/j.gie.2005.04.050]
- 7 **Spaander MC, Baron TH, Siersema PD, Fuccio L, Schumacher B, Escorsell A, Garcia-Pagan JC, Dumonceau JM, Conio M, de Ceglie A, Skowronek J, Nordmark M, Seufferlein T, Van Gossum A, Hassan C, Repici A, Bruno MJ.** Esophageal stenting for benign and malignant disease: European Society of Gastrointestinal Endoscopy (ESGE) Clinical Guideline. *Endoscopy* 2016; **48**: 939-948 [PMID: 27626318 DOI: 10.1055/s-0042-114210]
- 8 **Fuccio L, Hassan C, Frazzoni L, Miglio R, Repici A.** Clinical outcomes following stent placement in refractory benign esophageal stricture: a systematic review and meta-analysis. *Endoscopy* 2016; **48**: 141-148 [PMID: 26528754 DOI: 10.1055/s-0034-1393331]
- 9 **Young MM, Deschamps C, Trastek VF, Allen MS, Miller DL, Schleck CD, Pairolero PC.** Esophageal reconstruction for benign disease: early morbidity, mortality, and functional results. *Ann Thorac Surg* 2000; **70**: 1651-1655 [PMID: 11093504 DOI: 10.1016/S0003-4975(00)01916-0]
- 10 **Rao SV, McCoy LA, Spertus JA, Krone RJ, Singh M, Fitzgerald S, Peterson ED.** An updated bleeding model to predict the risk of post-procedure bleeding among patients undergoing percutaneous coronary intervention: a report using an expanded bleeding definition from the National Cardiovascular Data Registry CathPCI Registry. *JACC Cardiovasc Interv* 2013; **6**: 897-904 [PMID: 24050858 DOI: 10.1016/j.jcin.2013.04.016]
- 11 **McNamara RL, Kennedy KF, Cohen DJ, Diercks DB, Moscucci M, Ramee S, Wang TY, Connolly T, Spertus JA.** Predicting In-Hospital Mortality in Patients With Acute Myocardial Infarction. *J Am Coll Cardiol* 2016; **68**: 626-635 [PMID: 27491907 DOI: 10.1016/j.jacc.2016.05.049]
- 12 **Andreevski V, Deriban G, Isahi U, Mishevski J, Dimitrova M, Caloska V, Joksimovic N, Popova R, Serafimovski V.** Four Year Results of Conservative Treatment of Benign Strictures of the Esophagus with Savary Gilliard Technique of Bougienage: Cross-Sectional Study Representing First Experiences in Republic of Macedonia. *Pril (Makedon Akad Nauk Umet Odd Med Nauki)* 2018; **39**: 29-35 [PMID: 30110262 DOI: 10.2478/prilozi-2018-0021]
- 13 **Gumaste VV, Dave PB.** Ingestion of corrosive substances by adults. *Am J Gastroenterol* 1992; **87**: 1-5 [PMID: 1728104 DOI: 10.1148/radiology.186.3.794]
- 14 **Lew RJ, Kochman ML.** A review of endoscopic methods of esophageal dilation. *J Clin Gastroenterol* 2002; **35**: 117-126 [PMID: 12172355 DOI: 10.1097/00004836-200208000-00001]
- 15 **Nostrant TT, Nandi PS.** Esophageal dilation. *Gastroenterologist* 1998; **6**: 5-15 [PMID: 9531111 DOI: 10.1016/j.gie.2006.02.031]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

