

1 **Supplement Table 1. Case demographics.**

Patient	Age	Gender	Concomitant medication	Concomitant procedure /treatment	Prior medical history	Prior facial dermatological procedure
Case 1	29	Female	No	No	No	calcium hydroxylapatite the cheeks (2017)
Case 2	28	Female	No	No	No	hyaluronic acid injection in the jaw (2017)

2


3 **Supplement Table S2. Patient self-reported adverse events recorded after True Lift® procedures.**

Patient ID	Visit	Days after surgery	Continuous record (days)	Hypersensitive reactions (none, mild, moderate, severe)					
				Bruises	Erythema	Pain	Tenderness	Itching	Swelling
Case 1	V1	14 days	14	None	None	Moderate (Day 1) Mild (Day 2-3)	Moderate (Day 1) Mild (Day 2-4)	None	None
	V2	1 month	14	None	None	None	None	None	None
	V3	3 months	14	None	None	None	None	None	None
	V4	6 months	14	None	None	None	None	None	None
Case 2	V1	14 days	14	Mild (Day 1-3)	None	Moderate (Day 1-2) Mild (Day 3-6)	Moderate (Day 1-2) Mild (Day 3-7)	Mild (Day 1-7)	Mild (Day 1-2)
	V2	1 month	14	None	None	None	None	None	None
	V3	3 months	14	None	None	None	None	None	None
	V4	6 months	14	None	None	None	None	None	None


4

5

A)



B)



6

7 **Supplement Figure S1.** A) Illustration of True Lift® injections sites. High G' fillers (Restylane Lift
8 Lidocaine) was administered to the five sites as indicated (Figure adopted from Restylane information
9 pamphlet). B) Graphical representation of the facial retaining ligaments system. The five injection sites
10 in (A) correspond to the location of for true retaining ligaments (Figure adopted from Raskin &
11 Latrenta. (2007).