

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 58042

Manuscript Type: CASE REPORT

Coil embolization of arterioportal fistula complicated by gastrointestinal bleeding after Caesarian section: a case report

Arterioportal fistula embolization after C-section

Abstract

BACKGROUND

Most intrahepatic arterioportal fistulae (IAPF) are acquired. The few cases of congenital fistulae are diagnosed in infants and children.

CASE SUMMARY

We report a 31-year-old female patient presenting with haematemesis and melena 3 wk after delivering her second child. The patient had a 20-year history of abdominal

Match Overview

1	Internet 75 words crawled on 26-Dec-2019 www.wjgnet.com	4%
2	Crossref 51 words Lin Wu, Lu Zhao, Ying Lu, Lan He, Xihong Hu. "Intervention ... embolization of congenital intrahepatic shunts in children", Pe	3%
3	Crossref 24 words Thomas J. Ward, Michael L. Marin, Robert A. Lookstein. "Er ... bolization of a giant arterioportal fistula requiring multiple Amp	1%

Coil embolization of arterioportal fistula complicated by gastrointes



ALL

IMAGES

VIDEOS

8,420 Results

Any time ▾

Traumatic hepatic arteriohepatic venous fistula managed ...

<https://www.birpublications.org/doi/10.1259/bjr.20150512>

May 12, 2015 · Hepatic **arterioportal** fistulae are frequent vascular **complications** due to neoplasm, trauma and iatrogenic injury. On the other hand, fistulae between the hepatic arteries and hepatic veins (arteriohepatic venous **fistula**) are rare. We **report** the **case** of a 45-year-old male who suffered from a blunt abdominal trauma with abdominal distension.

Cited by: 4

Author: ChandrasekharanRajsekar, KPSreekumar, ...

Publish Year: 2017

[PDF] A rare case of simultaneous rectal and gastric carcinomas ...

<https://surgicalcasereports.springeropen.com/track...>

Background: Inferior mesenteric **arterioportal fistula** (APF) is rare as only 35 **case** reports in the literature. We herein presented a **case** of simultaneously double cancer in the rectum and **stomach** with inferior mesenteric APF, which is the first **case report** by searching using PubMed. Combination of interventional **embolization** and surgical operation

Gastric variceal bleeding caused by an intrahepatic ...

<https://link.springer.com/10.1007/s12328-011-0277-y> ▾

An intrahepatic **arterioportal fistula** is a rare cause of portal hypertension and variceal **bleeding**. We **report** on a patient with an intrahepatic **arterioportal fistula** following liver biopsy who was successfully treated by hepatectomy **after** unsuccessful arterial **embolization**. We also review the literature on symptomatic intrahepatic **arterioportal** fistulas **after** liver biopsy.

Cited by: 8

Author: Takaaki Iwaki, Hiroyuki Miyatani, Yukio Yos...

Publish Year: 2012

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

8,270 Results

Any time ▾

[Endovascular Management of a Post-Traumatic Complex ...](#)

<https://www.researchgate.net/publication/281733935...>

This **fistula** resulted in ongoing hemodynamically significant **gastrointestinal bleeding**. Initially, **coil embolization** was used to treat the large feeding arteries of the **AV fistula**, but with ...

[PDF] [A rare case of simultaneous rectal and gastric carcinomas ...](#)

<https://surgicalcasereports.springeropen.com/track...>

Background: Inferior mesenteric **arterioportal fistula** (APF) is rare as only 35 **case** reports in the literature. We herein presented a **case** of simultaneously double cancer in the rectum and **stomach** with inferior mesenteric APF, which is the first **case report** by searching using PubMed. Combination of interventional **embolization** and surgical operation

[A large congenital and solitary intrahepatic arterioportal ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2669954>

Apr 07, 2009 · **CASE REPORT**. A 73-year-old woman was admitted to our hospital with complaints of ascites, splenomegaly, abdominal distension and pain. She had been asymptomatic before and denied any medication history as well as history of cirrhosis and hepatic neoplasms, blunt or penetrating trauma, percutaneous liver biopsy, transhepatic cholangiography, gastrectomy and biliary surgery.

Cited by: 9**Author:** Zhen-Ya Lu, Jian-Yang Ao, Tian-An Jiang, Z...**Publish Year:** 2009

[Gastrointestinal bleeding caused by extrahepatic ...](#)

europepmc.org/articles/PMC3508647 ▾

Nov 01, 2012 · An extrahepatic **arterioportal fistula** (APF) involving the gastroduodenal artery and superior mesenteric vein is rare and mostly results from iatrogenic injuries. The clinical symptoms associated with APFs may include abdominal pain, **gastrointestinal bleeding**, ascites, nausea, vomiting, diarrhea, or even congestive heart failure.