

World Journal of *Gastrointestinal Oncology*

World J Gastrointest Oncol 2024 June 15; 16(6): 2264-2866



EDITORIAL

- 2264** Dual primary gastric and colorectal cancer: The known hereditary causes and underlying mechanisms
Azer SA
- 2271** Application of *Fusobacterium nucleatum* as a biomarker in gastrointestinal malignancies
Yu LC, Li YP, Xin YM, Mao M, Pan YX, Qu YX, Luo ZD, Zhang Y, Zhang X
- 2284** T1 colorectal cancer management in the era of minimally invasive endoscopic resection
Jiang SX, Zarrin A, Shahidi N
- 2295** Mixed neuroendocrine and adenocarcinoma of gastrointestinal tract: A complex diagnosis and therapeutic challenge
Shenoy S
- 2300** Advancements in breath-based diagnostics for pancreatic cancer: Current insights and future perspectives
Tez M, Şahingöz E, Marth HF
- 2304** Colorectal cancer and dormant metastases: Put to sleep or destroy?
Senchukova MA

REVIEW

- 2318** Advances in targeted therapy for human epidermal growth factor receptor 2 positive in advanced gastric cancer
Jiang YK, Li W, Qiu YY, Yue M
- 2335** Research progress of ferroptosis regulating lipid peroxidation and metabolism in occurrence and development of primary liver cancer
Shu YJ, Lao B, Qiu YY
- 2350** Early monitoring values of oncogenic signalling molecules for hepatocellular carcinoma
Yao M, Fang RF, Xie Q, Xu M, Sai WL, Yao DF

MINIREVIEWS

- 2362** Therapeutic strategies targeting the epidermal growth factor receptor signaling pathway in metastatic colorectal cancer
Zhou Y, Wu S, Qu FJ
- 2380** Predicting the prognosis of hepatic arterial infusion chemotherapy in hepatocellular carcinoma
Wang QF, Li ZW, Zhou HF, Zhu KZ, Wang YJ, Wang YQ, Zhang YW

- 2394** Unraveling colorectal cancer prevention: The vitamin D - gut flora - immune system nexus

Zhan ZS, Zheng ZS, Shi J, Chen J, Wu SY, Zhang SY

ORIGINAL ARTICLE

Retrospective Cohort Study

- 2404** Unveiling the secrets of gastrointestinal mucous adenocarcinoma survival after surgery with artificial intelligence: A population-based study

Song J, Yan XX, Zhang FL, Lei YY, Ke ZY, Li F, Zhang K, He YQ, Li W, Li C, Pan YM

- 2419** Analysis of metabolic characteristics of metabolic syndrome in elderly patients with gastric cancer by non-targeted metabolomics

Zhang H, Shen WB, Chen L

Retrospective Study

- 2429** Predictive value of preoperative routine examination for the prognosis of patients with pT2N0M0 or pT3N0M0 colorectal cancer

Jing PF, Chen J, Yu ED, Miao CY

- 2439** Simplified liver imaging reporting and data system for the diagnosis of hepatocellular carcinoma on gadoxetic acid-enhanced magnetic resonance imaging

Lyu R, Hu WJ, Wang D, Wang J, Ye YB, Jia KF

- 2449** Efficacy comparison of fruquintinib, regorafenib monotherapy or plus programmed death-1 inhibitors for microsatellite stable metastatic colorectal cancer

An TQ, Qiu H, Zhou QB, Zong H, Hu S, Lian YG, Zhao RH

- 2463** Development of a diagnostic nomogram for alpha-fetoprotein-negative hepatocellular carcinoma based on serological biomarkers

He L, Zhang C, Liu LL, Huang LP, Lu WJ, Zhang YY, Zou DY, Wang YF, Zhang Q, Yang XL

- 2476** Drug-eluting bead transarterial chemoembolization as neoadjuvant therapy pre-liver transplantation for advanced-stage hepatocellular carcinoma

Ye ZD, Zhuang L, Song MC, Yang Z, Zhang W, Zhang JF, Cao GH

- 2487** Association between *Helicobacter pylori* infection, mismatch repair, HER2 and tumor-infiltrating lymphocytes in gastric cancer

Castaneda CA, Castillo M, Bernabe LA, Sanchez J, Fassan M, Tello K, Wistuba II, Chavez Passiuri I, Ruiz E, Sanchez J, Barreda F, Valdivia D, Bazan Y, Abad-Licham M, Mengoa C, Fuentes H, Montenegro P, Poquioma E, Alatrasta R, Flores CJ, Taxa L

- 2504** Impact of baseline hepatitis B virus viral load on the long-term prognosis of advanced hepatocellular carcinoma treated with immunotherapy

Pan D, Liu HN, Yao ZY, Chen XX, Li YQ, Zhu JJ, Han ZX, Qin XB

- 2520** Prediction of pathological complete response and prognosis in locally advanced rectal cancer

Xu YJ, Tao D, Qin SB, Xu XY, Yang KW, Xing ZX, Zhou JY, Jiao Y, Wang LL

Observational Study

- 2531 Extrahepatic cholestasis associated with paracoccidioidomycosis: Challenges in the differential diagnosis of biliopancreatic neoplasia
dos Santos JS, de Moura Arrais V, Rosseto Ferreira WJ, Ribeiro Correa Filho R, Brunaldi MO, Kemp R, Sankanrakutty AK, Elias Junior J, Bellissimo-Rodrigues F, Martinez R, Zangiacomi Martinez E, Ardengh JC

Clinical and Translational Research

- 2541 Development of a novel staging classification for Siewert II adenocarcinoma of the esophagogastric junction after neoadjuvant chemotherapy
Zhang J, Liu H, Yu H, Xu WX
- 2555 N6-methyladenosine methylation regulates the tumor microenvironment of Epstein-Barr virus-associated gastric cancer
Zhang Y, Zhou F, Zhang MY, Feng LN, Guan JL, Dong RN, Huang YJ, Xia SH, Liao JZ, Zhao K
- 2571 Hepatocellular carcinoma: An analysis of the expression status of stress granules and their prognostic value
Ren QS, Sun Q, Cheng SQ, Du LM, Guo PX
- 2592 Comprehensive analysis of clinical and biological value of *ING* family genes in liver cancer
Liu SC
- 2610 Epidemiology and prognostic nomogram for locally advanced gastric signet ring cell carcinoma: A population-based study
Yu ZH, Zhang LM, Dai ZQ, Zhang MN, Zheng SM
- 2631 Socioeconomic traits and the risk of Barrett's esophagus and gastroesophageal reflux disease: A Mendelian randomization study
Liu YX, Bin CL, Zhang L, Yang WT, An BP

Basic Study

- 2646 Complement factor I knockdown inhibits colon cancer development by affecting Wnt/ β -catenin/c-Myc signaling pathway and glycolysis
Du YJ, Jiang Y, Hou YM, Shi YB
- 2663 Fine-needle aspiration technique under endoscopic ultrasound guidance: A technical approach for RNA profiling of pancreatic neoplasms
Seyfedinova SS, Freylikhman OA, Sokolnikova PS, Samochernykh KA, Kostareva AA, Kalinina OV, Solonitsyn EG
- 2673 Comprehensive analysis of gene mutations and mismatch repair in Chinese colorectal cancer patients
Chen H, Jiang RY, Hua Z, Wang XW, Shi XL, Wang Y, Feng QQ, Luo J, Ning W, Shi YF, Zhang DK, Wang B, Jie JZ, Zhong DR
- 2683 Action of circulating and infiltrating B cells in the immune microenvironment of colorectal cancer by single-cell sequencing analysis
Zhang JP, Yan BZ, Liu J, Wang W

- 2697** Bidirectional effects of the tryptophan metabolite indole-3-acetaldehyde on colorectal cancer
Dai Z, Deng KL, Wang XM, Yang DX, Tang CL, Zhou YP
- 2716** Sm-like 5 knockdown inhibits proliferation and promotes apoptosis of colon cancer cells by upregulating p53, CDKN1A and TNFRSF10B
Mo CJ, Deng XY, Ma RL, Zhu K, Shi L, Li K
- 2727** Shi-pi-xiao-ji formula suppresses hepatocellular carcinoma by reducing cellular stiffness through upregulation of acetyl-coA acetyltransferase 1
Jian HY, Liang ZC, Wen H, Zhang Z, Zeng PH
- 2742** Aspirin suppresses hepatocellular carcinoma progression by inhibiting platelet activity
Zhao LJ, Wang ZY, Liu WT, Yu LL, Qi HN, Ren J, Zhang CG
- 2757** Circ_0004592: An auxiliary diagnostic biomarker for gastric cancer
Kong S, Xu YH, Zheng M, Ju SQ, Shi HC
- 2769** N-glycosylation of Wnt3 regulates the progression of hepatocellular carcinoma by affecting Wnt/ β -catenin signal pathway
Zhang XZ, Mo XC, Wang ZT, Sun R, Sun DQ

SYSTEMATIC REVIEWS

- 2781** Ferroptosis regulating lipid peroxidation metabolism in the occurrence and development of gastric cancer
Wang LM, Zhang WW, Qiu YY, Wang F

META-ANALYSIS

- 2793** Meta-analysis of transarterial chemoembolization combined with cryoablation *vs* transarterial chemoembolization alone for ≥ 5 cm hepatocellular carcinoma
Cheng JF, Sun QL, Tang L, Xu XJ, Huang XZ
- 2804** Dynamic contrast enhanced ultrasound in differential diagnosis of hepatocellular carcinoma: A systematic review and meta-analysis
Esposto G, Santini P, Termite F, Galasso L, Mignini I, Ainora ME, Gasbarrini A, Zocco MA
- 2816** Correlation analysis of interstitial maturity and prognosis of colorectal cancer: Meta-analysis
Liu ZJ, Zhang XW, Liu QQ, Wang SZ

SCIENTOMETRICS

- 2826** Visualization analysis of research hotspots and trends on gastrointestinal tumor organoids
Wang G, Liu T, He WT
- 2842** Trends and hotspots in gastrointestinal neoplasms risk assessment: A bibliometric analysis from 1984 to 2022
Fu QQ, Ma L, Niu XM, Zhao HX, Ge XH, Jin H, Yu DH, Yang S

LETTER TO THE EDITOR

- 2862** New perspectives in prognostication of hepatocellular carcinoma: The role and clinical implications of transient receptor potential family genes

Guan SH, Hu WJ, Wang XY, Gu YX, Zhou DH

RETRACTION NOTE

- 2865** Retraction note to: RNA-binding protein CPSF6 regulates IBSP to affect pyroptosis in gastric cancer

Wang XJ, Liu Y, Ke B, Zhang L, Liang H

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Oncology*, Samy Azer, FACG, MD, PhD, Professor, Department of Medical Education, King Saud University College of Medicine, Riyadh 11461, Saudi Arabia. azer2000@optusnet.com.au

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Oncology* (WJGO, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, *etc.*

INDEXING/ABSTRACTING

The WJGO is now abstracted and indexed in PubMed, PubMed Central, Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGO as 3.0; IF without journal self cites: 2.9; 5-year IF: 3.0; Journal Citation Indicator: 0.49; Ranking: 157 among 241 journals in oncology; Quartile category: Q3; Ranking: 58 among 93 journals in gastroenterology and hepatology; and Quartile category: Q3. The WJGO's CiteScore for 2023 is 4.2 and Scopus CiteScore rank 2023: Gastroenterology is 80/167; Oncology is 196/404.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xiang Li; Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Oncology

ISSN

ISSN 1948-5204 (online)

LAUNCH DATE

February 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Monjur Ahmed, Florin Burada

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

PUBLICATION DATE

June 15, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Advances in targeted therapy for human epidermal growth factor receptor 2 positive in advanced gastric cancer

Ya-Kun Jiang, Wei Li, Ying-Yang Qiu, Meng Yue

Specialty type: Oncology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade B, Grade C, Grade C

Novelty: Grade A, Grade B, Grade C

Creativity or Innovation: Grade A, Grade B, Grade C

Scientific Significance: Grade A, Grade B, Grade C

P-Reviewer: Ding J, China; Jin C, China; Maslennikov R, Russia

Received: December 11, 2023

Revised: April 4, 2024

Accepted: April 18, 2024

Published online: June 15, 2024



Ya-Kun Jiang, Meng Yue, Department of Gastroenterology, Central Hospital Affiliated to Shandong First Medical University, Jinan 250013, Shandong Province, China

Wei Li, Health Management Center, Central Hospital Affiliated to Shandong First Medical University, Jinan 250013, Shandong Province, China

Ying-Yang Qiu, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 119077, Singapore

Corresponding author: Meng Yue, MM, Doctor, Department of Gastroenterology, Central Hospital Affiliated to Shandong First Medical University, No. 105 Jiefang Road, Jinan 250013, Shandong Province, China. mengtiger78@163.com

Abstract

Emerging therapeutic methods represented by targeted therapy are effective supplements to traditional first-line chemoradiotherapy resistance. Human epidermal growth factor receptor 2 (*HER2*) is one of the most important targets in targeted therapy for gastric cancer. Trastuzumab combined with chemotherapy has been used as the first-line treatment for advanced gastric cancer. The safety and efficacy of pertuzumab and margetuximab in the treatment of gastric cancer have been verified. However, monoclonal antibodies, due to their large molecular weight, inability to penetrate the blood-brain barrier, and drug resistance, lead to decreased therapeutic efficacy, so it is necessary to explore the efficacy of other *HER2*-targeting therapies in gastric cancer. Small-molecule tyrosine kinase inhibitors, such as lapatinib and pyrrotinib, have the advantages of small molecular weight, penetrating the blood-brain barrier and high oral bioavailability, and are expected to become the drugs of choice for perioperative treatment and neoadjuvant therapy of gastric cancer after validation by large-scale clinical trials in the future. Antibo-drug conjugate, such as T-DM1 and T-DXd, can overcome the resistance of monoclonal antibodies despite their different mechanisms of tumor killing, and are a supplement for the treatment of patients who have failed the treatment of monoclonal antibodies such as trastuzumab. Therefore, after more detailed stratification of gastric cancer patients, various gastric cancer drugs targeting *HER2* are expected to play a more significant role.

Key Words: Human epidermal growth factor receptor 2; Gastric cancer; Targeted therapy; Review

Core Tip: Human epidermal growth factor receptor 2 (*HER2*) is one of the most important targets in targeted therapy for gastric cancer. Trastuzumab combined with chemotherapy has been used as the first-line treatment for advanced gastric cancer. The safety and efficacy of pertuzumab and magtuximab in the treatment of gastric cancer have been verified. However, monoclonal antibodies, due to their large molecular weight, inability to penetrate the blood-brain barrier, and drug resistance, lead to decreased therapeutic efficacy, so it is necessary to explore the efficacy of other *HER2*-targeting therapies in gastric cancer.

Citation: Jiang YK, Li W, Qiu YY, Yue M. Advances in targeted therapy for human epidermal growth factor receptor 2 positive in advanced gastric cancer. *World J Gastrointest Oncol* 2024; 16(6): 2318-2334

URL: <https://www.wjgnet.com/1948-5204/full/v16/i6/2318.htm>

DOI: <https://dx.doi.org/10.4251/wjgo.v16.i6.2318>

INTRODUCTION

Stomach cancer is currently the fifth most common cancer worldwide, with the fourth highest case fatality rate[1-4]. Surgical resection and multidisciplinary treatment combined with chemoradiotherapy, targeted therapy and immunotherapy are the main methods in the treatment of gastric cancer[5]. Because there are no obvious specific symptoms in the early stage, most patients are not diagnosed until the tumor has developed to an advanced stage, resulting in some patients losing the opportunity for surgical treatment[6-8]. The 5-year survival rate of early gastric cancer is 70%, that of advanced gastric cancer is 32%, and that of distant metastatic gastric cancer is only 6%[9]. Therefore, there is an urgent need to find more new and effective therapeutic targets[10-12]. With the development of precision medicine, the score of cancer. Sub-targeted therapy has become the main treatment method for gastric cancer. Human epidermal growth factor receptor 2 (*HER2*) is an important therapeutic target of the epidermal growth factor receptor (*EGFR*) family, with an average positive rate of 17.9% in gastric cancer. In 2010, the ToGA trial was the first to demonstrate that trastuzumab combined with chemotherapy extended survival[13]. At present, in addition to trastuzumab, there are many other *HER2*-targeting therapies still under study, such as monoclonal antibodies (trastuzumab, pertuzumab and magtuximab, *etc.*), antitumor drug conjugate (ADC), such as T-DM1, T-DXd, *etc.*, tyrosine kinase inhibitors (TKI) such as lapatinib, pyrotinib, *etc.*[14-17].

Gastric cancer is a heterogeneous and highly aggressive malignancy with the 5th highest incidence and 3rd highest mortality worldwide[18]. Most patients are in the advanced stage when diagnosed, and the prognosis is very poor. Systemic therapy is currently the most important treatment for advanced gastric cancer, and *HER2* is an important therapeutic target for *HER2*-positive gastric cancer[19-22]. With the continuous optimization of *HER2*-targeting drugs and therapies, some gastric cancer patients have benefited from it. However, the high incidence of drug resistance and serious adverse reactions are still the bottlenecks limiting the application of *HER2*-targeted drugs[23]. Therefore, the development of novel anti-tumor drugs is of great significance for improving the long-term survival of patients with *HER2*-positive gastric cancer. ADCs are a new class of highly effective anti-tumor drugs, which are composed of specific targeted monoclonal antibodies, chemical junctions and small cytotoxic payloads[24]. Their main advantages are powerful therapeutic effect and moderate tissue toxicity[25]. In recent years, ADC has made great progress in the field of targeted therapy for *HER2*-positive advanced gastric cancer[26-30]. First of all, after years of development, a variety of ADCs, including DS-8201, RC48, *etc.*, have been used for second-line and postline treatment of gastric cancer. Secondly, with the progress of ADC bioengineering technology, including high ratio of drug antibodies, cleavable linkers, toxic payloads that can cause bystander effect, *etc.*, new ADCs can play a more significant therapeutic effect in the treatment of specific tumor targets. Some of these ADCs also have multi-targeting and can play an anti-cancer effect against multiple specific targets[31-33]. At the same time, the development of ADCs has reached the third stage, and the new generation of ADCs through site-specific coupling technology has higher homogeneity, more efficient cytotoxic payload, higher precision, and lower non-targeted toxic effects[34-36]. In addition, "target-free combination" therapy consisting of ADC and immune checkpoint inhibitors (ICI) may be a promising treatment strategy for advanced gastric cancer.

This reviews the progress and advantages of *HER2*-targeted therapies in the treatment of gastric cancer in recent years, and discusses the existing problems and future development directions.

TARGETED THERAPY FOR PATIENTS WITH *HER2* GENE POSITIVE GASTRIC CANCER

Many studies[37-40] have shown that the *HER2* gene is overexpressed in different tumors such as breast, ovarian, prostate, colorectal, and lung cancers. The *HER2* gene (also known as Neu or ErbB2) is located on human chromosome 17 (17q21) and encodes the transmembrane glycoprotein p185. Studies[41-45] have shown that the positive rate of *HER2* in gastric cancer is 4.4% to 53.4%, with an average of 17.9%[46-50]. The *HER2* protein has four extracellular domains with unique structural and functional features that interact with ligands and initiate signal transduction, domains I and III

have inherent ligand binding capabilities, domain II can form homologous or heterodimers with other *HER* molecules, and domain IV at the proximal end of the membrane[51-54]. Different ligands are involved in the proliferation and apoptosis of tumor cells by activating Ras/MAPK and PI3K-Akt by binding to different domains of *HER2* receptor[55]. In addition, *HER2* overexpression is closely related to the formation of neovascularization in gastric cancer[56-60]. At present, the recommended methods for detecting *HER2* expression are mainly immunohistochemistry, which can be combined with fluorescence *in situ* hybridization if necessary. Second-generation sequencing, although expensive, is also gaining interest as it helps to find mutations in *HER* genes. Liquid biopsy techniques can also obtain tumor lesion information by detecting circulating tumor cells or circulating tumor DNA. In gastric cancer, *Helicobacter pylori* infection was positively correlated with the positive rate of *HER2* in gastric cancer[61-65]. The difference in the positive rate of *HER2* in gastric cancer was statistically significant between the killed and non-killed groups, suggesting that the detection of *Helicobacter pylori* and *HER2* is of great significance for the detection of early gastric cancer and the evaluation of tumor malignancy[66]. At present, the correlation between *HER2* overexpression and clinical outcome of gastric cancer is uncertain. Some studies[67-70] have shown that *HER2* positivity is associated with a significantly worse prognosis. Other studies[71-75] have found no correlation between *HER2* status and prognosis, so the prognostic value of *HER2* in gastric cancer remains uncertain and needs further study (Figure 1).

HER2 is currently the only clear therapeutic target for gastric cancer, and the ToGA trial is the first large-scale phase III clinical trial in the world to prove that targeted therapy for gastric cancer can prolong survival, making trastuzumab combined with first-line chemotherapy agents the standard first-line treatment for *HER2*-positive advanced gastroesophageal adenocarcinoma[76-80]. However, targeting *HER2* has achieved significant efficacy in the treatment of breast cancer, such as trastuzumab combined with paclitaxel or pertuzumab, the combination therapy of lapatinib and the combination therapy of the antibody coupling drug T-DM1, *etc.*, but these therapies have not achieved positive results in the targeted therapy of *HER2*-positive gastric cancer (Figure 2). At present, the combination of trastuzumab, pertuzumab and magtuximab with immunotherapy drugs is the main direction of clinical trials of targeted therapy for gastric cancer [81]. Due to the emergence of drug resistance, the efficacy of trastuzumab is not good in some patients, so TKI and ADC are the choice of targeted therapy for this part of patients[82-85].

MONOCLONAL ANTIBODY TARGETING *HER2*

Monoclonal antibodies targeting *HER2* recognize the *HER2* antigen through the Fab segment, and the Fc segment binds to immune cells to exert anti-tumor function through antibody-dependent cell-mediated cytotoxicity (ADCC), antibody-dependent cell-mediated phagocytosis, and complement-dependent cytotoxicity[86-88]. Compared with other targeted drugs, *HER2*-targeting monoclonal antibodies have stronger anti-tumor specificity, but because of their relatively large molecular weight, they cannot cross the blood-brain barrier, so the treatment of patients with brain metastases has limitations (Figure 3).

TRASTUZUMAB

Trastuzumab is the only monoclonal antibody approved by the US Food and Drug Administration (FDA) for the first-line treatment of *HER2*-positive advanced gastric cancer[89-93]. It can bind to the extracellular domain IV of *HER2*, block the *HER2* signaling pathway and inhibit the proliferation of tumor cells[94]. The ToGA trial showed that overall survival (OS) in the trastuzumab plus chemotherapy group was superior to chemotherapy alone [median OS (mOS) 13.8 months *vs* 11.1 months] compared with the traditional standard chemotherapy regimens for gastric cancer (capecitabine/cisplatin or fluorouracil/cisplatin)[95]. The ToGA trial also yielded consistent results in subgroup analyses in Japan[96-98]. In addition to survival, patients' recovery adjustment time for health-related quality of life and absence of disease or toxic symptoms was also longer than that for chemotherapy alone[99]. In addition, in the HERXO trial, trastuzumab combined with capecitabine and oxaliplatin were significantly effective in the treatment of advanced gastric cancer patients (mOS was 13.8 months *vs* 7.1 months), the 18-month disease-free survival (DFS) rate was 71% in perioperative treatment of gastric cancer[100-104]. The main results of trastuzumab combined with capecitabine and oxaliplatin in the treatment of advanced gastric cancer were consistent with those of the CGOG1001 trial[105-108]. The above results further established the basic position of trastuzumab in the first-line treatment of gastric cancer.

In the second-line chemotherapy regimen for gastric cancer, 5-FU, calcium folinate, oxaliplatin, irinotecan, docetaxel and paclitaxel are the main chemotherapy drugs at present[109-112]. A clinical trial of trastuzumab combined with 5-FU, leucovorin, oxaliplatin, and docetaxel for perioperative treatment of gastric cancer (median DFS was 42.5 months) demonstrated the efficacy of trastuzumab in second-line chemotherapy for gastric cancer[113-115]. However, the combination of trastuzumab and paclitaxel in the treatment of advanced gastric cancer did not effectively prolong the survival of patients in the WJOG7112G trial and T-CORE1203 trial, which is consistent with the current mainstream view that trastuzumab is not a recommended choice in second-line chemotherapy for gastric cancer[116-118]. Therefore, the efficacy of trastuzumab in second-line chemotherapy for gastric cancer still needs to be further explored[119]. The drug resistance of trastuzumab is often the main reason limiting its efficacy[120]. The use of other drugs in combination with trastuzumab is still the main research direction of clinical trials.




Figure 1 Human epidermal growth factor receptor 2 gene mutation site diagram. Created with BioRender.com (see Supplementary material).




Figure 2 Activation pathway of human epidermal growth factor receptor 2 gene in gastric cancer immunotherapy. Created with BioRender.com (see Supplementary material).

PERTUZUMAB

Pertuzumab is a recombinant humanized monoclonal antibody that binds to the extracellular dimerization domain II of *HER2*, prevents the formation of ligand-induced *HER2* heterodimer, and can play a synergistic role with trastuzumab to inhibit tumor growth[121-126]. The JACOB trial applied pertuzumab and trastuzumab combined chemotherapy in the treatment of gastric cancer. The mOS of the pertuzumab group was 17.5 months, while that of the control group was 14.2 months.

mOS was extended by 3.3 months, but there was no statistical difference between the two, nor was it confirmed that the addition of pertuzumab effectively improved survival time[127-130]. Another clinical trial evaluating the dose and safety of pertuzumab in combination with trastuzumab, capecitabine, and cisplatin in patients with *HER2*-positive advanced gastric cancer showed that patients receiving a valley concentration (62.7 µg/mL) of pertuzumab (840 mg) every 3 wk had a higher response rate than those receiving the first week of pertuzumab loading dose (840 mg). Partial response rates (reduction in tumor volume) were 86% and 55%, respectively, in patients receiving weekly valley concentrations of 40.0 µg/mL of pertuzumab (420 mg)[131-134]. The INNOVATION trial applied pertuzumab combined with trastuzumab and chemotherapy drugs in perioperative chemotherapy for gastric cancer, which will further clarify the significance of pertuzumab in the treatment of gastric cancer[135].

MAGTUXIMAB

Magtuximab is a monoclonal antibody with an engineered Fc domain that targets the *HER2* receptor[136]. The optimized Fc domain improves binding to CD16A, a low-affinity stimulatory receptor found on macrophages and natural killer




Figure 3 Mechanism of action of anti-human epidermal growth factor receptor 2 therapies in gastric cancer. Created with BioRender.com (see Supplementary material).

cells. The optimized Fc domain imparts enhanced ADCC action to the antibody[137-140]. A Phase I clinical trial evaluating magtuximab in patients with *HER2*-positive solid tumors, including gastric cancer, showed that 12% of patients were assessed as having a partial response and 50% were assessed as having stable disease. In addition, magtuximab combined with *PD-1* inhibitors inhibits tumor growth by enhancing innate immunity. The CP-MGAH22-05 trial evaluated the response of magtuximab in combination with pabrolizumab in *HER2*-positive gastroesophageal adenocarcinoma patients with an objective response rate (ORR) of 18.48% and was safe and well tolerated[141-144]. The combination of an FC-optimized anti-*HER2* drug (Magtuximab) with an anti-*PD-1* checkpoint blocker (pembrolizumab) synergistically enhances the activity of these anti-tumor agents[145]. The MAHOGANY trial combined *PD-L1* inhibitor riflezumab with magtuximab in the first-line treatment of gastric cancer, and the results showed that ORR was 53%, the median duration of response was 10.3 months, and the disease control rate was 73%[146-148]. The above results suggest that the combination of dual-targeted drugs is effective in the treatment of gastric cancer.

TKI

HER2 is a transmembrane tyrosine kinase receptor, and TKI represented by Lapatinib blocks *HER2* signaling by competing with ATP in the catalytic domain of the *HER2* molecule, thereby preventing autophosphorylation and subsequent downstream signal transduction events[149]. In theory, TKI has an advantage over monoclonal antibodies because TKI binds to the intracellular domain and thus avoids monoclonal antibody resistance caused by the truncated form (P95) of *HER2* (Figure 4). Monoclonal antibody has a large molecular weight and cannot effectively cross the blood-brain barrier, which has certain limitations in the treatment of patients with brain metastases[150]. In contrast, TKI has a small molecular weight and can play a role across the blood-brain barrier. At the same time, TKI has good oral bioavailability and is an ideal therapeutic drug (Figure 5).




Figure 4 Strategies to overcome mutation-based TKI resistance. A: PDB: 5XFF; B: PDB: 4V01; C: PDB: 4R6V; D: PDB: 6JPJ. Created with BioRender.com (see [Supplementary material](#)).

LAPATINIB

Lapatinib is an oral TKI that blocks *HER1* and *HER2* tyrosine kinase activity by binding to the ATP-binding site of the intracellular domain of the recipient cell, thereby inhibiting tumor cell growth[151,152]. In the LOGiC trial, lapatinib was combined with first-line chemotherapy capecitabine plus oxaliplatin in the treatment of *HER2*-positive gastric cancer. mOS in the lapatinib group and placebo group was 12.2 months and 10.5 months, respectively[153-155]. Although the OS in the lapatinib group was longer, the difference was not statistically significant. In addition, the response rate was 53% in the lapatinib group compared with 39% in the placebo group, and the OS was prolonged in the Asian population and younger patients, suggesting that lapatinib may still play a large role in the treatment of gastric cancer. In the EORTC 40071 trial, mPFS was 8.0 months in the lapatinib group and 5.9 months in the placebo group, suggesting that lapatinib combined with epirubicin, cisplatin, 5-FU, and capecitabine can significantly improve PFS in gastric cancer. Lapatinib did not improve OS in chemotherapy for second-line gastric cancer (mOS 11 months *vs* 8.9 months for lapatinib and paclitaxel alone)[156-160]. In the treatment of metastatic gastric cancer, a trial found that PFS and OS were comparable when treated with lapatinib alone *vs* lapatinib combined with capecitabine after platinum therapy failed. In another trial, lapatinib was used as a neoadjuvant therapy for gastric cancer, but the complete response rate was only 8%[161-164]. Although the above trial results did not reach a consistent positive conclusion, it can suggest that lapatinib can still play a significant role in the treatment of gastric cancer after selecting the right target population.

PYRROTINIB

Pyrrotinib is a novel and irreversible TKI with extensive anti-*EGFR* (*HER1*, *HER2* and *HER4*) activities[165]. By covalently binding to intracellular ATP binding sites, pyrrotinib inhibits the formation of homologous/heterodimer of *HER* family and self-phosphorylation, thereby blocking tumor cell cycle and limiting tumor development. mPFS of gastric cancer patients treated with pyrrotinib were 2.9 months, mOS was 5.9 months, disease control rate was 100%, and ORR was 50%, indicating that the treatment of pyrrotinib has a good effect on *HER2*-positive advanced gastric cancer patients, and its toxic effects can be well controlled[166-170]. In another retrospective study, gastric cancer patients treated with pyrrotinib had PFS of 142 d and mOS of 179 d, which also revealed that pyrrotinib has significant antitumor activity and good safety profile (Figure 6). However, the sample sizes collected in these trials are small, and the role of pyrrotinib in gastric cancer remains to be demonstrated by further larger clinical studies[171-173].




Figure 5 Schematic diagram of TKI-FGFR pathway activation in gastric cancer cells. Created with BioRender.com (see [Supplementary material](#)).

ADC

ADCs consist of recombinant monoclonal antibodies that are covalently bound to cytotoxic drugs. Antibodies can be internalized after binding with specific antigens on the surface of tumor cells, so that cytotoxic drugs can be released inside cells. Another ADC is designed to promote the release of drugs from target cells to extracellular space, or release cytotoxic drugs from antibodies after binding with internalized preantigens, which can achieve the purpose of killing target cells and surrounding cells. High levels of *HER2* expression are not necessarily required[174-176]. This may be beneficial for the treatment of stomach cancer. At present, the clinical ADCs for *HER2*-positive gastric cancer mainly include T-DM1 and T-DXd.

T-DM1

T-DM1 consists of trastuzumab and the small molecule microtubule inhibitor DM1 (maidenin)[177]. In the treatment of breast cancer, T-DM1 has achieved good efficacy and has been approved by the US FDA as a second-line treatment for *HER2*-positive advanced breast cancer. Based on this, many trials[178-180] have investigated its efficacy in the treatment of gastric cancer. The GATSBY trial evaluated the efficacy of T-DM1 and taxanes in *HER2*-positive advanced gastric or gastroesophageal junction carcinoma patients who had previously received chemotherapy and failed targeted therapy. The results showed that T-DM1 was not superior to taxanes in *HER2*-positive advanced gastric cancer patients who had previously received chemotherapy and failed targeted therapy (mOS was 7.9 months *vs* 8.6 months), and the results of the secondary efficacy endpoint PFS analysis did not reach a positive conclusion, which may be related to the *HER2*-positive patients with *HER2* negative conversion[181]. Therefore, there is still a need to explore other therapeutic drugs for these patients.




Figure 6 Mechanisms of alternative signaling activation induced FGFR-TKI resistance in gastric cancer. Created with BioRender.com (see Supplementary material).

T-DXd

T-DXd is a novel ADC composed of trastuzumab, cleavable tetrapeptidase splitter and human DNA topoisomerase I inhibitor[182]. In the Phase I dose-escalation study of T-DXd in patients with advanced *HER2*-positive gastric cancer, the recommended dose of 5.4 mg/kg or 6.4 mg/kg T-DXd was administered intravenously every 3 wk, and 43.2% of patients responded objectively in the dose-expansion trial[183-185]. When T-DXd was used in *HER2*-positive gastric cancer patients who had previously failed treatment with two or more fluoropyrimidine, platinum-based drugs, and trastuzumab (or an approved biosimilar), the difference in OS was statistically significant (12.5 months *vs* 8.4 months) [186]. It provides a new and effective drug for *HER2*-positive gastric cancer patients who have failed chemotherapy and targeted therapy.

RESEARCH PROGRESS OF ADC AND ICI IN *HER2* ADVANCED GASTRIC CANCER

Immunotherapy is a new tumor treatment model. In advanced gastric cancer, the strategies of immunotherapy mainly include adoptive cell therapy, tumor vaccine and immune checkpoint regulation[187]. In recent years, the field of immunotherapy for advanced gastric cancer, especially targeted immune checkpoint therapy, has developed rapidly. The CheckMate 649 study established the important position of ICI in the immunotherapy of advanced gastric cancer. The latest research results were updated at the European Society of Medical Oncology Congress in 2022, which further confirmed the effect of nabulizumab combined with chemotherapy in the whole population, and the ORR of all randomized patients increased from 46% to 58%[188-190]. Both PFS and OS are extended and secure. These follow-up results further support the use of nabulizumab in combination with chemotherapy as the first-line standard of care for patients with advanced gastric cancer. In the Phase III KEYNOTE 062 trial, pembrolizumab, a *PD-1* inhibitor, was no less effective than chemotherapy as a first-line treatment for advanced gastric cancer, either alone or in combination with chemotherapy, and had fewer aes. *PD-1* inhibitors are also beneficial for the treatment of *HER2*-positive gastric cancer. In the Phase III KEYNOTE-811 study, adding pembrolizumab to trastuzumab plus chemotherapy for *HER2*-positive gastric cancer or gastroesophageal junction carcinoma, trastuzumab plus chemotherapy combined with pembrolizumab significantly reduced tumor size. The ORR of the patients was significantly improved. This study is another major breakthrough after the ToGA study, rewriting the first-line treatment guidelines for *HER2*-positive gastric cancer, and opening a new era of "immunotherapy + targeted therapy + chemotherapy" treatment for *HER2*-positive gastric cancer [191]. On this basis, a series of explorations have been carried out on new anti-*HER2* drugs and their combination methods. ZW25, a bi-specific antibody targeting the ECD4 and ECD2 domains of *HER2*, also achieved an ORR and a median PFS of 10.9 months similar to KEYMATE-811 in *HER2*-positive gastric cancer patients treated in combination with *PD-1* inhibitors and chemotherapy[192]. In addition, cytotoxic T lymphocyte-associated antigen 4 (*CTLA-4*) is another important checkpoint (Figure 7).

However, *CTLA-4* checkpoint inhibitors have not been successful in gastric cancer. A *PD-1/CTLA-4* dual-target inhibitor currently under development, cadonilimab (AK104) is the first *PD-1/CTLA-4* bispecific antibody. In a Phase I b/ II study of AK104 in combination with chemotherapy for the first-line treatment of gastric or gastroesophageal borderline cancer (NCT03852251), AK104 showed good anticancer activity and a manageable safety profile[193]. A Phase III study of AK104 combined with chemotherapy as first-line treatment for gastric or gastroesophageal borderline cancer (NCT05008783) is ongoing[194].




Figure 7 Schematic diagram of the effect of PD-1/CTLA-4 signaling on the immune microenvironment of gastric cancer. ICB: Immune checkpoint blockade; ADR: Adverse drug reaction; MHC: Major histocompatibility complex. Created with BioRender.com (see [Supplementary material](#)).

ADC can improve the degree of T lymphocyte infiltration in gastric cancer tissue, enhance the anti-tumor immune response, and thus improve the therapeutic effect of *PD-1* inhibitors. Therefore, a promising therapeutic strategy is ADC combined with ICI. ADC enhances anti-tumor immune response in the following three aspects: First, ADC can induce anti-tumor immune activity by releasing cytotoxic payloads within cancer cells; Secondly, ADC can significantly improve the immune response by directly killing cancer cells or non-targeted cancer cells through enrichment on the tumor surface; Third, cytotoxic payloads can induce antigen-presenting dendritic cells to mature to enhance anti-tumor immune responses. At present, some preclinical studies have confirmed the potential of ADC combined with ICI in the treatment of gastric cancer. Sato *et al*[195] found that XMT-1522 can induce immune cell death *in vitro* and enhance CD8+T cell infiltration and *PD-1* expression level of CD8+ cells after treating multiple gastric cancer cell lines. Yuan *et al*[196] found that T-DXd could increase the number of tumor infiltrating dendritic cells and up-regulate their CD86 expression. In addition, T-DXd also increased the number of tumor-infiltrating CD8+T cells and enhanced the expression levels of *PD-L1* and major histocompatibility complex class I molecules on tumor cells. When combined with ICI, both XMT-1522 and T-DXd showed synergistic antitumor activity. Therefore, the combination therapy composed of ADC and ICI shows an anticancer effect of "1 + 1 > 2", which is expected to become a promising treatment strategy for *HER2*-positive gastric cancer, but it needs to be confirmed by large randomized controlled clinical studies.

CHALLENGES IN THE APPLICATION AND DEVELOPMENT OF ADCS

As the fastest growing *HER2*-targeting drug at present, ADC faces three major challenges in the application and development process: First, how to improve the uptake of ADC by tumor cells is the main difficulty in its development [197]. ADC mainly relies on the high expression of target antigen on the surface of cancer cells to ensure its effective endocytosis and release cytotoxic payloads. The killing effect of *HER2* ADC is closely related to the expression level of *HER2* on the surface of gastric cancer cells, and requires high expression of *HER2* on the surface of gastric cancer cells. Gastric cancer limits the therapeutic effectiveness of existing ADCs. Therefore, improving the cancer cell uptake of ADC can significantly improve its anticancer activity, especially for *HER2*-low expressing gastric cancer. Second, non-targeted toxicity is one of the main factors leading to the failure of ADC clinical trials. Non-targeted toxicity is associated with a variety of factors, including mAb, cytotoxic payloads, linkers and target antigens, as well as the internalization of ADC by cancer cells, non-specific binding of antibodies to Fc receptors, premature hydrolysis of linkers to release payloads, and the "bystander" effect of overstrong ADCC action on normal cells. Thirdly, low expression of target antigen is another important factor in off-target toxicity of ADC[198]. ADC resistance is another problem that needs to be overcome. The results suggest that ineffective internalization of ADC and lysosomal degradation may be the main mechanism of T-DM1 inactivation. It is also possible that cancer cells develop resistance to T-DM1 through upregulation of the drug efflux pump or by alternating the use of tubulin/microtubule-associated proteins. In addition, trastuzumab related changes

may also contribute to the development of T-DM1 resistance, including reduced *HER2* expression levels, the emergence of a truncated form of *HER2*, or mutations in the *ERBB2* gene. At the same time, some key issues should also be addressed in the future, including optimizing each ADC component to improve its anti-cancer activity; Accelerate the exploration of other targets in addition to *HER2*, such as ADCs against Trop-2, C-MET, claudin 18.2, and guanylate cyclase C.

CONCLUSION

Since the ToGA trial started the treatment of *HER2*-positive gastric cancer with trastuzumab, trastuzumab has played a huge role in the first-line treatment of gastric cancer, but in the second-line treatment, the efficacy of trastuzumab is not significant due to the emergence of drug resistance. At the same time, both pertuzumab and magtuximab have shown certain clinical application value in the treatment of gastric cancer. Small molecule TKI drugs lapatinib and pyrrotinib have shown certain efficacy in some clinical trials, but when combined with chemotherapy drugs, they do not significantly improve the survival rate of all patients. However, they still show irreplaceable therapeutic effects in some gastric cancer patients. In addition, the ADC drug T-DM1 did not significantly extend survival in patients with gastric cancer who had failed chemotherapy or targeted therapy, but T-DXd may be the treatment of choice for this subset of patients. Therefore, more detailed classification of *HER2*-positive gastric cancer patients is needed, which will help the use of various targeted drugs to better exert anti-tumor effects. At present, many drugs are still under study, such as monoclonal antibody inatumab, TKI drugs lenatinib, Tucatinib, and ADC SYD985, which may play a good anti-tumor role in the treatment of gastric cancer in the future. At the same time, it is also expected that new and more effective anti-tumor drugs will be developed against the target *HER2*.

FOOTNOTES

Author contributions: Jiang YK wrote the manuscript; Li W and Qiu YY collected the data; Yue M guided the manuscript; all authors reviewed, edited, and approved the final manuscript and revised it critically for important intellectual content, gave final approval of the version to be published, and agreed to be accountable for all aspects of the work.

Conflict-of-interest statement: All the authors declare no conflict of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: China

ORCID number: Meng Yue 0009-0003-7827-5928.

S-Editor: Lin C

L-Editor: A

P-Editor: Zhao S

REFERENCES

- 1 Zeng Y, Jin RU. Molecular pathogenesis, targeted therapies, and future perspectives for gastric cancer. *Semin Cancer Biol* 2022; **86**: 566-582 [PMID: 34933124 DOI: 10.1016/j.semcancer.2021.12.004]
- 2 Patel TH, Cecchini M. Targeted Therapies in Advanced Gastric Cancer. *Curr Treat Options Oncol* 2020; **21**: 70 [PMID: 32725377 DOI: 10.1007/s11864-020-00774-4]
- 3 Meric-Bernstam F, Johnson AM, Dumbrava EEI, Raghav K, Balaji K, Bhatt M, Murthy RK, Rodon J, Piha-Paul SA. Advances in HER2-Targeted Therapy: Novel Agents and Opportunities Beyond Breast and Gastric Cancer. *Clin Cancer Res* 2019; **25**: 2033-2041 [PMID: 30442682 DOI: 10.1158/1078-0432.CCR-18-2275]
- 4 Chen Z, Li Y, Tan B, Zhao Q, Fan L, Li F, Zhao X. Progress and current status of molecule-targeted therapy and drug resistance in gastric cancer. *Drugs Today (Bare)* 2020; **56**: 469-482 [PMID: 32648857 DOI: 10.1358/dot.2020.56.7.3112071]
- 5 Narita Y, Muro K. Challenges in molecular targeted therapy for gastric cancer: considerations for efficacy and safety. *Expert Opin Drug Saf* 2017; **16**: 319-327 [PMID: 27976952 DOI: 10.1080/14740338.2017.1273348]
- 6 Yang W, Raufi A, Klempner SJ. Targeted therapy for gastric cancer: molecular pathways and ongoing investigations. *Biochim Biophys Acta* 2014; **1846**: 232-237 [PMID: 24858418 DOI: 10.1016/j.bbcan.2014.05.003]
- 7 Zeng XQ, Wang J, Chen SY. Methylation modification in gastric cancer and approaches to targeted epigenetic therapy (Review). *Int J Oncol* 2017; **50**: 1921-1933 [PMID: 28498404 DOI: 10.3892/ijo.2017.3981]
- 8 Liu L, Cao L, Gong B, Yu J. Novel biomarkers for the identification and targeted therapy of gastric cancer. *Expert Rev Gastroenterol Hepatol* 2015; **9**: 1217-1226 [PMID: 26220043 DOI: 10.1586/17474124.2015.1072466]

- 9 **Gambardella V**, Castillo J, Tarazona N, Gimeno-Valiente F, Martínez-Ciarpaglini C, Cabeza-Segura M, Roselló S, Roda D, Huerta M, Cervantes A, Fleitas T. The role of tumor-associated macrophages in gastric cancer development and their potential as a therapeutic target. *Cancer Treat Rev* 2020; **86**: 102015 [PMID: 32248000 DOI: 10.1016/j.ctrv.2020.102015]
- 10 **Cann C**, Ciombor KK. Systemic therapy for gastric cancer: Perioperative strategies and beyond. *J Surg Oncol* 2022; **125**: 1151-1160 [PMID: 35230696 DOI: 10.1002/jso.26834]
- 11 **Zhao TT**, Xu H, Xu HM, Wang ZN, Xu YY, Song YX, Yin SC, Liu XY, Miao ZF. The efficacy and safety of targeted therapy with or without chemotherapy in advanced gastric cancer treatment: a network meta-analysis of well-designed randomized controlled trials. *Gastric Cancer* 2018; **21**: 361-371 [PMID: 29455269 DOI: 10.1007/s10120-018-0813-2]
- 12 **Jiang L**, Gong X, Liao W, Lv N, Yan R. Molecular targeted treatment and drug delivery system for gastric cancer. *J Cancer Res Clin Oncol* 2021; **147**: 973-986 [PMID: 33550445 DOI: 10.1007/s00432-021-03520-x]
- 13 **Catanese S**, Lordick F. Targeted and immunotherapy in the era of personalised gastric cancer treatment. *Best Pract Res Clin Gastroenterol* 2021; **50-51**: 101738 [PMID: 33975679 DOI: 10.1016/j.bpg.2021.101738]
- 14 **Rao X**, Zhang C, Luo H, Zhang J, Zhuang Z, Liang Z, Wu X. Targeting Gastric Cancer Stem Cells to Enhance Treatment Response. *Cells* 2022; **11** [PMID: 36139403 DOI: 10.3390/cells11182828]
- 15 **Rosenbaum MW**, Gonzalez RS. Targeted therapy for upper gastrointestinal tract cancer: current and future prospects. *Histopathology* 2021; **78**: 148-161 [PMID: 33382497 DOI: 10.1111/his.14244]
- 16 **Ucaryilmaz Metin C**, Ozcan G. Comprehensive bioinformatic analysis reveals a cancer-associated fibroblast gene signature as a poor prognostic factor and potential therapeutic target in gastric cancer. *BMC Cancer* 2022; **22**: 692 [PMID: 35739492 DOI: 10.1186/s12885-022-09736-5]
- 17 **Garrido M**, Fonseca PJ, Vieitez JM, Frunza M, Lacave AJ. Challenges in first line chemotherapy and targeted therapy in advanced gastric cancer. *Expert Rev Anticancer Ther* 2014; **14**: 887-900 [PMID: 24953238 DOI: 10.1586/14737140.2014.915194]
- 18 **Xu W**, Yang Z, Lu N. Molecular targeted therapy for the treatment of gastric cancer. *J Exp Clin Cancer Res* 2016; **35**: 1 [PMID: 26728266 DOI: 10.1186/s13046-015-0276-9]
- 19 **Cetin B**, Gumusay O, Cengiz M, Ozet A. Advances of Molecular Targeted Therapy in Gastric Cancer. *J Gastrointest Cancer* 2016; **47**: 125-134 [PMID: 26875080 DOI: 10.1007/s12029-016-9806-8]
- 20 **Yazici O**, Sendur MA, Ozdemir N, Aksoy S. Targeted therapies in gastric cancer and future perspectives. *World J Gastroenterol* 2016; **22**: 471-489 [PMID: 26811601 DOI: 10.3748/wjg.v22.i2.471]
- 21 **Al-Batran SE**, Werner D. Recent advances and future trends in the targeted therapy of metastatic gastric cancer. *Expert Rev Gastroenterol Hepatol* 2014; **8**: 555-569 [PMID: 24665840 DOI: 10.1586/17474124.2014.902304]
- 22 **Tomasello G**, Ghidini M, Liguigli W, Ratti M, Toppo L, Passalacqua R. Targeted therapies in gastric cancer treatment: where we are and where we are going. *Invest New Drugs* 2016; **34**: 378-393 [PMID: 26873643 DOI: 10.1007/s10637-016-0330-2]
- 23 **Pellino A**, Riello E, Nappo F, Brignola S, Murgioni S, Djabballah SA, Lonardi S, Zagonel V, Rugge M, Loupakis F, Fassan M. Targeted therapies in metastatic gastric cancer: Current knowledge and future perspectives. *World J Gastroenterol* 2019; **25**: 5773-5788 [PMID: 31636471 DOI: 10.3748/wjg.v25.i38.5773]
- 24 **Pectasides E**. Genomic Alterations and Targeted Therapy in Gastric and Esophageal Adenocarcinoma. *Clin Ther* 2016; **38**: 1589-1599 [PMID: 27041412 DOI: 10.1016/j.clinthera.2016.03.016]
- 25 **Lim SM**, Lim JY, Cho JY. Targeted therapy in gastric cancer: personalizing cancer treatment based on patient genome. *World J Gastroenterol* 2014; **20**: 2042-2050 [PMID: 24587678 DOI: 10.3748/wjg.v20.i8.2042]
- 26 **Yoo C**, Park YS. Companion diagnostics for the targeted therapy of gastric cancer. *World J Gastroenterol* 2015; **21**: 10948-10955 [PMID: 26494953 DOI: 10.3748/wjg.v21.i39.10948]
- 27 **Aichler M**, Lubner B, Lordick F, Walch A. Proteomic and metabolic prediction of response to therapy in gastric cancer. *World J Gastroenterol* 2014; **20**: 13648-13657 [PMID: 25320503 DOI: 10.3748/wjg.v20.i38.13648]
- 28 **Pazo Cid RA**, Antón A. Advanced HER2-positive gastric cancer: current and future targeted therapies. *Crit Rev Oncol Hematol* 2013; **85**: 350-362 [PMID: 23021388 DOI: 10.1016/j.critrevonc.2012.08.008]
- 29 **Wang S**, Yuan L. Predictive biomarkers for targeted and cytotoxic agents in gastric cancer for personalized medicine. *Biosci Trends* 2016; **10**: 171-180 [PMID: 27251446 DOI: 10.5582/bst.2016.01078]
- 30 **Feng W**, Zhu X. Efficacy prediction of targeted therapy for gastric cancer: The current status (Review). *Mol Med Rep* 2018; **18**: 1238-1246 [PMID: 29901092 DOI: 10.3892/mmr.2018.9145]
- 31 **Kim JG**. Molecular targeted therapy for advanced gastric cancer. *Korean J Intern Med* 2013; **28**: 149-155 [PMID: 23525404 DOI: 10.3904/kjim.2013.28.2.149]
- 32 **Harada K**, Yamashita K, Iwatsuki M, Baba H, Ajani JA. Intraperitoneal therapy for gastric cancer peritoneal carcinomatosis. *Expert Rev Clin Pharmacol* 2022; **15**: 43-49 [PMID: 35184625 DOI: 10.1080/17512433.2022.2044790]
- 33 **Luis M**, Tavares A, Carvalho LS, Lara-Santos L, Araújo A, de Mello RA. Personalizing therapies for gastric cancer: molecular mechanisms and novel targeted therapies. *World J Gastroenterol* 2013; **19**: 6383-6397 [PMID: 24151357 DOI: 10.3748/wjg.v19.i38.6383]
- 34 **Aoyama T**, Yoshikawa T. Adjuvant therapy for locally advanced gastric cancer. *Surg Today* 2017; **47**: 1295-1302 [PMID: 28251375 DOI: 10.1007/s00595-017-1493-y]
- 35 **Huang Y**, Wei J, Liu BR. [Research advances of K-ras mutation in the prognosis and targeted therapy of gastric cancer]. *Zhonghua Zhong Liu Za Zhi* 2016; **38**: 81-85 [PMID: 26899324 DOI: 10.3760/cma.j.issn.0253-3766.2016.02.001]
- 36 **Farran B**, Müller S, Montenegro RC. Gastric cancer management: Kinases as a target therapy. *Clin Exp Pharmacol Physiol* 2017; **44**: 613-622 [PMID: 28271563 DOI: 10.1111/1440-1681.12743]
- 37 **Wu L**, Zhong Y, Yu X, Wu D, Xu P, Lv L, Ruan X, Liu Q, Feng Y, Liu J, Li X. Selective poly adenylation predicts the efficacy of immunotherapy in patients with lung adenocarcinoma by multiple omics research. *Anticancer Drugs* 2022; **33**: 943-959 [PMID: 35946526 DOI: 10.1097/CAD.0000000000001319]
- 38 **Ding J**, Kang X, Feng M, Tan J, Feng Q, Wang X, Wang J, Liu J, Li Z, Guan W, Qiao T. A novel active mitochondrion-selective fluorescent probe for the NIR fluorescence imaging and targeted photodynamic therapy of gastric cancer. *Biomater Sci* 2022; **10**: 4756-4763 [PMID: 35837996 DOI: 10.1039/d2bm00684g]
- 39 **Ang YL**, Yong WP, Tan P. Translating gastric cancer genomics into targeted therapies. *Crit Rev Oncol Hematol* 2016; **100**: 141-146 [PMID: 26947813 DOI: 10.1016/j.critrevonc.2016.02.007]

- 40 **Murayama Y**, Oritani K, Tsutsui S. Novel CD9-targeted therapies in gastric cancer. *World J Gastroenterol* 2015; **21**: 3206-3213 [PMID: 25805926 DOI: 10.3748/wjg.v21.i11.3206]
- 41 **Tabernero J**, Macarulla T, Ramos FJ, Baselga J. Novel targeted therapies in the treatment of gastric and esophageal cancer. *Ann Oncol* 2005; **16**: 1740-1748 [PMID: 15980157 DOI: 10.1093/annonc/mdi355]
- 42 **Zheng T**, Jin J, Zhang Y, Zhou L. Efficacy and safety of paclitaxel with or without targeted therapy as second-line therapy in advanced gastric cancer: A meta-analysis. *Medicine (Baltimore)* 2020; **99**: e20734 [PMID: 32569214 DOI: 10.1097/MD.00000000000020734]
- 43 **Lyons TG**, Ku GY. Systemic therapy for esophagogastric cancer: targeted therapies. *Chin Clin Oncol* 2017; **6**: 48 [PMID: 29129088 DOI: 10.21037/cco.2017.07.02]
- 44 **Cappetta A**, Lonardi S, Pastorelli D, Bergamo F, Lombardi G, Zagonel V. Advanced gastric cancer (GC) and cancer of the gastro-oesophageal junction (GEJ): focus on targeted therapies. *Crit Rev Oncol Hematol* 2012; **81**: 38-48 [PMID: 21256046 DOI: 10.1016/j.critrevonc.2010.12.006]
- 45 **Zhang J**, Zhao T, Han F, Hu Y, Li Y. Photothermal and gene therapy combined with immunotherapy to gastric cancer by the gold nanoshell-based system. *J Nanobiotechnology* 2019; **17**: 80 [PMID: 31277667 DOI: 10.1186/s12951-019-0515-x]
- 46 **Ye M**, Xiu LJ, Ji QQ, Zhang YC, Sun YW, Zhao Y, Wang D, Li YJ, Wang XW, Yue XQ, Sun DZ. Research progress in targeted therapies for gastric cancer. *Int J Clin Pharmacol Ther* 2022; **60**: 509-514 [PMID: 36197788 DOI: 10.5414/CP204250]
- 47 **Zhou H**, Yuan M, Yu Q, Zhou X, Min W, Gao D. Autophagy regulation and its role in gastric cancer and colorectal cancer. *Cancer Biomark* 2016; **17**: 1-10 [PMID: 27314289 DOI: 10.3233/CBM-160613]
- 48 **Jou E**, Rajdev L. Current and emerging therapies in unresectable and recurrent gastric cancer. *World J Gastroenterol* 2016; **22**: 4812-4823 [PMID: 27239108 DOI: 10.3748/wjg.v22.i20.4812]
- 49 **Bekaii-Saab T**, El-Rayes B. Identifying and targeting cancer stem cells in the treatment of gastric cancer. *Cancer* 2017; **123**: 1303-1312 [PMID: 28117883 DOI: 10.1002/cncr.30538]
- 50 **Zhou YL**, Li YM, He WT. Application of Mesenchymal Stem Cells in the Targeted Gene Therapy for Gastric Cancer. *Curr Stem Cell Res Ther* 2016; **11**: 434-439 [PMID: 26496889 DOI: 10.2174/1574888x10666151026113818]
- 51 **Lin YH**, Chen ZR, Lai CH, Hsieh CH, Feng CL. Active Targeted Nanoparticles for Oral Administration of Gastric Cancer Therapy. *Biomacromolecules* 2015; **16**: 3021-3032 [PMID: 26286711 DOI: 10.1021/acs.biomac.5b00907]
- 52 **Zhu ZG**. [Current status and prospect of perioperative therapy for locally advanced gastric cancer]. *Zhonghua Wei Chang Wai Ke Za Zhi* 2021; **24**: 101-106 [PMID: 33508913 DOI: 10.3760/cma.j.cn.441530-20210105-00003]
- 53 **Lee SY**, Oh SC. Changing strategies for target therapy in gastric cancer. *World J Gastroenterol* 2016; **22**: 1179-1189 [PMID: 26811656 DOI: 10.3748/wjg.v22.i3.1179]
- 54 **Costa G**, Younes H, Kourie HR, Kattan J. The rapidly evolving landscape of advanced gastric cancer therapy. *Future Oncol* 2022; **18**: 1413-1416 [PMID: 35129395 DOI: 10.2217/fon-2021-1499]
- 55 **Wu L**, Zheng Y, Ruan X, Wu D, Xu P, Liu J, Li X. Long-chain noncoding ribonucleic acids affect the survival and prognosis of patients with esophageal adenocarcinoma through the autophagy pathway: construction of a prognostic model. *Anticancer Drugs* 2022; **33**: e590-e603 [PMID: 34338240 DOI: 10.1097/CAD.0000000000001189]
- 56 **Gerson JN**, Skariah S, Denlinger CS, Astsaturov I. Perspectives of HER2-targeting in gastric and esophageal cancer. *Expert Opin Investig Drugs* 2017; **26**: 531-540 [PMID: 28387541 DOI: 10.1080/13543784.2017.1315406]
- 57 **Majeed W**, Iftikhar A, Khaliq T, Aslam B, Muzaffar H, Atta K, Mahmood A, Waris S. Gastric Carcinoma: Recent Trends in Diagnostic Biomarkers and Molecular Targeted Therapies. *Asian Pac J Cancer Prev* 2016; **17**: 3053-3060 [PMID: 27509928]
- 58 **Davidson M**, Okines AF, Starling N. Current and Future Therapies for Advanced Gastric Cancer. *Clin Colorectal Cancer* 2015; **14**: 239-250 [PMID: 26524924 DOI: 10.1016/j.clcc.2015.05.013]
- 59 **Sunkar S**, Necharika D, Nellore J, Nachiyar VC, Peela S. Small-Molecule-Targeted Therapies for Gastrointestinal Cancer: Successes and Failures. *Crit Rev Oncog* 2020; **25**: 311-333 [PMID: 33639060 DOI: 10.1615/CritRevOncog.2020036206]
- 60 **Joo MK**, Park JJ, Chun HJ. Recent updates of precision therapy for gastric cancer: Towards optimal tailored management. *World J Gastroenterol* 2016; **22**: 4638-4650 [PMID: 27217696 DOI: 10.3748/wjg.v22.i19.4638]
- 61 **Durães C**, Almeida GM, Seruca R, Oliveira C, Carneiro F. Biomarkers for gastric cancer: prognostic, predictive or targets of therapy? *Virchows Arch* 2014; **464**: 367-378 [PMID: 24487788 DOI: 10.1007/s00428-013-1533-y]
- 62 **Asaoka Y**, Ikenoue T, Koike K. New targeted therapies for gastric cancer. *Expert Opin Investig Drugs* 2011; **20**: 595-604 [PMID: 21406037 DOI: 10.1517/13543784.2011.566863]
- 63 **Chen J**, Röcken C, Malferteiner P, Ebert MP. Recent advances in molecular diagnosis and therapy of gastric cancer. *Dig Dis* 2004; **22**: 380-385 [PMID: 15812163 DOI: 10.1159/000083602]
- 64 **Zhang LH**, Ji JF. [Prognostic biomarkers and targeted therapies for gastric cancer]. *Zhonghua Wei Chang Wai Ke Za Zhi* 2013; **16**: 193-196 [PMID: 23446485]
- 65 **Le DT**, Ott PA, Korytowsky B, Le H, Le TK, Zhang Y, Maglinte GA, Abraham P, Patel D, Shanguan T, Chau I. Real-world Treatment Patterns and Clinical Outcomes Across Lines of Therapy in Patients With Advanced/Metastatic Gastric or Gastroesophageal Junction Cancer. *Clin Colorectal Cancer* 2020; **19**: 32-38.e3 [PMID: 31813769 DOI: 10.1016/j.clcc.2019.09.001]
- 66 **Aguiar PN Jr**, Muniz TP, Miranda RR, Tadokoro H, Forones NM, Monteiro ID, Castelo-Branco P, Janjigian YY, De Mello RA. Current advances in targeted therapies for metastatic gastric cancer: improving patient care. *Future Oncol* 2016; **12**: 839-854 [PMID: 26838766 DOI: 10.2217/fon.15.348]
- 67 **Wu L**, Liu Q, Ruan X, Luan X, Zhong Y, Liu J, Yan J, Li X. Multiple Omics Analysis of the Role of RBM10 Gene Instability in Immune Regulation and Drug Sensitivity in Patients with Lung Adenocarcinoma (LUAD). *Biomedicines* 2023; **11** [PMID: 37509501 DOI: 10.3390/biomedicines11071861]
- 68 **Yoong J**, Michael M, Leong T. Targeted therapies for gastric cancer: current status. *Drugs* 2011; **71**: 1367-1384 [PMID: 21812503 DOI: 10.2165/11592530-000000000-00000]
- 69 **Lianos GD**, Mangano A, Cho WC, Roukos DH. From standard to new genome-based therapy of gastric cancer. *Expert Rev Gastroenterol Hepatol* 2015; **9**: 1023-1026 [PMID: 26108988 DOI: 10.1586/17474124.2015.1060856]
- 70 **Mu YP**, Sun WJ, Lu CW, Su XL. MicroRNAs May Serve as Emerging Molecular Biomarkers for Diagnosis and Prognostic Assessment or as Targets for Therapy in Gastric Cancer. *Asian Pac J Cancer Prev* 2015; **16**: 4813-4820 [PMID: 26163596 DOI: 10.7314/apjcp.2015.16.12.4813]
- 71 **Athauda A**, Segelov E, Ali Z, Chau I. Integrative molecular analysis of colorectal cancer and gastric cancer: What have we learnt? *Cancer*

- Treat Rev* 2019; **73**: 31-40 [PMID: 30597479 DOI: 10.1016/j.ctrv.2018.12.004]
- 72 **Yong J**, Li Y, Lin S, Wang Z, Xu Y. Inhibitors Targeting YAP in Gastric Cancer: Current Status and Future Perspectives. *Drug Des Devel Ther* 2021; **15**: 2445-2456 [PMID: 34140763 DOI: 10.2147/DDDT.S308377]
- 73 **Ouyang J**, Long Z, Li G. Circular RNAs in Gastric Cancer: Potential Biomarkers and Therapeutic Targets. *Biomed Res Int* 2020; **2020**: 2790679 [PMID: 32685459 DOI: 10.1155/2020/2790679]
- 74 **Irmak-Yazicioglu MB**. Mechanisms of MicroRNA Deregulation and MicroRNA Targets in Gastric Cancer. *Oncol Res Treat* 2016; **39**: 136-139 [PMID: 27031864 DOI: 10.1159/000443224]
- 75 **Shin VY**, Chu KM. MiRNA as potential biomarkers and therapeutic targets for gastric cancer. *World J Gastroenterol* 2014; **20**: 10432-10439 [PMID: 25132759 DOI: 10.3748/wjg.v20.i30.10432]
- 76 **Zhang M**, Deng W, Cao X, Shi X, Zhao H, Duan Z, Lv B, Liu B. Concurrent apatinib and local radiation therapy for advanced gastric cancer: A case report and review of the literature. *Medicine (Baltimore)* 2017; **96**: e6241 [PMID: 28248891 DOI: 10.1097/MD.0000000000006241]
- 77 **Iyikesici MS**. Survival outcomes of metabolically supported chemotherapy combined with ketogenic diet, hyperthermia, and hyperbaric oxygen therapy in advanced gastric cancer. *Niger J Clin Pract* 2020; **23**: 734-740 [PMID: 32367884 DOI: 10.4103/njcp.njcp_509_18]
- 78 **Janjigian YY**, Shah MA. Molecularly targeted therapies in advanced gastric cancer. *Minerva Gastroenterol Dietol* 2011; **57**: 75-88 [PMID: 21372772]
- 79 **Ngeow J**, Tan IB, Choo SP. Targeted therapies in the treatment of gastric cancer. *Asia Pac J Clin Oncol* 2011; **7**: 224-235 [PMID: 21884434 DOI: 10.1111/j.1743-7563.2011.01407.x]
- 80 **Lin F**, Tan HJ, Guan JS, Lim YP. Divide and conquer: subproteomic approaches toward gastric cancer biomarker and drug target discovery. *Expert Rev Proteomics* 2014; **11**: 515-530 [PMID: 24684179 DOI: 10.1586/14789450.2014.904751]
- 81 **Zhang F**, Yin J, Zhang C, Han M, Wang X, Fu S, Du J, Zhang H, Li W. Affibody-Conjugated RALA Polymers Delivering Oligomeric 5-Fluorodeoxyuridine for Targeted Therapy of HER2 Overexpressing Gastric Cancer. *Macromol Biosci* 2020; **20**: e2000083 [PMID: 32558229 DOI: 10.1002/mabi.202000083]
- 82 **Kang X**, Li W, Liu W, Liang H, Deng J, Wong CC, Zhao S, Kang W, To KF, Chiu PWY, Wang G, Yu J, Ng EKW. LIMK1 promotes peritoneal metastasis of gastric cancer and is a therapeutic target. *Oncogene* 2021; **40**: 3422-3433 [PMID: 33883692 DOI: 10.1038/s41388-021-01656-1]
- 83 **De Vita F**, Giuliani F, Silvestris N, Rossetti S, Pizzolorusso A, Santabarbara G, Galizia G, Colucci G, Ciardiello F, Orditura M. Current status of targeted therapies in advanced gastric cancer. *Expert Opin Ther Targets* 2012; **16** Suppl 2: S29-S34 [PMID: 22443228 DOI: 10.1517/14728222.2011.652616]
- 84 **Tyczyńska M**, Kędzierawski P, Karakula K, Januszewski J, Kozak K, Sitarz M, Forma A. Treatment Strategies of Gastric Cancer-Molecular Targets for Anti-angiogenic Therapy: a State-of-the-art Review. *J Gastrointest Cancer* 2021; **52**: 476-488 [PMID: 33761051 DOI: 10.1007/s12029-021-00629-7]
- 85 **Ren Z**, Sun J, Sun X, Hou H, Li K, Ge Q. Efficacy and safety of different molecular targeted agents based on chemotherapy for gastric cancer patients treatment: a network meta-analysis. *Oncotarget* 2017; **8**: 48253-48262 [PMID: 28477027 DOI: 10.18632/oncotarget.17192]
- 86 **Kuo HY**, Yeh KH. Molecular-targeted therapy for chemotherapy-refractory gastric cancer: a case report and literature review. *Anticancer Res* 2014; **34**: 3695-3699 [PMID: 24982389]
- 87 **Ciliberto D**, Staropoli N, Caglioti F, Gualtieri S, Fiorillo L, Chiellino S, De Angelis AM, Mendicino F, Botta C, Caraglia M, Tassone P, Tagliaferri P. A systematic review and meta-analysis of randomized trials on the role of targeted therapy in the management of advanced gastric cancer: Evidence does not translate? *Cancer Biol Ther* 2015; **16**: 1148-1159 [PMID: 26061272 DOI: 10.1080/15384047.2015.1056415]
- 88 **Zagouri F**, Papadimitriou CA, Dimopoulos MA, Pectasides D. Molecularly targeted therapies in unresectable-metastatic gastric cancer: a systematic review. *Cancer Treat Rev* 2011; **37**: 599-610 [PMID: 21676549 DOI: 10.1016/j.ctrv.2011.03.007]
- 89 **Weidle UH**, Birzele F, Brinkmann U, Auslaender S. Gastric Cancer: Identification of microRNAs Inhibiting Druggable Targets and Mediating Efficacy in Preclinical In Vivo Models. *Cancer Genomics Proteomics* 2021; **18**: 497-514 [PMID: 34183383 DOI: 10.21873/cgp.20275]
- 90 **Kim HS**, Lee H, Shin SJ, Beom SH, Jung M, Bae S, Lee EY, Park KH, Choi YY, Son T, Kim HI, Cheong JH, Hyung WJ, Park JC, Shin SK, Lee SK, Lee YC, Koom WS, Lim JS, Chung HC, Noh SH, Rha SY, Kim H, Paik S. Complementary utility of targeted next-generation sequencing and immunohistochemistry panels as a screening platform to select targeted therapy for advanced gastric cancer. *Oncotarget* 2017; **8**: 38389-38398 [PMID: 28418920 DOI: 10.18632/oncotarget.16409]
- 91 **Chandra R**, Balachandran N, Wang S, Reznik S, Zeh H, Porembka M. The changing face of gastric cancer: epidemiologic trends and advances in novel therapies. *Cancer Gene Ther* 2021; **28**: 390-399 [PMID: 33009508 DOI: 10.1038/s41417-020-00234-z]
- 92 **Aprile G**, Giampieri R, Bonotto M, Bittoni A, Ongaro E, Cardellino GG, Graziano F, Giuliani F, Fasola G, Cascinu S, Scartozzi M. The challenge of targeted therapies for gastric cancer patients: the beginning of a long journey. *Expert Opin Investig Drugs* 2014; **23**: 925-942 [PMID: 24806575 DOI: 10.1517/13543784.2014.912631]
- 93 **Liu J**, Liu YP. [Progress and prospects of international cancer drug clinical research on gastric cancer]. *Zhonghua Wei Chang Wai Ke Za Zhi* 2020; **23**: 196-200 [PMID: 32074802 DOI: 10.3760/cma.j.issn.1671-0274.2020.02.018]
- 94 **Weidle UH**, Birzele F, Auslaender S, Brinkmann U. Down-regulated MicroRNAs in Gastric Carcinoma May Be Targets for Therapeutic Intervention and Replacement Therapy. *Anticancer Res* 2021; **41**: 4185-4202 [PMID: 34475038 DOI: 10.21873/anticancer.15223]
- 95 **Xie S**, Zhang H, Wang X, Ge Q, Hu J. The relative efficacy and safety of targeted agents used in combination with chemotherapy in treating patients with untreated advanced gastric cancer: a network meta-analysis. *Oncotarget* 2017; **8**: 26959-26968 [PMID: 28460479 DOI: 10.18632/oncotarget.15923]
- 96 **Zou K**, Yang S, Zheng L, Yang C, Xiong B. Efficacy and safety of target combined chemotherapy in advanced gastric cancer: a meta-analysis and system review. *BMC Cancer* 2016; **16**: 737 [PMID: 27633381 DOI: 10.1186/s12885-016-2772-5]
- 97 **Hescheler DA**, Plum PS, Zander T, Quaa A, Korenkov M, Gassa A, Michel M, Bruns CJ, Alakus H. Identification of targeted therapy options for gastric adenocarcinoma by comprehensive analysis of genomic data. *Gastric Cancer* 2020; **23**: 627-638 [PMID: 32107691 DOI: 10.1007/s10120-020-01045-9]
- 98 **Wagner AD**, Moehler M. Development of targeted therapies in advanced gastric cancer: promising exploratory steps in a new era. *Curr Opin Oncol* 2009; **21**: 381-385 [PMID: 19412098 DOI: 10.1097/CCO.0b013e32832c42e0]
- 99 **Koizumi K**, Kato S, Sakurai H, Hashimoto I, Yasumoto K, Saiki I. Therapeutics target of CXCR4 and its downstream in peritoneal carcinomatosis of gastric cancer. *Front Biosci (Schol Ed)* 2012; **4**: 269-276 [PMID: 22202059 DOI: 10.2741/s267]
- 100 **Carneiro F**, Oliveira C, Leite M, Seruca R. Molecular targets and biological modifiers in gastric cancer. *Semin Diagn Pathol* 2008; **25**: 274-287 [PMID: 19013893 DOI: 10.1053/j.semdp.2008.07.004]

- 101 **Wu L**, Zhong Y, Wu D, Xu P, Ruan X, Yan J, Liu J, Li X. Immunomodulatory Factor TIM3 of Cytolytic Active Genes Affected the Survival and Prognosis of Lung Adenocarcinoma Patients by Multi-Omics Analysis. *Biomedicines* 2022; **10** [PMID: 36140350 DOI: 10.3390/biomedicines10092248]
- 102 **Cao HY**, Yuan AH, Chen W, Shi XS, Miao Y. A DNA aptamer with high affinity and specificity for molecular recognition and targeting therapy of gastric cancer. *BMC Cancer* 2014; **14**: 699 [PMID: 25248985 DOI: 10.1186/1471-2407-14-699]
- 103 **Ohtsu A**, Sasako M. Overview of adjuvant therapy for resected gastric cancer: differences in Japan and the United States. *Semin Oncol* 2005; **32**: S101-S104 [PMID: 16399444 DOI: 10.1053/j.seminoncol.2005.06.001]
- 104 **Dionisi F**, Avery S, Lukens JN, Ding X, Kralik J, Kirk M, Roses RE, Amichetti M, Metz JM, Plastaras JP. Proton therapy in adjuvant treatment of gastric cancer: planning comparison with advanced x-ray therapy and feasibility report. *Acta Oncol* 2014; **53**: 1312-1320 [PMID: 24797885 DOI: 10.3109/0284186X.2014.912351]
- 105 **Wang J**, Du L, Chen X. Adenosine signaling: Optimal target for gastric cancer immunotherapy. *Front Immunol* 2022; **13**: 1027838 [PMID: 36189223 DOI: 10.3389/fimmu.2022.1027838]
- 106 **Kim S**, Barzi A, Rajdev L. Biomarker-driven targeted therapies for gastric/gastro-esophageal junction malignancies. *Semin Oncol* 2018; **45**: 133-150 [PMID: 30262395 DOI: 10.1053/j.seminoncol.2018.03.001]
- 107 **Wang X**, Li G, Zhang Y, Bai S, Xu F, Wei Y, Gong Y. Single-arc volumetric-modulated arc therapy (sVMAT) as adjuvant treatment for gastric cancer: dosimetric comparisons with three-dimensional conformal radiotherapy (3D-CRT) and intensity-modulated radiotherapy (IMRT). *Med Dosim* 2013; **38**: 395-400 [PMID: 23791504 DOI: 10.1016/j.meddos.2013.04.007]
- 108 **Curea FG**, Hebbar M, Ilie SM, Bacinschi XE, Trifanescu OG, Botnariuc I, Anghel RM. Current Targeted Therapies in HER2-Positive Gastric Adenocarcinoma. *Cancer Biother Radiopharm* 2017; **32**: 351-363 [PMID: 29265917 DOI: 10.1089/cbr.2017.2249]
- 109 **Lee JA**, Ahn YC, Lim DH, Park HC, Asranbaeva MS. Dosimetric and Clinical Influence of 3D Versus 2D Planning in Postoperative Radiation Therapy for Gastric Cancer. *Cancer Res Treat* 2015; **47**: 727-737 [PMID: 25672580 DOI: 10.4143/crt.2014.018]
- 110 **Wu L**, Li X, Qian X, Wang S, Liu J, Yan J. Lipid Nanoparticle (LNP) Delivery Carrier-Assisted Targeted Controlled Release mRNA Vaccines in Tumor Immunity. *Vaccines (Basel)* 2024; **12** [PMID: 38400169 DOI: 10.3390/vaccines12020186]
- 111 **Du S**, Fang Z, Ye L, Sun H, Deng G, Wu W, Zeng F. Pretreatment neutrophil-to-lymphocyte ratio predicts the benefit of gastric cancer patients with systemic therapy. *Aging (Albany NY)* 2021; **13**: 17638-17654 [PMID: 34245559 DOI: 10.18632/aging.203256]
- 112 **Shi J**, Liu S, Yu Y, He C, Tan L, Shen YM. RGD peptide-decorated micelles assembled from polymer-paclitaxel conjugates towards gastric cancer therapy. *Colloids Surf B Biointerfaces* 2019; **180**: 58-67 [PMID: 31028965 DOI: 10.1016/j.colsurf.2019.04.042]
- 113 **Arkenau HT**, Saggese M, Lemech C. Advanced gastric cancer: is there enough evidence to call second-line therapy standard? *World J Gastroenterol* 2012; **18**: 6376-6378 [PMID: 23197882 DOI: 10.3748/wjg.v18.i44.6376]
- 114 **Hashemi M**, Aparviz R, Beickzade M, Paskeh MDA, Kheirabad SK, Koohpar ZK, Moravej A, Dehghani H, Saebfar H, Zandieh MA, Salimimoghadam S, Rashidi M, Taheriazam A, Entezari M, Samarghandian S. Advances in RNAi therapies for gastric cancer: Targeting drug resistance and nanoscale delivery. *Biomed Pharmacother* 2023; **169**: 115927 [PMID: 38006616 DOI: 10.1016/j.biopha.2023.115927]
- 115 **Abdel-Rahman O**. Targeting vascular endothelial growth factor (VEGF) pathway in gastric cancer: preclinical and clinical aspects. *Crit Rev Oncol Hematol* 2015; **93**: 18-27 [PMID: 24970311 DOI: 10.1016/j.critrevonc.2014.05.012]
- 116 **Padmanabhan N**, Ushijima T, Tan P. How to stomach an epigenetic insult: the gastric cancer epigenome. *Nat Rev Gastroenterol Hepatol* 2017; **14**: 467-478 [PMID: 28513632 DOI: 10.1038/nrgastro.2017.53]
- 117 **Fährmann M**. Targeting protein kinase C (PKC) in physiology and cancer of the gastric cell system. *Curr Med Chem* 2008; **15**: 1175-1191 [PMID: 18473812 DOI: 10.2174/092986708784310413]
- 118 **Yoshida K**, Yamaguchi K, Okumura N, Osada S, Takahashi T, Tanaka Y, Tanabe K, Suzuki T. The roles of surgical oncologists in the new era: minimally invasive surgery for early gastric cancer and adjuvant surgery for metastatic gastric cancer. *Pathobiology* 2011; **78**: 343-352 [PMID: 22104206 DOI: 10.1159/000328197]
- 119 **Yu L**, Zhou GQ, Li DC. MiR-136 triggers apoptosis in human gastric cancer cells by targeting AEG-1 and BCL2. *Eur Rev Med Pharmacol Sci* 2018; **22**: 7251-7256 [PMID: 30468468 DOI: 10.26355/eurrev_201811_16259]
- 120 **Khan U**, Shah MA. Optimizing Therapies in the Perioperative Management of Gastric Cancer. *Curr Treat Options Oncol* 2019; **20**: 57 [PMID: 31144054 DOI: 10.1007/s11864-019-0654-1]
- 121 **Li M**, Chang J, Ren H, Song D, Guo J, Peng L, Zhou X, Zhao K, Lu S, Liu Z, Hu P. Downregulation of CCKBR Expression Inhibits the Proliferation of Gastric Cancer Cells, Revealing a Potential Target for Immunotoxin Therapy. *Curr Cancer Drug Targets* 2022; **22**: 257-268 [PMID: 34994328 DOI: 10.2174/1568009622666220106113616]
- 122 **Pan SY**, Fang JY. [Research progresses in targeted therapies for gastric cancer]. *Zhonghua Nei Ke Za Zhi* 2020; **59**: 148-152 [PMID: 32074690 DOI: 10.3760/cma.j.issn.0578-1426.2020.02.011]
- 123 **Azimi M**, Totonchi M, Rahimi M, Firouzi J, Sahranavard P, Emami Razavi A, Memari F, Kamali F, Ebrahimi M. An integrated analysis to predict micro-RNAs targeting both stemness and metastasis in human gastric cancer. *J Gastroenterol Hepatol* 2021; **36**: 436-445 [PMID: 32633423 DOI: 10.1111/jgh.15176]
- 124 **Lei Z**, Fan J, Li X, Chen Y, Shi D, Xie H, Liu B. Biomimetic graphene oxide quantum dots nanoparticles targeted photothermal-chemotherapy for gastric cancer. *J Drug Target* 2023; **31**: 320-333 [PMID: 36647791 DOI: 10.1080/1061186X.2022.2162060]
- 125 **Wu L**, Li H, Liu Y, Fan Z, Xu J, Li N, Qian X, Lin Z, Li X, Yan J. Research progress of 3D-bioprinted functional pancreas and *in vitro* tumor models. *IJB* 2024; **10**: 1256 [DOI: 10.36922/ijb.1256]
- 126 **Briasoulis E**, Liakakos T, Dova L, Fatouros M, Tsekeris P, Roukos DH, Kappas AM. Selecting a specific pre- or postoperative adjuvant therapy for individual patients with operable gastric cancer. *Expert Rev Anticancer Ther* 2006; **6**: 931-939 [PMID: 16761937 DOI: 10.1586/14737140.6.6.931]
- 127 **Abdel-Rahman O**. Targeting the hepatocyte growth factor/mesenchymal epithelial transition pathway in gastric cancer: biological rationale and clinical applications. *Expert Rev Anticancer Ther* 2015; **15**: 235-245 [PMID: 25353620 DOI: 10.1586/14737140.2014.974564]
- 128 **Zheng Y**, Li Z, Wang Y, Chen W, Lin Y, Guo J, Ye G. CircRNA: A new class of targets for gastric cancer drug resistance therapy. *Pathol Oncol Res* 2023; **29**: 1611033 [PMID: 37065861 DOI: 10.3389/pore.2023.1611033]
- 129 **Kim HM**, Jeong I, Kim HJ, Kang SK, Kwon WS, Kim TS, Park KH, Jung M, Soong J, Lin SC, Chung HC, Rha SY. Casein Kinase 2 Inhibitor, CX-4945, as a Potential Targeted Anticancer Agent in Gastric Cancer. *Anticancer Res* 2018; **38**: 6171-6180 [PMID: 30396934 DOI: 10.21873/anticancer.12970]
- 130 **Geng H**, Qian R, Zhang L, Yang C, Xia X, Wang C, Zhao G, Zhang Z, Zhu C. Clinical outcomes and potential therapies prediction of

- subgroups based on a ferroptosis-related long non-coding RNA signature for gastric cancer. *Aging (Albany NY)* 2022; **14**: 6358-6376 [PMID: 35969182 DOI: 10.18632/aging.204227]
- 131 Lee J, Kim ST, Kim K, Lee H, Kozarewa I, Mortimer PGS, Odegaard JI, Harrington EA, Lee J, Lee T, Oh SY, Kang JH, Kim JH, Kim Y, Ji JH, Kim YS, Lee KE, Kim J, Sohn TS, An JY, Choi MG, Lee JH, Bae JM, Kim S, Kim JJ, Min YW, Min BH, Kim NKD, Luke S, Kim YH, Hong JY, Park SH, Park JO, Park YS, Lim HY, Talasaz A, Hollingsworth SJ, Kim KM, Kang WK. Tumor Genomic Profiling Guides Patients with Metastatic Gastric Cancer to Targeted Treatment: The VIKTORY Umbrella Trial. *Cancer Discov* 2019; **9**: 1388-1405 [PMID: 31315834 DOI: 10.1158/2159-8290.CD-19-0442]
- 132 Ayyappan S, Prabhakar D, Sharma N. Epidermal growth factor receptor (EGFR)-targeted therapies in esophagogastric cancer. *Anticancer Res* 2013; **33**: 4139-4155 [PMID: 24122977]
- 133 Yoshida K, Tsujimoto H, Matsumura K, Kinoshita M, Takahata R, Matsumoto Y, Hiraki S, Ono S, Seki S, Yamamoto J, Hase K. CD47 is an adverse prognostic factor and a therapeutic target in gastric cancer. *Cancer Med* 2015; **4**: 1322-1333 [PMID: 26077800 DOI: 10.1002/cam4.478]
- 134 Feng Y, Sun T, Yu Y, Gao Y, Wang X, Chen Z. MicroRNA-370 inhibits the proliferation, invasion and EMT of gastric cancer cells by directly targeting PAQR4. *J Pharmacol Sci* 2018; **138**: 96-106 [PMID: 30322804 DOI: 10.1016/j.jphs.2018.08.004]
- 135 Xu L, Zhang XY, Zhang SY. [In vitro and in vivo targeting therapy of immunoliposomes against human gastric cancer]. *Zhonghua Yi Xue Za Zhi* 1994; **74**: 83-86, 126 [PMID: 8069726]
- 136 Moehler M, Galle PR, Gockel I, Junginger T, Schmidberger H. The multidisciplinary management of gastrointestinal cancer. Multimodal treatment of gastric cancer. *Best Pract Res Clin Gastroenterol* 2007; **21**: 965-981 [PMID: 18070698 DOI: 10.1016/j.bpg.2007.10.003]
- 137 Xia J, Wang H, Li S, Wu Q, Sun L, Huang H, Zeng M. Ion channels or aquaporins as novel molecular targets in gastric cancer. *Mol Cancer* 2017; **16**: 54 [PMID: 28264681 DOI: 10.1186/s12943-017-0622-y]
- 138 Chen Z, Huang W, Tian T, Zang W, Wang J, Liu Z, Li Z, Lai Y, Jiang Z, Gao J, Shen L. Characterization and validation of potential therapeutic targets based on the molecular signature of patient-derived xenografts in gastric cancer. *J Hematol Oncol* 2018; **11**: 20 [PMID: 29433585 DOI: 10.1186/s13045-018-0563-y]
- 139 Zhang W. TCGA divides gastric cancer into four molecular subtypes: implications for individualized therapeutics. *Chin J Cancer* 2014; **33**: 469-470 [PMID: 25223913 DOI: 10.5732/cjc.014.10117]
- 140 Pentheroudakis G, Stoyianni A. Incorporation of targeted agents in the management of patients with advanced gastric cancer. *Curr Med Chem* 2011; **18**: 1629-1639 [PMID: 21428884 DOI: 10.2174/092986711795471365]
- 141 Zhang SJ, Feng JF, Wang L, Guo W, Du YW, Ming L, Zhao GQ. miR-1303 targets claudin-18 gene to modulate proliferation and invasion of gastric cancer cells. *Dig Dis Sci* 2014; **59**: 1754-1763 [PMID: 24647998 DOI: 10.1007/s10620-014-3107-5]
- 142 Yashiro M, Matsuoka T. Fibroblast growth factor receptor signaling as therapeutic targets in gastric cancer. *World J Gastroenterol* 2016; **22**: 2415-2423 [PMID: 26937130 DOI: 10.3748/wjg.v22.i8.2415]
- 143 Matsukura N, Onda M, Shimada T. [Possibility and future problems of gene therapy for gastric cancer]. *Nihon Geka Gakkai Zasshi* 2001; **102**: 778-782 [PMID: 11681005]
- 144 Lu M, Wu Y, Zhang Y, Yu Y, Wang S, Su X. Immunotherapeutic strategy in the management of gastric cancer: molecular profiles, current practice, and ongoing trials. *J Egypt Natl Canc Inst* 2023; **35**: 32 [PMID: 37779128 DOI: 10.1186/s43046-023-00192-1]
- 145 Kataoka K, Deleersnijder A, Lordick F. Will molecular target agents enable the multidisciplinary treatment in stage IV gastric cancer? *Eur J Surg Oncol* 2017; **43**: 1835-1845 [PMID: 28888797 DOI: 10.1016/j.ejso.2017.08.006]
- 146 Carneiro P, Figueiredo J, Bordeira-Carriço R, Fernandes MS, Carvalho J, Oliveira C, Seruca R. Therapeutic targets associated to E-cadherin dysfunction in gastric cancer. *Expert Opin Ther Targets* 2013; **17**: 1187-1201 [PMID: 23957294 DOI: 10.1517/14728222.2013.827174]
- 147 Wen L, Cheng F, Zhou Y, Yin C. MiR-26a enhances the sensitivity of gastric cancer cells to cisplatin by targeting NRAS and E2F2. *Saudi J Gastroenterol* 2015; **21**: 313-319 [PMID: 26458859 DOI: 10.4103/1319-3767.166206]
- 148 Wu L, Zheng Y, Liu J, Luo R, Wu D, Xu P, Li X. Comprehensive evaluation of the efficacy and safety of LPV/r drugs in the treatment of SARS and MERS to provide potential treatment options for COVID-19. *Aging (Albany NY)* 2021; **13**: 10833-10852 [PMID: 33879634 DOI: 10.18632/aging.202860]
- 149 Sun W, Jiang C, Ji Y, Xiao C, Song H. Long Noncoding RNAs: New Regulators of Resistance to Systemic Therapies for Gastric Cancer. *Biomed Res Int* 2021; **2021**: 8853269 [PMID: 33506041 DOI: 10.1155/2021/8853269]
- 150 Ma YY, Tao HQ. Role of urokinase plasminogen activator receptor in gastric cancer: a potential therapeutic target. *Cancer Biother Radiopharm* 2012; **27**: 285-290 [PMID: 22702495 DOI: 10.1089/cbr.2012.1232]
- 151 Liu Y, Li D, Chen Y, Liu Y, Lin Y, Huang X, Wu T, Wang C, Ding J. Integrated bioinformatics analysis for conducting a prognostic model and identifying immunotherapeutic targets in gastric cancer. *BMC Bioinformatics* 2023; **24**: 191 [PMID: 37161430 DOI: 10.1186/s12859-023-05312-1]
- 152 Zhang Y, Zhu WG, Han JH, Zhang XY, Tao GZ, Li T, Yu CH. [Application of simplified intensity modulated radiation therapy in gastric cancer after operation]. *Zhonghua Wei Chang Wai Ke Za Zhi* 2013; **16**: 268-272 [PMID: 23536350]
- 153 Li Y, Qin C. MiR-1179 inhibits the proliferation of gastric cancer cells by targeting HMGB1. *Hum Cell* 2019; **32**: 352-359 [PMID: 30903594 DOI: 10.1007/s13577-019-00244-6]
- 154 Labots M, Buffart TE, Haan JC, van Grieken NC, Tijssen M, van de Velde CJ, Grabsch HI, Ylstra B, Carvalho B, Fijneman RJ, Verheul HM, Meijer GA. High-level copy number gains of established and potential drug target genes in gastric cancer as a lead for treatment development and selection. *Cell Oncol (Dordr)* 2014; **37**: 41-52 [PMID: 24379144 DOI: 10.1007/s13402-013-0162-4]
- 155 Pandian J, Ganesan K. Delineation of gastric tumors with activated ERK/MAPK signaling cascades for the development of targeted therapeutics. *Exp Cell Res* 2022; **410**: 112956 [PMID: 34864005 DOI: 10.1016/j.yexcr.2021.112956]
- 156 Ali SM, Sanford EM, Klempner SJ, Robinson DA, Wang K, Palma NA, Chmielecki J, Yelensky R, Palmer GA, Morosini D, Lipson D, Catenacci DV, Braithe F, Erlich R, Stephens PJ, Ross JS, Ou SH, Miller VA. Prospective comprehensive genomic profiling of advanced gastric carcinoma cases reveals frequent clinically relevant genomic alterations and new routes for targeted therapies. *Oncologist* 2015; **20**: 499-507 [PMID: 25882375 DOI: 10.1634/theoncologist.2014-0378]
- 157 Deng N, Goh LK, Wang H, Das K, Tao J, Tan IB, Zhang S, Lee M, Wu J, Lim KH, Lei Z, Goh G, Lim QY, Tan AL, Sin Poh DY, Riahi S, Bell S, Shi MM, Linnartz R, Zhu F, Yeoh KG, Toh HC, Yong WP, Cheong HC, Rha SY, Boussioutas A, Grabsch H, Rozen S, Tan P. A comprehensive survey of genomic alterations in gastric cancer reveals systematic patterns of molecular exclusivity and co-occurrence among distinct therapeutic targets. *Gut* 2012; **61**: 673-684 [PMID: 22315472 DOI: 10.1136/gutjnl-2011-301839]

- 158 **Alani S**, Soyfer V, Strauss N, Schifter D, Corn BW. Limited advantages of intensity-modulated radiotherapy over 3D conformal radiation therapy in the adjuvant management of gastric cancer. *Int J Radiat Oncol Biol Phys* 2009; **74**: 562-566 [PMID: 19427558 DOI: 10.1016/j.ijrobp.2008.09.061]
- 159 **Choi ES**, Kim H, Kim HP, Choi Y, Goh SH. CD44v8-10 as a potential theranostic biomarker for targeting disseminated cancer cells in advanced gastric cancer. *Sci Rep* 2017; **7**: 4930 [PMID: 28694503 DOI: 10.1038/s41598-017-05247-7]
- 160 **Lee J**, Ou SH. Towards the goal of personalized medicine in gastric cancer--time to move beyond HER2 inhibition. Part I: Targeting receptor tyrosine kinase gene amplification. *Discov Med* 2013; **15**: 333-341 [PMID: 23819947]
- 161 **Corso S**, Giordano S. How Can Gastric Cancer Molecular Profiling Guide Future Therapies? *Trends Mol Med* 2016; **22**: 534-544 [PMID: 27260398 DOI: 10.1016/j.molmed.2016.05.004]
- 162 **Di Cicilia R**, Mordenti P, Anselmi E, Paties C, Carella E, Cavanna L. HER2-positive gastric cancer showing marked thickening of the gastric wall on ultrasonographic and computed tomographic scans: a chance phenomenon or a specific behaviour of this cancer type? Report of three cases. *Anticancer Res* 2014; **34**: 5083-5086 [PMID: 25202095]
- 163 **Xu C**, Ooi WF, Qamra A, Tan J, Chua BY, Ho SWT, Das K, Adam Isa ZF, Li Z, Yao X, Yan T, Xing M, Huang KK, Lin JS, Nandi T, Tay ST, Lee MH, Tan ALK, Ong X, Ashktorab H, Smoot D, Li S, Ng SC, Teh BT, Tan P. HNF4 α pathway mapping identifies wild-type IDH1 as a targetable metabolic node in gastric cancer. *Gut* 2020; **69**: 231-242 [PMID: 31068366 DOI: 10.1136/gutjnl-2018-318025]
- 164 **Choi KM**, Cho E, Bang G, Lee SJ, Kim B, Kim JH, Park SG, Han EH, Chung YH, Kim JY, Kim E. Activity-Based Protein Profiling Reveals Potential Dasatinib Targets in Gastric Cancer. *Int J Mol Sci* 2020; **21** [PMID: 33291786 DOI: 10.3390/ijms21239276]
- 165 **Liu J**, Wang H, Liao X. MicroRNA-223-5p targets long non-coding RNA TP73 antisense RNA1 to promote the invasion of gastric cancer. *Hum Cell* 2020; **33**: 676-682 [PMID: 32248369 DOI: 10.1007/s13577-020-00349-3]
- 166 **Zou P**, Xia Y, Chen W, Chen X, Ying S, Feng Z, Chen T, Ye Q, Wang Z, Qiu C, Yang S, Liang G. EF24 induces ROS-mediated apoptosis via targeting thioredoxin reductase 1 in gastric cancer cells. *Oncotarget* 2016; **7**: 18050-18064 [PMID: 26919110 DOI: 10.18632/oncotarget.7633]
- 167 **Zhang H**, Tian B, Yu H, Yao H, Gao Z. LAPTM4B-35 protein as a potential therapeutic target in gastric cancer. *Tumour Biol* 2014; **35**: 12737-12742 [PMID: 25416438 DOI: 10.1007/s13277-014-2599-0]
- 168 **Zhang X**, Zhang J, Ma Y, Pei X, Liu Q, Lu B, Jin L, Wang J, Liu J. A cell-based single-stranded DNA aptamer specifically targets gastric cancer. *Int J Biochem Cell Biol* 2014; **46**: 1-8 [PMID: 24501772]
- 169 **Iwagami S**, Watanabe M, Baba H. Current status and problems in development of molecular target agents for gastric cancer in Japan. *Nihon Shokakibyo Gakkai Zasshi* 2011; **108**: 11-18 [PMID: 21212590]
- 170 **Zhi Q**, Guo X, Guo L, Zhang R, Jiang J, Ji J, Zhang J, Chen X, Cai Q, Li J, Liu B, Zhu Z, Yu Y. Oncogenic miR-544 is an important molecular target in gastric cancer. *Anticancer Agents Med Chem* 2013; **13**: 270-275 [PMID: 22934698 DOI: 10.2174/1871520611313020013]
- 171 **Gong M**, Lu Z, Fang G, Bi J, Xue X. A small interfering RNA targeting osteopontin as gastric cancer therapeutics. *Cancer Lett* 2008; **272**: 148-159 [PMID: 18694621 DOI: 10.1016/j.canlet.2008.07.004]
- 172 **Diniz F**, Lamas S, Osório H, Aguiar P, Freitas D, Gärtner F, Sarmento B, Reis CA, Gomes J. Nanoparticles targeting Sialyl-Tn for efficient tyrosine kinase inhibitor delivery in gastric cancer. *Acta Biomater* 2023; **170**: 142-154 [PMID: 37586448 DOI: 10.1016/j.actbio.2023.08.014]
- 173 **Sohal DP**, Sun W. What is the reference cytotoxic regimen for advanced gastric cancer: standard versus optimal therapy? *Expert Rev Anticancer Ther* 2011; **11**: 1843-1850 [PMID: 22117152 DOI: 10.1586/era.11.170]
- 174 **Fuereder T**, Hoeflmayer D, Jaeger-Lansky A, Rasin-Streden D, Strommer S, Fisker N, Hansen BJ, Crevenna R, Wacheck V. Sphingosine kinase 1 is a relevant molecular target in gastric cancer. *Anticancer Drugs* 2011; **22**: 245-252 [PMID: 21360847 DOI: 10.1097/cad.0b013e328340bd95]
- 175 **Wang K**, Gong Z, Chen Y, Zhang M, Wang S, Yao S, Liu Z, Huang Z, Fei B. KDM4C-mediated senescence defense is a targetable vulnerability in gastric cancer harboring TP53 mutations. *Clin Epigenetics* 2023; **15**: 163 [PMID: 37848946 DOI: 10.1186/s13148-023-01579-6]
- 176 **Seo JH**, Jeong ES, Choi YK. Therapeutic effects of lentivirus-mediated shRNA targeting of cyclin D1 in human gastric cancer. *BMC Cancer* 2014; **14**: 175 [PMID: 24618206 DOI: 10.1186/1471-2407-14-175]
- 177 **Chang HR**, Nam S, Lee J, Kim JH, Jung HR, Park HS, Park S, Ahn YZ, Huh I, Balch C, Ku JL, Powis G, Park T, Jeong JH, Kim YH. Systematic approach identifies RHOA as a potential biomarker therapeutic target for Asian gastric cancer. *Oncotarget* 2016; **7**: 81435-81451 [PMID: 27806312 DOI: 10.18632/oncotarget.12963]
- 178 **Kim Y**, Hosoda Y, Nishino M, Okano M, Kawada J, Yamasaki M, Nagai K, Yasui M, Okuyama M, Tsujinaka T. [Hepatic Resection of Multiple Liver Metastases from Gastric Cancer after Molecular Targeted Chemotherapy(S-I plus Cisplatin plus Trastuzumab)]. *Gan To Kagaku Ryoho* 2015; **42**: 1635-1637 [PMID: 26805121]
- 179 **Wu L**, Chen X, Zeng Q, Lai Z, Fan Z, Ruan X, Li X, Yan J. NR5A2 gene affects the overall survival of LUAD patients by regulating the activity of CSCs through SNP pathway by OCLR algorithm and immune score. *Heliyon* 2024; **10**: e28282 [PMID: 38601554 DOI: 10.1016/j.heliyon.2024.e28282]
- 180 **Inoue M**, Matsumoto S, Saito H, Tsujitani S, Ikeguchi M. Intraperitoneal administration of a small interfering RNA targeting nuclear factor-kappa B with paclitaxel successfully prolongs the survival of xenograft model mice with peritoneal metastasis of gastric cancer. *Int J Cancer* 2008; **123**: 2696-2701 [PMID: 18798274 DOI: 10.1002/ijc.23867]
- 181 **Michiura T**, Miyazaki S, Fujita S, Hirose H, Miyazaki Y, Sugimoto S, Yamabe K, Nagaoka M. [A case of advanced HER2- positive gastric cancer with a partial response to molecular targeted chemotherapy (S-I+CDDP+Trastuzumab)]. *Gan To Kagaku Ryoho* 2014; **41**: 2346-2348 [PMID: 25731518]
- 182 **Yamamoto K**, Nishikawa K, Hirao M, Fujitani K, Tsujinaka T, Maeda S, Uemura M, Miyake M, Hama N, Miyamoto A, Omiya H, Ikeda M, Takami K, Nakamori S, Sekimoto M. [Optimal Target Population for Adjuvant Chemotherapy for Pathological T1 Gastric Cancer]. *Gan To Kagaku Ryoho* 2015; **42**: 2084-2087 [PMID: 26805272]
- 183 **Akamo Y**, Mizuno I, Ichino T, Yamamoto T, Yasui T, Itabashi Y, Saito T, Kurahashi S, Tanimoto N, Shinagawa N. [Endoscopic injection of liposomal adriamycin targeting lymph node metastasis of gastric cancer]. *Gan To Kagaku Ryoho* 1991; **18**: 1822-1824 [PMID: 1877824]
- 184 **Akamo Y**, Mizuno I, Ichino T, Yamamoto T, Yasui T, Itabashi Y, Saito T, Kurahashi S, Tanimoto N, Shinagawa N. [Efficacy of preoperative adjuvant chemotherapy using adriamycin targeting the regional lymph nodes for gastric cancer--lymph node-targeting delivery of adriamycin in rabbits]. *Gan To Kagaku Ryoho* 1992; **19**: 451-455 [PMID: 1558394]
- 185 **D'Ugo D**, Agnes A, Grieco M, Biondi A, Persiani R. Global updates in the treatment of gastric cancer: a systematic review. Part 2: perioperative management, multimodal therapies, new technologies, standardization of the surgical treatment and educational aspects. *Updates*

- Surg* 2020; **72**: 355-378 [PMID: 32306277 DOI: 10.1007/s13304-020-00771-0]
- 186 **Zhang Y**, Wen Y. The Role of LncRNA NEAT1 in Gastric Cancer: Tumor Development and Potential Therapeutic Target. *Curr Pharm Des* 2023; **29**: 2213-2222 [PMID: 37779403 DOI: 10.2174/0113816128257113230921035222]
- 187 **Iinuma H**, Maruyama K, Okinaga K, Sasaki K, Sekine T, Ishida O, Ogiwara N, Johkura K, Yonemura Y. Intracellular targeting therapy of cisplatin-encapsulated transferrin-polyethylene glycol liposome on peritoneal dissemination of gastric cancer. *Int J Cancer* 2002; **99**: 130-137 [PMID: 11948504 DOI: 10.1002/ijc.10242]
- 188 **He H**, Song F, Gao Q, Lu Z, Yuan Y, Li X, Chen L, Jia C, Yang R, Yang J, Duan W, Hou Y. The APEX1/miRNA-27a-5p axis plays key roles in progression, metastasis and targeted chemotherapy of gastric cancer. *Int J Pharm* 2021; **599**: 120446 [PMID: 33675923 DOI: 10.1016/j.ijpharm.2021.120446]
- 189 **Yang Y**, Ma Y, Xiang X, Xing P, Wu Y, Zhang L, Tian Y. The prognostic value of the lymph node ratio for local advanced gastric cancer patients with intensity-modulated radiation therapy and concurrent chemotherapy after radical gastrectomy in China. *Radiat Oncol* 2020; **15**: 237 [PMID: 33054848 DOI: 10.1186/s13014-020-01687-0]
- 190 **Aoyama T**, Yoshikawa T, Fujikawa H, Hayashi T, Ogata T, Cho H, Yamada T, Hasegawa S, Tsuchida K, Yukawa N, Oshima T, Rino Y, Masuda M. Exploratory Analysis to Find Unfavorable Subset of Stage II Gastric Cancer for Which Surgery Alone Is the Standard Treatment; Another Target for Adjuvant Chemotherapy. *Int Surg* 2014; **99**: 835-841 [PMID: 25437596 DOI: 10.9738/INTSURG-D-13-00176.1]
- 191 **Liu Y**, Chen S, Xue R, Zhao J, Di M. Mefloquine effectively targets gastric cancer cells through phosphatase-dependent inhibition of PI3K/Akt/mTOR signaling pathway. *Biochem Biophys Res Commun* 2016; **470**: 350-355 [PMID: 26780727 DOI: 10.1016/j.bbrc.2016.01.046]
- 192 **Liang H**, Li Z, Huang Z, Wu C, Qiu Y, Liang Y, Chen X, Li F, Xu Z, Li G, Liu H, Zhao L. Prognostic characteristics and clinical response to immunotherapy targeting programmed cell death 1 for patients with advanced gastric cancer with liver metastases. *Front Immunol* 2022; **13**: 1015549 [PMID: 36211361 DOI: 10.3389/fimmu.2022.1015549]
- 193 **Li X**, Xu J, Xie J, Yang W. Research progress in targeted therapy and immunotherapy for gastric cancer. *Chin Med J (Engl)* 2022; **135**: 1299-1313 [PMID: 35830242 DOI: 10.1097/CM9.00000000000002185]
- 194 **Wieland P**, Dobler B, Mai S, Hermann B, Tiefenbacher U, Steil V, Wenz F, Lohr F. IMRT for postoperative treatment of gastric cancer: covering large target volumes in the upper abdomen: a comparison of a step-and-shoot and an arc therapy approach. *Int J Radiat Oncol Biol Phys* 2004; **59**: 1236-1244 [PMID: 15234061 DOI: 10.1016/j.ijrobp.2004.02.051]
- 195 **Sato T**, Saito M, Nakajima S, Saito K, Katagata M, Fukai S, Okayama H, Sakamoto W, Saze Z, Momma T, Mimura K, Kono K. ARID1A deficiency is targetable by AKT inhibitors in HER2-negative gastric cancer. *Gastric Cancer* 2023; **26**: 379-392 [PMID: 36811690 DOI: 10.1007/s10120-023-01373-6]
- 196 **Yuan DD**, Zhu ZX, Zhang X, Liu J. Targeted therapy for gastric cancer: Current status and future directions (Review). *Oncol Rep* 2016; **35**: 1245-1254 [PMID: 26718131 DOI: 10.3892/or.2015.4528]
- 197 **Pandey SS**, Shaikh FI, Gupta AR, Vaidya RJ. Mannosylated Solid Lipid Nanocarriers of Chrysin to Target Gastric Cancer: Optimization and Cell Line Study. *Curr Drug Deliv* 2021; **18**: 1574-1584 [PMID: 33745434 DOI: 10.2174/1567201818666210319142206]
- 198 **Dong W**, Li B, Wang J, Song Y, Zhang Z, Fu C. Exploration of the Radiotherapeutic Clinical Target Volume Delineation for Gastric Cancer from Lymph Node Metastases. *Anticancer Res* 2017; **37**: 4629-4634 [PMID: 28739763 DOI: 10.21873/anticancer.11864]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

