



国内版 国际版

Fasciola Hepatica presenting as Colon Cancer Liver Metastasis on 18F-Fluorodeox



All Images Videos

翻译成中文 关闭取词

12,200 Results Any time ▾

### (PDF) Fasciola Hepatica Mimicking Malignancy on 18F ...

[https://www.researchgate.net/publication/308753629\\_Fasciola\\_Hepatica\\_Mimicking...](https://www.researchgate.net/publication/308753629_Fasciola_Hepatica_Mimicking...)

Fasciola Hepatica Mimicking Malignancy on 18F-Fluorodeoxyglucose-Positron Emission Tomography/Computed Tomography Article (PDF Available) · September 2016 with 59 Reads DOI: 10.4274/mirt.97759

### Fasciola Hepatica Mimicking Malignancy on 18F ...

[europepmc.org/articles/PMC5100087](http://europepmc.org/articles/PMC5100087)

Oct 01, 2016 · Positron emission tomography/computed tomography (PET/CT) findings of **Fasciola hepatica**. PET/CT scan was performed 60 min after i.v. injection of 256 MBq of 18 F-fluorodeoxyglucose (FDG), using a lutetium oxyorthosilicate crystal equipped PET/CT (mCT20 Siemens, Germany). Maximum intensity projection images (A) show high FDG uptakes in different areas of the liver.

Author: Erdem Sürücü, Yusuf Demir, Ahmet C... Publish Year: 2016

### Selection of Patients for Resection of Hepatic Metastases ...

[https://pubs.rsna.org/doi/10.1148/radiographics.21.suppl\\_1.g01oc05s55](https://pubs.rsna.org/doi/10.1148/radiographics.21.suppl_1.g01oc05s55)

Oct 01, 2001 · A rapidly emerging clinical application of **positron emission tomography (PET)** is the detection of tumor tissue at whole-body studies performed with the glucose analogue 2-[fluorine-18]fluoro-2-deoxy-d-glucose (FDG). High rates of recurrence after partial hepatic resection in patients with colorectal cancer liver metastases indicate that current presurgical imaging strategies are failing to ...

Cited by: 63 Author: Ian A. Zealley, Stephen J. Skehan, John ...

Publish Year: 2001

### Performance of imaging modalities in diagnosis of liver ...

<https://onlinelibrary.wiley.com/doi/abs/10.1002/jmri.22010>

Dec 20, 2009 · Surgery of **liver metastases** can be effective, and the appropriate selection of surgical candidates relies first on imaging. Different techniques are available, but information on their relative performance is unclear. The aim of this overview is to assess the imaging modality performance in the diagnosis of colorectal cancer (CRC) **liver metastases**.

Cited by: 200 Author: Irene Floriani, Valter Torri, Eliana Rulli, Da...

Publish Year: 2010

### Dual-time point images of the liver with 18F-FDG PET/CT in ...

<https://www.sciencedirect.com/science/article/pii/S2253808912000523>

Dual-time point images of the liver with 18 F-FDG PET/CT in suspected recurrence from colorectal cancer 18 F-FDG PET/TC con imagen hepática ... imaging modalities in diagnosis of **liver metastases** from colorectal cancer: a systematic review ... fluorodeoxyglucose-positron emission tomography

**Name of Journal:** *World Journal of Hepatology*

**Manuscript NO:** 48690

**Manuscript Type:** CASE REPORT

**Fascioliasis presenting as colon cancer liver metastasis on <sup>18</sup>F-fluorodeoxyglucose positron emission tomography/computed tomography: A case report**

Akbulut S *et al.* Misdiagnosis and overtreatment of hepatic fasciola hepatica

Sami Akbulut, Egemen Ozdemir, Emine Samdanci, Selver Unsal, Murat Harputluoglu,  
Sezai Yilmaz

## Match Overview

1	<b>Internet</b> 46 words crawled on 29-Mar-2017 <a href="http://emedicine.medscape.com">emedicine.medscape.com</a>	2%
2	<b>Publications</b> 15 words KAYA, Muhsin, BEŞTAŞ, Remzi, GİRGİN, Sadullah, ÇİÇEK, Muttalip and KAPLAN, Mehmet Ali. "Increased anti-Echin...	1%
3	<b>Internet</b> 10 words crawled on 21-Mar-2013 <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	<1%



国内版

国际版

Fascioliasis presenting as colon cancer liver metastasis on 18F-fluorodeoxyglucose



All

Images

Videos

翻译成中文

关闭取词

29,000 Results

Any time ▾

### False-Positive Mediastinal Lymphadenopathy on 18F ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3251247>

**18F-fluorodeoxyglucose (FDG) positron emission tomography and computed tomography**

(integrated FDG PET/CT) has been used to **diagnose recurrence** and differentiate postoperative changes from **lymph node metastasis in colorectal cancer**, although its accuracy is questionable.

### [18F]fluorodeoxyglucose positron emission tomography ...

[https://www.researchgate.net/publication/10636495\\_18Ffluorodeoxyglucose\\_positron...](https://www.researchgate.net/publication/10636495_18Ffluorodeoxyglucose_positron...)

**[18F]fluorodeoxyglucose positron emission tomography** surveillance of **hepatic metastases** from **prostate cancer** following radiofrequency ablation: A case report Article in The American surgeon 69(7 ...

### 18F-fluorodeoxyglucose PET/CT findings of a solitary ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3544051>

Dec 28, 2012 · Primary hepatocarcinoma, **liver abscess**, and **metastasis** from the gastrointestinal tract were initially suspected. In order to make an accurate diagnosis and stage the lesion, the patient was referred for a whole body **positron emission tomography/computed tomography** (PET/CT) scan to identify other sites of involvement.

Cited by: 2

Author: Bo Pan, Cun-Shi Wang, Jian-Kui Han, Li...

Publish Year: 2012

### 18F-fluorodeoxyglucose positron emission tomography ...

<https://www.sciencedirect.com/science/article/pii/S1590865813000625>

**Positron emission tomography-computed tomography** may be more effective for the diagnosis of **bone metastasis** in the initial staging workup. Conversely, **bone scanning and positron emission tomography-computed tomography** may be similarly effective for the detection of **metachronous bone metastasis**.

Cited by: 6

Author: Dae Won Ma, Jie Hyun Kim, Tae Joo Je...

Publish Year: 2013

### Value of Positron Emission Tomography With [F-18 ...

[https://www.researchgate.net/publication/314015190\\_Value\\_of\\_Positron\\_Emission...](https://www.researchgate.net/publication/314015190_Value_of_Positron_Emission...)

**Positron Emission Tomography (PET)/Computed Tomography (CT)** is widely used in the work-up of



All

Images

Videos

翻译成中文

开启取词

27,900 Results

Any time ▾

## False-Positive Mediastinal Lymphadenopathy on 18F ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3251247>

18F-fluorodeoxyglucose (FDG) positron emission tomography and computed tomography (integrated FDG PET/CT) has been used to diagnose recurrence and differentiate postoperative changes from lymph node metastasis in colorectal cancer, although its accuracy is questionable.

## [18F]fluorodeoxyglucose positron emission tomography ...

[https://www.researchgate.net/publication/10636495\\_18Ffluorodeoxyglucose\\_positron...](https://www.researchgate.net/publication/10636495_18Ffluorodeoxyglucose_positron...)

[18F]fluorodeoxyglucose positron emission tomography surveillance of hepatic metastases from prostate cancer following radiofrequency ablation: A case report Article in The American surgeon 69(7 ...

## 18F-fluorodeoxyglucose PET/CT findings of a solitary ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3544051>

Dec 28, 2012 · Primary hepatocarcinoma, liver abscess, and metastasis from the gastrointestinal tract were initially suspected. In order to make an accurate diagnosis and stage the lesion, the patient was referred for a whole body positron emission tomography/computed tomography (PET/CT) scan to identify other sites of involvement.

Cited by: 2

Author: Bo Pan, Cun-Shi Wang, Jian-Kui Han, Lin...

Publish Year: 2012

## Value of Positron Emission Tomography With [F-18 ...

[https://www.researchgate.net/publication/314015190\\_Value\\_of\\_Positron\\_Emission...](https://www.researchgate.net/publication/314015190_Value_of_Positron_Emission...)

Positron Emission Tomography (PET)/Computed Tomography (CT) is widely used in the work-up of patients with various types of cancer. In our case, a patient with colon cancer ...

## Usefulness 18F-FDG positron emission ...

<https://aasldpubs.onlinelibrary.wiley.com/doi/full/10.1002/lt.22069>

May 25, 2010 · 18 F-fluoro-2-deoxy-D-glucose (18 F-FDG) positron emission tomography (PET)/computed tomography (CT) has been used at some institutions for clinical staging of HCC. The principle of 18 F-FDG PET is that 18 F-FDG uptake allows cellular glucose metabolism to ...

Cited by: 54

Author: Young-Kyu Kim, Kwang-Woong Lee, Seo...

Publish Year: 2010

## Management of potentially resectable colorectal cancer ...

<https://www.uptodate.com/contents/management-of-potentially-resectable-colorectal...>

The role of whole-body positron emission tomography with [18F]fluorodeoxyglucose in identifying operable colorectal cancer metastases to the liver. Arch Surg 1996; 131:703. Valk PE, Abella-Columa E, Haseman MK, et al. Whole-body PET imaging with [18F]fluorodeoxyglucose in management of recurrent colorectal cancer.

## Fasciola Hepatica: Missing Malincom on 18F