

Name of Journal: *World Journal of Hepatology*

Manuscript NO: 38394

Manuscript Type: Original Article

Retrospective Study

Paracentesis in cirrhotics is associated with increased risk of 30-d readmission

Sobotka LA *et al.* Paracentesis in cirrhotics and readmission risk

Lindsay A Sobotka, Rohan M Modi, Akshay Vijayaraman, A James Hanje, Anthony J

Michaels, Lanla F Conteh, Alice Hinton, Ashraf El-Hinnawi, Khalid Mumtaz

Match Overview

1	Internet 42 words crawled on 18-Jan-2018 www.wjgnet.com	1%
2	Internet 30 words crawled on 12-Sep-2017 scholarworks.iupui.edu	1%
3	Internet 15 words crawled on 18-Nov-2017 www.frontiersin.org	<1%
4	Crossref 15 words Desai, Archita P., and Nancy Reau. "The Burden of Rehospitalization for Patients With Liver Cirrhosis." <i>Journal of Hepatology</i> 64, no. 1 (2016): 105-112.	<1%
5	Publications 15 words "Studies from State University of New York Research Institute for Liver and Gastrointestinal New Findings on Breast Cancer (Report)" <i>World Journal of Hepatology</i> 6, no. 1 (2014): 1-10.	<1%
6	Crossref 14 words McDaniel, John, Gangarao Davuluri, Elizabeth Ann Hill, Michelle Moyer, Ashok Runkana, Rich	<1%
7	Internet 13 words crawled on 02-Jul-2017 linknovate.com	<1%
8	Crossref 12 words Mitchell Ramsey, Somashekar G Krishna, Peter P Stanich, Sved Husain, Edward J Levine, Darw	<1%
9	Crossref 11 words Lindsay, Aaron, James Burton, and Charles Ray. "Paracentesis-Induced Circulatory Dysfunction	<1%
10	Internet 11 words crawled on 25-Feb-2014 www.ncbi.nlm.nih.gov	<1%



All

Images

News

Shopping

Videos

More

Settings

Tools

About 69,700 results (0.49 seconds)

P584 - Paracentesis Is Associated With Increased Risk of 30-Day ...

<https://www.eventscribe.com/2017/wcogacg2017/ajaxcalls/PosterInfo.asp?efp...> ▼

P584 - Paracentesis Is Associated With Increased Risk of 30-Day Readmission in Patients With Cirrhosis and Ascites. Sunday, Oct 15. 3:30 PM – 7:00 PM. Category: Liver. Lindsay Sobotka, DO¹, Akshay Vijayaraman, MD¹, Rohan M. Modi, MD¹, Alice Hinton, PhD², Lanla F. Conteh, MD, MPH¹, Anthony J. Michaels, MD¹, ...

American Journal of Gastroenterology - Could Adherence to Quality of ...

www.nature.com › [Journal home](#) › [Archive](#) › [Original Contributions](#)

Jan 5, 2016 - **Patients** who received an abdominal **paracentesis** within **30 days** of **ascites** diagnosis (QI 1, relative **risk** (RR) 0.41, P=0.004) or during index hospitalization (QI 2, RR 0.57, ... Baseline serum bilirubin >2.5 mg/dl was **associated with increased 30-day cirrhosis-related readmission** (RR 1.51, P=0.03). A total of ...

Ascites, refractory ascites and hyponatremia in cirrhosis - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5421465/>

by B Fortune - 2017 - [Cited by 2](#) - [Related articles](#)

Apr 24, 2017 - The development of **ascites**, as mentioned previously, is **associated** with significant decline in morbidity and mortality. The 1-year mortality **risk** for **patients with cirrhosis increases** from 1–3% to 15–20% after the development of **ascites** and as **high** as 50% in 5 years [5,25]. Liver transplantation is the only ...

Cirrhosis and its complications: Evidence based treatment - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4017060/> ▼

by S Nusrat - 2014 - [Cited by 59](#) - [Related articles](#)

May 14, 2014 - **Patients with cirrhosis** are at **increased risk** of numerous complications and have a decreased life expectancy[3]. The American Association of Liver Disease (AASLD) recommends

全部

新闻

图片

购物

视频

更多

设置

工具

已启用安全搜索

找到约 53,600 条结果 (用时 0.63 秒)

您是不是要找：**Paracentesis in cirrhosis is associated with increased risk of 30-d readmission**

Google 学术：**Paracentesis in cirrhotics is associated with increased risk of 30-d readmission**

Complications associated with liver transplantation in ... - LaMattina - 被引用次数：63

Alcoholic hepatitis 2010: a clinician's guide to ... - Amini - 被引用次数：69

Hospital readmissions in decompensated cirrhotics: Factors pointing ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5645619/> - 翻译此页

作者：SM Seraj - 2017 - 被引用次数：2 - 相关文章

2017年10月7日 - Readmissions are challenging aspect of care for cirrhotic patients and risk continues beyond 30 d. ... fluid without evidence for an intra-abdominal secondary source such as a perforated viscus; (4) Volume related, defined as ascites or hepatic hydrothorax requiring paracentesis during the hospitalization.

Paracentesis is Associated with Reduced Mortality in Patients ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3944409/> - 翻译此页

作者：ES Orman - 2014 - 被引用次数：33 - 相关文章

2013年8月24日 - Ascites is the most common complication of cirrhosis, and its development is associated with substantially increased mortality. One of the most ... SBP is present in 10–30% of all hospitalized patients with ascites, and the risk of complications from diagnostic paracentesis is negligible. Therefore, for more ...

The benefit of paracentesis on hospitalized adults with cirrhosis and ...

<https://www.ncbi.nlm.nih.gov/pubmed/26642977> - 翻译此页

作者：M Cozzani - 2016 - 被引用次数：4 - 相关文章

[全部](#)[新闻](#)[图片](#)[购物](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 275,000 条结果 (用时 0.64 秒)

您是不是要找: [Paracentesis in *cirrhosis* is associated with increased risk of 30-d readmission](#)

[Hospital readmissions in decompensated cirrhotics: Factors pointing ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5645619/> - 翻译此页

作者: SM Seraj - 2017 - 被引用次数: 2 - 相关文章

2017年10月7日 - Readmissions are challenging aspect of care for cirrhotic patients and risk continues beyond 30 d. fluid without evidence for an intra-abdominal secondary source such as a perforated viscus; (4) Volume related, defined as ascites or hepatic hydrothorax requiring paracentesis during the hospitalization.

[Paracentesis is Associated with Reduced Mortality in Patients ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3944409/> - 翻译此页

作者: ES Orman - 2014 - 被引用次数: 33 - 相关文章

2013年8月24日 - Ascites is the most common complication of cirrhosis, and its development is associated with substantially increased mortality. One of the most ... SBP is present in 10–30% of all hospitalized patients with ascites, and the risk of complications from diagnostic paracentesis is negligible. Therefore, for more ...

[The benefit of paracentesis on hospitalized adults with cirrhosis and ...](#)

<https://www.ncbi.nlm.nih.gov/pubmed/26642977> - 翻译此页

作者: JN Gaetano - 2016 - 被引用次数: 1 - 相关文章

J Gastroenterol Hepatol. 2016 May;31(5):1025-30. doi: 10.1111/jgh.13255. The benefit of paracentesis on hospitalized adults with cirrhosis and ascites. Gaetano JN(1), Micic D(1), Aronsohn A(1), Reddy G(1), Te H(1), Reau NS(2), Jensen D(1). Author information: (1)University of Chicago Medicine, Department of Medicine, ...