

## FoxO3a and disease progression

Richard Seonghun Nho, Polla Hergert

Richard Seonghun Nho, Polla Hergert, Division of Pulmonary, Allergy and Critical Care Medicine, Department of Medicine, University of Minnesota, Minneapolis, MN 55455, United States  
Author contributions: Nho RS and Hergert P both contributed to this paper.

Supported by the National Institutes of Health R01 HL 114662 to Nho R  
Correspondence to: Richard Seonghun Nho, PhD, Division of Pulmonary, Allergy and Critical Care Medicine, Department of Medicine, University of Minnesota, Box 276, 420 Delaware Street SE., Minneapolis, Minnesota, MN 55455, United States. [nhox002@umn.edu](mailto:nhox002@umn.edu)

Telephone: +1-612-6250686 Fax: +1-612-6252174

Received: November 29, 2013 Revised: April 21, 2014

Accepted: May 16, 2014

Published online: August 26, 2014

highlighted as a critical protein that regulates numerous cell functions from proliferation/apoptosis to stress-resistance and aging. FoxO3a has been found to be deregulated in several diseases and FoxO3a targeting approaches are currently underway to treat various types of cancers. This review will describe the current concept of FoxO3a's pathological role in various diseases and elucidate the regulatory mechanisms involved. It will also provide the clinical significance and strategies to target FoxO3a to limit the progression of human diseases.

Nho RS, Hergert P. FoxO3a and disease progression. *World J Biol Chem* 2014; 5(3): 346-354 Available from: URL: <http://www.wjgnet.com/1949-8454/full/v5/i3/346.htm> DOI: <http://dx.doi.org/10.4331/wjbc.v5.i3.346>

### Abstract

The Forkhead box O (FoxO) family has recently been highlighted as an important transcriptional regulator of crucial proteins associated with the many diverse functions of cells. So far, FoxO1, FoxO3a, FoxO4 and FoxO6 proteins have been identified in humans. Although each FoxO family member has its own role, unlike the other FoxO families, FoxO3a has been extensively studied because of its rather unique and pivotal regulation of cell proliferation, apoptosis, metabolism, stress management and longevity. FoxO3a alteration is closely linked to the progression of several types of cancers, fibrosis and other types of diseases. In this review, we will examine the function of FoxO3a in disease progression and also explore FoxO3a's regulatory mechanisms. We will also discuss FoxO3a as a potential target for the treatment of several types of disease.


© 2014 Baishideng Publishing Group Inc. All rights reserved.

**Key words:** Forkhead box O; Cell proliferation; Apoptosis; Stress; Aging

**Core tip:** Forkhead box O (FoxO)3a has recently been

### INTRODUCTION

Forkhead box O (FoxO) transcription factors are the human homologues of the *C. elegans* transcription factor DAF-16 and share a highly conserved 110-amino acid DNA binding domain, forkhead box or winged-helix domain<sup>[1,2]</sup>. Forkhead box proteins comprise more than 100 members in humans, classified from FOXA to FOXR<sup>[3-5]</sup>. Members of class O share the characteristic of being regulated by the insulin/PI3K/Akt signaling pathway<sup>[4]</sup>. Four principal members of the mammalian FoxO subfamily, FoxO1, FoxO3a, FoxO4 and FoxO6 have been previously described<sup>[3]</sup>. Although they seem to bind a common set of DNA sites, FoxO6 is mainly specific to neurons while the other 3 FoxO family members are expressed in most tissues. These FoxO members are linked to cell survival, cellular proliferation and DNA damage repair response<sup>[5,6]</sup>. Among them, FoxO3a has recently been studied extensively as a crucial protein that is involved in the regulation of several essential cellular functions (see page 349). Prior studies have shown that FoxO3a functions as a tumor suppressor by regulating expression of genes in-



**Figure 1 Forkhead box O3a target genes.** Forkhead box O (FoxO)3a transcriptionally activates several target genes. FoxO3a binds to the promoter of apoptosis inducing genes, such as *Bim*, *FasL* and *TRAIL*, and to the promoter of cell cycle inhibitors, such as p27 and p21. FoxO3a also activates autophagy genes *Gabarrap1*, *ATG12*, etc. A recent study showed that FoxO3a also participates in the activation of stress response genes, such as MnSOD and catalase in response to oxidative stress.

involved in apoptosis, cell cycle arrest, oxidative stress resistance and autophagy<sup>[3,7-9]</sup> (Figure 1). In general, FoxO3a is known to suppress cell cycle progression and promote cell death. Thus, it has been thought that FoxO3a can be an important target to inhibit cancer cell progression. However, recent studies have discovered other functions of FoxO3a, such as stress response and longevity, as described on page 349. FoxO3a alteration is also linked to many different types of disease. Interestingly, FoxO3a increases autophagy to protect cells from environmental stresses<sup>[10,11]</sup>. Thus, under this situation, unlike the general concept of FoxO3a's role, FoxO3a potentially has a protective role in maintaining a cell's homeostasis. Perhaps the most interesting feature of FoxO3a is its biological role associated with longevity (page 349). Based on this, it becomes clear that FoxO3a has diverse roles in response to many environmental stimuli and these recent findings certainly change our view on the previous roles of FoxO3a. Therefore, from the perspective of disease progression, it is imperative to define the potential role of FoxO3a in cells and elucidate how alteration of FoxO3a is linked to the development of several types of disease.


## FOXO3A STRUCTURE

Recent technologies have revealed that the primary structure of FoxO3a contains highly conserved residues of the helix H3 (motif NXXRHXXS/T), which is the main DNA recognition element that binds into a major groove, which comprises the majority of the direct base-specific contacts<sup>[1,6]</sup>. Recent studies further revealed that FoxO proteins recognize two consensus sequences, 5'-GTAAA(T/C)AA-3' known as the Daf-16 family member-binding element<sup>[6,7]</sup> and 5'-(C/A)(A/C)AAA(C/T)AA-3' known as the insulin-responsive sequence (IRE)<sup>[8,9]</sup>. Crystal structure revealed that the recognition helix H3 docked perpendicular to the major groove making extensive contacts with the DNA<sup>[7]</sup>. FoxO3a contains several crucial domains<sup>[12]</sup> (Figure 2) such as a nuclear localization signal (NLS), a nuclear export signal (NES) and a transactivation domain (TA).


## FOXO3A REGULATORY MECHANISMS

### Phosphorylation and dephosphorylation

FoxO3a is regulated by posttranslational modifications such as phosphorylation, acetylation and ubiquitination, each of which affects the transcriptional activity of FoxO proteins<sup>[11-16]</sup> (Figure 2). The potency of FoxO3a is carefully regulated by phosphorylation. The phosphorylation of FoxO3a by several kinases is well established. Among them, protein kinase B (Akt) is an important kinase that directly phosphorylates FoxOs. In the case of FoxO3a, T32, S253 and S315 residues are phosphorylated by Akt and, in particular, the phosphorylation of S253 is a crucial residue regulating the nuclear/cytoplasmic shuttling of FoxO3a. For example, when cells are cultured in the presence of growth factors or insulin, FoxO3a is phosphorylated by Akt and mainly localized to the cytoplasm, which prevents its transcriptional activity. The phosphorylation event of FoxO3a by Akt facilitates FoxO3a interaction with the 14-3-3 nuclear export protein, further preventing nuclear re-import by concealing nuclear localization signals<sup>[13]</sup>. Furthermore, the phosphorylation of FoxO3a by activated Akt promotes an association with an ubiquitin E3 ligase, subsequently polyubiquitinating FoxO3a, which facilitates FoxO3a degradation by proteasomes<sup>[13-17]</sup>. Thus, the activation of Akt is thought to be critical in FoxO3a regulation. However, in some tumors, FoxO3a remains in the cytoplasm even in the absence of active Akt<sup>[14]</sup>. It has been found that IκB kinase (IKK) phosphorylates FoxO3a at serine 644, thereby inhibiting its transcriptional activity in an Akt-independent manner<sup>[15]</sup>. The phosphorylation of FoxO3a by IKK also leads to its cytoplasmic localization, although the underlying export mechanism is not understood. The insulin/IGF-1 and integrin-dependent signaling pathways activate Akt *via* PTEN suppression which phosphorylates FoxO3a, thereby rendering it functionally inactive. In contrast, FoxO3a is localized to the nucleus to activate its target genes when growth factors or serum are deprived. Additionally, serum and glucocorticoid regulated kinase (SGK), casein kinase 1 (CK 1), dual specificity tyrosine-phosphorylated and regulated kinase 1A (DYRK 1A), janus N-terminal kinase (JNK), mitogen-activated protein kinases (MAPKs), IκappaB kinase (IKKβ), mammalian sterile 20-like kinase 1 (MST1) and AMP activated protein kinase (AMPK) are also known to regulate FoxO3a and other family members<sup>[18-23]</sup> by phosphorylating multiple residues. Interestingly, SGK 1 is transcriptionally up-regulated in response to a variety of external stimuli, including growth factors. SGK 1 is also known to phosphorylate the pivotal ser 253 residue, which triggers its location to the cytoplasm, thereby inhibiting its function<sup>[19]</sup>. In contrast, AMPK activates FoxO3a function. 6 threonine/serine residues (T179, S309, S413, S555, S588 and S626) in mammalian FoxO3a are found to be phosphorylated by AMPK<sup>[24,25]</sup>. Mutation of these phosphorylation residues to alanine severely impairs its function, yet it does not alter its ability to bind to cognate



**Figure 2 Major phosphorylation and acetylation residues of FoxO3a.** Post-translational modification sites of FoxO3a. Shown are sites of serine/threonine phosphorylation by Akt/SGK, MST1, IKKβ or the residues acetylated by CBP or unidentified acetyl transferases (?) on FoxO3a domains<sup>[12]</sup>. FKH: Forkhead DNA binding domain; NLS: Nuclear localization signal; NES: Nuclear export sequence; TA: Transactivation domain; Akt: Protein kinase B; MST1: Mammalian sterile 20 like kinase-1; CBP: The cyclic-AMP responsive element binding (CREB) binding protein, IKKβ: Iκβ kinase; SGK: Serum-and glucocorticoid-induced protein kinase.



**Figure 3 Forkhead box O3a localization by phosphorylation and dephosphorylation.** Forkhead box O (FoxO)3a becomes translocated to the cytoplasm when phosphorylated on ser 253 residue by Akt or SGK. FoxO3a is then bound to 14-3-3 and this interaction promotes its degradation by the proteasome. In contrast, FoxO3a is dephosphorylated by protein phosphatase-2A and this opposite event facilitates its relocation into the nucleus, thereby activating its target genes. SGK: Serum-and glucocorticoid-induced protein kinase; Akt: Protein kinase B.

sequences or to participate in nucleocytoplasmic shuttling depending on external cues<sup>[25]</sup>. Likewise, JNK also phosphorylates FoxO3a, activating FoxO3a function by enhancing its location into the nucleus which subsequently increases its transcriptional activity<sup>[18,22]</sup>.

Unlike kinases, very few phosphatases have been found to regulate FoxO3a. One particular phosphatase, protein phosphatase-2A (PP2A), has been shown to regulate FoxO3a function. Nho *et al.*<sup>[26]</sup> showed that when fibroblasts attach to 2D type collagen coated plates, PP2A activity is suppressed, which facilitates FoxO3a inactivation by enhanced Akt, promoting fibroblast proliferation. But the over-expression of PP2A reverses this inactivation and increases dephosphorylated FoxO3a, thereby suppressing their proliferation. Singh *et al.*<sup>[24]</sup> also demonstrated that FoxO3a interacts with PP2A C/A subunits in HeLa cells, dephosphorylating its T32/S253 residues, which subsequently inhibits the interaction of the 14-3-3 protein to FoxO3a by Akt. This study showed that PP2A is required for the reactivation of FoxO3a by promoting its translocation to the nucleus (Figure 3). Interestingly, recent studies also showed that the adenovirus E1A stabilizes FoxO3a by inducing the expression of PP2A/C, which suppresses βTrCP-mediated degradation of FoxO3a<sup>[25]</sup>. Thus, these studies clearly suggest that the imbalance between kinases and phosphatase(s) can greatly affect a cell's fate by curbing FoxO3a function and the alteration of these kinases and phosphatases are directly linked to certain disease progression.

**Ubiquitin proteasome degradation**

As we briefly described above, FoxO3a degradation is

also an important step to regulate its function. The single molecule RING-finger E3 ligase murine double minute 2 (MDM2) promotes ubiquitination of FoxO3a as well as FoxO1 and FoxO4, facilitating their degradation<sup>[27]</sup>. Intriguingly, knockout or knockdown of MDM2 alone increases FoxO3a protein levels. This effect was shown to be mediated by MDM2-induced polyubiquitination of FoxO proteins<sup>[27,28]</sup>, whereas another study showed that MDM2 catalyzes multiple monoubiquitination of FoxO4 rather than polyubiquitination<sup>[28]</sup>. When FoxO3a is located to the cytoplasm by Akt, FoxO3a becomes ubiquitinated and this event triggers a proteasome-dependent degradation process. Like MDM2, FoxO3a phosphorylation by IKK also leads to its ubiquitination and degradation<sup>[15]</sup>. Thus, these studies document that FoxO3a localization in the cytoplasm not only deactivates FoxO3a function but also becomes a crucial step leading to FoxO3a degradation.

**Acetylation, transcriptional regulation, microRNA and others**

Acetylation also plays an important role in regulating FoxO3a. Oxidative stress triggers FoxO3a acetylation/deacetylation and affects the localization of FoxO3a. For example, protein acetylase CREB binding protein (CBP)<sup>[29-31]</sup>, p300<sup>[32,33]</sup> and deacetylase Sirt are known to modulate FoxO3a function<sup>[34-36]</sup>, although a precise mechanism describing the effects of acetylation and deacetylation is not known. A recent piece of evidence suggests that the FoxO family is also regulated by microRNA. mir155, mir96 and mir21 are thought to directly regulate FoxO3a, while mir205 regulates FoxO3a *via* its



upstream target PTEN<sup>[39-43]</sup>. FoxO3a is also known to be regulated by a transcription factor. E2F-1 can bind to the promoter region of FoxO1 and FoxO3a, thereby regulating FoxO3a at the mRNA level<sup>[44]</sup>. FoxO3a mRNAs are modulated as a function of age in rat muscle, peaking at 6 and 23 mo, suggesting that FoxO3a may also affect longevity in mammals<sup>[45]</sup>.

## FOXO3A FUNCTION

### Cell proliferation and apoptosis

Perhaps the two most significant cellular processes that are regulated by FoxO transcription factor are the suppression of cell cycle progression and the promotion of apoptosis<sup>[46-50]</sup>. FoxO3a activation increases cell cycle inhibitor proteins p21 and p27, both of which subsequently suppress G1 to S cell cycle transition<sup>[51-54]</sup>. Although p27 is transcriptionally regulated by FoxO3a *via* the PI3K/Akt-dependent axis, it has been shown that p27 is also regulated *via* the FoxO3a/NF- $\kappa$ B/c-Myc-dependent pathway. Chandramohan *et al.*<sup>[55]</sup> showed that in WEHI 231 cells, the suppression of PI3K activity promotes a decrease in c-Myc dependent p27 expression *via* NF- $\kappa$ B inhibition. Since NF- $\kappa$ B is frequently altered in many types of cancers and NF- $\kappa$ B transcriptionally activates *c-Myc* gene expression, this finding suggests that p27 is reciprocally regulated by FoxO3a and c-Myc. A recent study further suggests that FoxO3a inhibits NF- $\kappa$ B function and that the alteration of FoxO3a is associated with hyper-proliferative helper T cells, cigarette smoke-induced inflammation, airspace enlargement and chronic obstructive pulmonary disease<sup>[56,57]</sup>. Likewise, FoxO3a also increases several target genes, such as Bim, TRAIL, PUMA and Fas ligand, which all promote cell apoptosis. For example, FoxO3a directly binds to the promoter region of Bim, causing sympathetic neuron cell death<sup>[44]</sup>. The activation of the transcription factor FoxO3a led to increased TRAIL transcription and induction of G1 arrest in the absence of v-Abl inhibition; this effect could be inhibited by the expression of a constitutively active Akt mutant in BCR-Abl-transformed human cells. Ghaffari *et al.*<sup>[49]</sup> also demonstrated that cytokine and BCR-Abl suppression of TRAIL transcription is mediated through phosphorylation and inhibition of the FoxO3a transcription factor. This study showed that BCR-Abl-induced inhibition of TRAIL transcription is linked to the tumorigenicity in chronic myeloid leukemia<sup>[50]</sup>. FoxO3a is also associated with the regulation of PUMA and Noxa proteins in lymphoid and neuroblastoma cells, respectively<sup>[58,59]</sup>. Thus, these findings clearly demonstrate that FoxO3a-dependent cell cycle arrest and apoptosis induction are important for tumor suppression (Table 1) and further indicate that the pathological alteration of FoxO3a can potentially contribute to the acquisition of uncontrolled cell proliferation and an apoptosis-resistant cell phenotype.

### Stress resistant effect

The most recent discovery regarding FoxO3a's function

is that it is also associated with stress response and longevity. In contrast to FoxO3a's better known functions of inhibiting cell proliferation and promoting apoptosis as described above, FoxO3a also participates in protecting cells when exposed to unfavorable conditions. This seemingly contradictory effect of FoxO3a has been observed in various cell models and it has been found that the reactive oxygen species (ROS) are linked to the activation of FoxO3a to protect cells from a stress inducing environment<sup>[60,61]</sup>. In *C. elegans*, DAF-16 is thought to regulate 230 genes on the ablated germ cell line background and most of these genes are related to the resistance of external stress<sup>[62,63]</sup>. Deregulated ROS induce apoptosis and are associated with various diseases and aging. Sirtuin-1 (Sirt1) decreases ROS levels and promotes cell survival under oxidative stress conditions. Interestingly, FoxO3a and other FoxO family members increase superoxide dismutase (SOD) and protect cells from oxidative stress in a Sirt1-dependent manner<sup>[34,38]</sup>. A Sirt1/FoxO3a-dependent cell regulatory function that has been linked to stress management was previously studied. Brunet *et al.* showed that Sirt1 and FoxO3a form a complex in cells in response to oxidative stress and Sirt1 increases the ability of FoxO3a to induce cell cycle arrest and resistance to oxidative stress but inhibited FoxO3a's function to induce cell death<sup>[38]</sup>. These results showed that FoxO3a deacetylation by Sirt1 in response to ROS can be an important self defense mechanism to detoxifying harmful reactive molecules, further suggesting that Sirt1 is linked to protect cells from a stress inducing environment by tipping FoxO dependent response away from apoptosis and toward stress resistance<sup>[38]</sup>. Studies also found that Sirt3 which belongs to class III of HDACs, is linked to the resistance of stress inducing environments by detoxifying ROS. The role of Sirt3 and FoxO3a function is particularly well described in myocytes<sup>[64]</sup>. At the cellular level, when cardiomyocytes are exposed to stressful stimuli, Sirt3 levels are elevated, which subsequently deacetylate FoxO3a and facilitate its location into the nucleus to activate anti-oxidant genes<sup>[65]</sup>. Among them, catalase (Cat) and manganese superoxide dismutase (MnSOD) are direct targets of detoxifying enzymes by FoxO3a. Thus, the increased level of Cat and MnSOD by FoxO3a activation may efficiently and effectively manage ROS, which can be beneficial for reducing stress induced by ROS. Interestingly, a prior study found a potential FoxO activator as a way to protect cells from oxidative stress. Resveratrol, a polyphenolic flavonoid abundant in red wine with potent antioxidant activity, is known to up-regulate the FoxO family and block caspase 3, 8, and 9 activation, protecting photoreceptor cells from oxidative stress<sup>[66]</sup>. Thus, it is believed that when cells are exposed to a stress inducing environment, FoxO3a protects cells by utilizing SOD, catalase, *etc.*, and this action is ultimately beneficial to cells. Given the fact that FoxO3a is linked to stress response and cells utilize FoxO3a to respond to ROS, it is a plausible scenario that the activation of FoxO3a under stress inducing conditions triggers the cell's defense sys-

**Table 1 FoxO3a target genes in various cell types**

FoxO3a target genes	Cell types
<i>Bim</i>	Neuron cells <sup>[48]</sup>
<i>TRAIL</i>	Bcr/Abl transformed cells <sup>[57]</sup>
<i>TRAIL</i>	Chronic myeloid leukemia <sup>[46]</sup>
<i>PUMA</i>	Lymphoid cells <sup>[58]</sup>
<i>Noxa</i>	Neuroblastoma <sup>[59]</sup>
<i>FasL</i>	Glomerular mesangial cells <sup>[102]</sup>
<i>p27, Caveolin-1</i>	Glomerular mesangial cells <sup>[102]</sup>
<i>p21</i>	Glomerular mesangial cells <sup>[102]</sup>

Shown are previously known FoxO3a target genes that regulate cell proliferation and apoptosis in different cell types.

tem, which can protect cells from harmful environments.

### Longevity

However, perhaps the most intriguing recent discovery in FoxO3a function is that the *FoxO3a* gene is associated with aging. Because FoxO3a is regulated by insulin-IGF1 signaling (IIS) which influences metabolism and lifespan in model organisms<sup>[67]</sup>, FoxO3a had been proposed to be an ideal candidate to study longevity as the link between FoxO3a and longevity that has previously been described. Willcox *et al.*<sup>[68]</sup> described 3 single nucleotide polymorphisms (SNPs) in the *FoxO3a* gene that were statistically significantly associated with longevity and different aging phenotypes in a sample of long-lived Americans of Japanese ancestry. Furthermore, Flachsbart *et al.*<sup>[69]</sup> found that not only were certain FoxO3a variants very common in 90 year olds, they were even more common in 100 year olds, emphasizing the importance of genetics for aging well. It becomes clear that increases in cellular ROS levels are known to be associated with aging<sup>[70-75]</sup>. Increased cellular oxidative stress regulates FoxO post-translational modifications and the activation of the FoxO family has been shown to regulate cellular oxidative-stress resistance<sup>[76-81]</sup>. Interestingly, to support these findings, recent studies suggest a possibility that Sirt3 and FoxO3a have been linked to an extended life span in humans<sup>[75-78,82]</sup>.

### FOXO3A IN CLINICAL APPLICATION

FoxO3, FoxO1 and FoxO4 are present at chromosomal translocation break points in cells of rhabdomyosarcomas and acute myeloid leukemia. Among the FoxO family, FoxO3a has been shown to be deregulated in several tumor types, including breast cancer<sup>[83-85]</sup>, prostate cancer<sup>[86-88]</sup>, glioblastoma<sup>[89]</sup> and leukemia<sup>[90,91]</sup>. Therefore, FoxO3a has been targeted as a way to treat several types of cancers. Interestingly, Akt, IKK and Erk are three commonly activated oncogenic kinases in human cancers and all three kinases target FoxO3a in an identical manner to inhibit its tumor suppressor function<sup>[92]</sup>. All three kinase-mediated phosphorylations stimulate FoxO3a ubiquitination, resulting in its proteasomal degradation. Thus, a FoxO3a targeting approach *via* the modulation of above kinases is currently underway. For example,

the chemotherapeutic drugs paclitaxel<sup>[93]</sup> and KP372-1 (a multiple kinase inhibitor)<sup>[30]</sup>, currently used in the treatment of breast carcinoma, activate FoxO3a by reducing Akt activity. Doxorubicin activates FoxO3a to induce the expression of the multidrug resistance gene ABCB1 (MDR1) in K562 doxorubicin-sensitive leukemic cells<sup>[94]</sup>. Imatinib activates FoxO3a and induces Bim-dependent apoptosis through inhibition of BCR-ABL in chronic myeloid leukemia<sup>[95]</sup>. Imatinib also induces erythroid differentiation through repressing ID1 gene transcription by FoxO3a activation<sup>[96]</sup>. BMS-345541, a selective IKK inhibitor, promotes apoptosis in T-cell acute lymphoblastic leukemia (T-ALL) cell lines<sup>[97]</sup>. Several pieces of evidence in recent years further suggest that a FoxO3a targeting approach may be helpful for the treatment of other types of human diseases. For example, FoxO3a causes the induction of apoptosis in prostate cancer cells *via* up-regulating PUMA<sup>[98]</sup>. Low levels of FoxO3a may link to chemotherapy resistance in liver cancer and FoxO3a appears to present antitumor properties in hepatocellular carcinoma<sup>[99-101]</sup>. FoxO3a also plays a role in the neuroprotective effect of the erythropoietin (EPO) role in Parkinson's disease *via* Akt<sup>[102]</sup>. Thus, all these studies indicate that as our knowledge for FoxO3a targeting approaches continuously develop, the clinical application of FoxO3 is potentially promising to limit the progression of human diseases in the future.

### FUTURE APPLICATION OF FOXO3A

FoxO3a has recently been recognized as a promising therapeutic target to treat cancers and other types of diseases. To improve therapeutic outcomes, FoxO3a-dependent chemosensitization is being currently tested. Studies suggest that precise FoxO3a regulation is essential for homeostasis and if there is deregulation of FoxO3a by environmental factors, such as chronic exposure to ROS or genetic/epigenetic alteration, this pathological condition can directly lead to abnormal proliferation or changes in apoptotic signals, which subsequently are responsible for disease progression. In particular, age-dependent FoxO3a modulation is an interesting concept to help understand the pathogenesis of certain types of disease models. If FoxO3a is a crucial protein mainly deregulated by aging, maintaining optimum FoxO3a activity in a patient's specific clinical condition can be beneficial to minimize age-dependent disease. For example, the preservation of optimum FoxO3a activity using drugs such as paclitaxel may be helpful for patients with age-related diseases. Clearly, more studies are required to elucidate FoxO3a's function as an effective and useful target capable of preventing or limiting the progression of diseases without clinical compromise.

### REFERENCES

1 Clark KL, Halay ED, Lai E, Burley SK. Co-crystal structure of the HNF-3/fork head DNA-recognition motif resembles histone H5. *Nature* 1993; **364**: 412-420 [PMID: 8332212 DOI:

- 10.1038/364412a0]
- 2 **Arden KC.** Multiple roles of FOXO transcription factors in mammalian cells point to multiple roles in cancer. *Exp Gerontol* 2006; **41**: 709-717 [PMID: 16806782 DOI: 10.1016/j.exger.2006.05.015]
  - 3 **Huang H, Tindall DJ.** Dynamic FoxO transcription factors. *J Cell Sci* 2007; **120**: 2479-2487 [PMID: 17646672 DOI: 10.1242/jcs.001222]
  - 4 **Dillin A, Crawford DK, Kenyon C.** Timing requirements for insulin/IGF-1 signaling in *C. elegans*. *Science* 2002; **298**: 830-834 [PMID: 12399591 DOI: 10.1126/science.1074240]
  - 5 **Monsalve M, Olmos Y.** The complex biology of FOXO. *Curr Drug Targets* 2011; **12**: 1322-1350 [PMID: 21443460 DOI: 10.2174/138945011796150307]
  - 6 **Biggs WH, Cavenee WK, Arden KC.** Identification and characterization of members of the FKHR (FOX O) subclass of winged-helix transcription factors in the mouse. *Mamm Genome* 2001; **12**: 416-425 [PMID: 11353388 DOI: 10.1007/s003350020002]
  - 7 **Furuyama T, Nakazawa T, Nakano I, Mori N.** Identification of the differential distribution patterns of mRNAs and consensus binding sequences for mouse DAF-16 homologues. *Biochem J* 2000; **349**: 629-634 [PMID: 10880363 DOI: 10.1042/0264-6021.]
  - 8 **Hall RK, Yamasaki T, Kucera T, Waltner-Law M, O'Brien R, Granner DK.** Regulation of phosphoenolpyruvate carboxykinase and insulin-like growth factor-binding protein-1 gene expression by insulin. The role of winged helix/forkhead proteins. *J Biol Chem* 2000; **275**: 30169-30175 [PMID: 10913147 DOI: 10.1074/jbc.M004898200]
  - 9 **Hribal ML, Nakae J, Kitamura T, Shutter JR, Accili D.** Regulation of insulin-like growth factor-dependent myoblast differentiation by Foxo forkhead transcription factors. *J Cell Biol* 2003; **162**: 535-541 [PMID: 12925703 DOI: 10.1083/jcb.200212107]
  - 10 **Warr MR, Binnewies M, Flach J, Reynaud D, Garg T, Malhotra R, Debnath J, Passegué E.** FOXO3A directs a protective autophagy program in haematopoietic stem cells. *Nature* 2013; **494**: 323-327 [PMID: 23389440 DOI: 10.1038/nature11895]
  - 11 **Xia N, Strand S, Schlufter F, Siuda D, Reifenberg G, Kleintert H, Förstermann U, Li H.** Role of SIRT1 and FOXO factors in eNOS transcriptional activation by resveratrol. *Nitric Oxide* 2013; **32**: 29-35 [PMID: 23583951 DOI: 10.1016/j.niox.2013.04.001]
  - 12 **Greer EL, Brunet A.** FOXO transcription factors at the interface between longevity and tumor suppression. *Oncogene* 2005; **24**: 7410-7425 [PMID: 16288288 DOI: 10.1038/sj.onc.1209086]
  - 13 **Plas DR, Thompson CB.** Akt activation promotes degradation of tuberin and FOXO3a via the proteasome. *J Biol Chem* 2003; **278**: 12361-12366 [PMID: 12517744 DOI: 10.1074/jbc.M213069200]
  - 14 **Brunet A, Park J, Tran H, Hu LS, Hemmings BA, Greenberg ME.** Protein kinase SGK mediates survival signals by phosphorylating the forkhead transcription factor FKHL1 (FOXO3a). *Mol Cell Biol* 2001; **21**: 952-965 [PMID: 11154281 DOI: 10.1128/MCB.21.3.952-965.2001]
  - 15 **Hu MC, Lee DF, Xia W, Golfman LS, Ou-Yang F, Yang JY, Zou Y, Bao S, Hanada N, Saso H, Kobayashi R, Hung MC.** IkappaB kinase promotes tumorigenesis through inhibition of forkhead FOXO3a. *Cell* 2004; **117**: 225-237 [PMID: 15084260 DOI: 10.1016/S0092-8674(04)00302-2]
  - 16 **Vogt PK, Jiang H, Aoki M.** Triple layer control: phosphorylation, acetylation and ubiquitination of FOXO proteins. *Cell Cycle* 2005; **4**: 908-913 [PMID: 15917664 DOI: 10.4161/cc.4.7.1796]
  - 17 **Fu W, Ma Q, Chen L, Li P, Zhang M, Ramamoorthy S, Nawaz Z, Shimojima T, Wang H, Yang Y, Shen Z, Zhang Y, Zhang X, Nicosia SV, Zhang Y, Pledger JW, Chen J, Bai W.** MDM2 acts downstream of p53 as an E3 ligase to promote FOXO ubiquitination and degradation. *J Biol Chem* 2009; **284**: 13987-14000 [PMID: 19321440 DOI: 10.1074/jbc.M901758200]
  - 18 **Sunters A, Madureira PA, Pomeranz KM, Aubert M, Brokens JJ, Cook SJ, Burgering BM, Coombes RC, Lam EW.** Paclitaxel-induced nuclear translocation of FOXO3a in breast cancer cells is mediated by c-Jun NH2-terminal kinase and Akt. *Cancer Res* 2006; **66**: 212-220 [PMID: 16397234 DOI: 10.1158/0008-5472.CAN-05-1997]
  - 19 **Huang H, Tindall DJ.** Regulation of FOXO protein stability via ubiquitination and proteasome degradation. *Biochim Biophys Acta* 2011; **1813**: 1961-1964 [PMID: 21238503 DOI: 10.1016/j.bbamcr.2011.01.007]
  - 20 **Chatterjee A, Chatterjee U, Ghosh MK.** Activation of protein kinase CK2 attenuates FOXO3a functioning in a PML-dependent manner: implications in human prostate cancer. *Cell Death Dis* 2013; **4**: e543 [PMID: 23492774 DOI: 10.1038/cddis.2013.63]
  - 21 **Gao J, Yang X, Yin P, Hu W, Liao H, Miao Z, Pan C, Li N.** The involvement of FoxO in cell survival and chemosensitivity mediated by Mirk/Dyrk1B in ovarian cancer. *Int J Oncol* 2012; **40**: 1203-1209 [PMID: 22159921 DOI: 10.3892/ijo.2011.1293]
  - 22 **Wang X, Chen WR, Xing D.** A pathway from JNK through decreased ERK and Akt activities for FOXO3a nuclear translocation in response to UV irradiation. *J Cell Physiol* 2012; **227**: 1168-1178 [PMID: 21604264 DOI: 10.1002/jcp.22839]
  - 23 **Clavel S, Siffroi-Fernandez S, Coldefy AS, Boulukos K, Pisani DF, Dérjard B.** Regulation of the intracellular localization of Foxo3a by stress-activated protein kinase signaling pathways in skeletal muscle cells. *Mol Cell Biol* 2010; **30**: 470-480 [PMID: 19917721 DOI: 10.1128/MCB.00666-09]
  - 24 **Nho RS, Kahm J.** beta1-Integrin-collagen interaction suppresses FoxO3a by the coordination of Akt and PP2A. *J Biol Chem* 2010; **285**: 14195-14209 [PMID: 20223831 DOI: 10.1074/jcb.M109.052845]
  - 25 **Singh A, Ye M, Bucur O, Zhu S, Tanya Santos M, Rabinovitz I, Wei W, Gao D, Hahn WC, Khosravi-Far R.** Protein phosphatase 2A reactivates FOXO3a through a dynamic interplay with 14-3-3 and AKT. *Mol Biol Cell* 2010; **21**: 1140-1152 [PMID: 20110348 DOI: 10.1091/mbc.E09-09-0795]
  - 26 **Chiacchiera F, Simone C.** The AMPK-FoxO3A axis as a target for cancer treatment. *Cell Cycle* 2010; **9**: 1091-1096 [PMID: 20190568 DOI: 10.4161/cc.9.6.11035]
  - 27 **Milkiewicz M, Roudier E, Doyle JL, Trifonova A, Birot O, Haas TL.** Identification of a mechanism underlying regulation of the anti-angiogenic forkhead transcription factor FoxO1 in cultured endothelial cells and ischemic muscle. *Am J Pathol* 2011; **178**: 935-944 [PMID: 21281824 DOI: 10.1016/j.ajpath.2010.10.042]
  - 28 **Brenkman AB, de Keizer PL, van den Broek NJ, Jochemsen AG, Burgering BM.** Mdm2 induces mono-ubiquitination of FOXO4. *PLoS One* 2008; **3**: e2819 [PMID: 18665269 DOI: 10.1371/journal.pone.0002819]
  - 29 **Wang F, Marshall CB, Li GY, Yamamoto K, Mak TW, Ikura M.** Synergistic interplay between promoter recognition and CBP/p300 coactivator recruitment by FOXO3a. *ACS Chem Biol* 2009; **4**: 1017-1027 [PMID: 19821614 DOI: 10.1021/cb900190u]
  - 30 **Zeng Z, Samudio IJ, Zhang W, Estrov Z, Pelicano H, Harris D, Frolova O, Hail N, Chen W, Kornblau SM, Huang P, Lu Y, Mills GB, Andreeff M, Konopleva M.** Simultaneous inhibition of PDK1/AKT and Fms-like tyrosine kinase 3 signaling by a small-molecule KP372-1 induces mitochondrial dysfunction and apoptosis in acute myelogenous leukemia. *Cancer Res* 2006; **66**: 3737-3746 [PMID: 16585200 DOI: 10.1158/0008-5472.CAN-05-1278]
  - 31 **Wang F, Marshall CB, Yamamoto K, Li GY, Gasmi-Seabrook GM, Okada H, Mak TW, Ikura M.** Structures of KIX do-



- main of CBP in complex with two FOXO3a transactivation domains reveal promiscuity and plasticity in coactivator recruitment. *Proc Natl Acad Sci USA* 2012; **109**: 6078-6083 [PMID: 22474372 DOI: 10.1073/pnas.1119073109]
- 32 **Emerling BM**, Weinberg F, Liu JL, Mak TW, Chandel NS. PTEN regulates p300-dependent hypoxia-inducible factor 1 transcriptional activity through Forkhead transcription factor 3a (FOXO3a). *Proc Natl Acad Sci USA* 2008; **105**: 2622-2627 [PMID: 18268343 DOI: 10.1073/pnas.0706790105]
- 33 **Senf SM**, Sandesara PB, Reed SA, Judge AR. p300 Acetyltransferase activity differentially regulates the localization and activity of the FOXO homologues in skeletal muscle. *Am J Physiol Cell Physiol* 2011; **300**: C1490-C1501 [PMID: 21389279 DOI: 10.1152/ajpcell.00255.2010]
- 34 **Hori YS**, Kuno A, Hosoda R, Horio Y. Regulation of FOXOs and p53 by SIRT1 modulators under oxidative stress. *PLoS One* 2013; **8**: e73875 [PMID: 24040102 DOI: 10.1371/journal.pone.0073875]
- 35 **Kim MJ**, Ahn K, Park SH, Kang HJ, Jang BG, Oh SJ, Oh SM, Jeong YJ, Heo JI, Suh JG, Lim SS, Ko YJ, Huh SO, Kim SC, Park JB, Kim J, Kim JI, Jo SA, Lee JY. SIRT1 regulates tyrosine hydroxylase expression and differentiation of neuroblastoma cells via FOXO3a. *FEBS Lett* 2009; **583**: 1183-1188 [PMID: 19285077 DOI: 10.1016/j.febslet.2009.03.007]
- 36 **Ni HM**, Du K, You M, Ding WX. Critical role of FoxO3a in alcohol-induced autophagy and hepatotoxicity. *Am J Pathol* 2013; **183**: 1815-1825 [PMID: 24095927 DOI: 10.1016/j.ajpath.2013.08.011]
- 37 **Lai L**, Yan L, Gao S, Hu CL, Ge H, Davidow A, Park M, Bravo C, Iwatsubo K, Ishikawa Y, Auwerx J, Sinclair DA, Vatner SF, Vatner DE. Type 5 adenylyl cyclase increases oxidative stress by transcriptional regulation of manganese superoxide dismutase via the SIRT1/FoxO3a pathway. *Circulation* 2013; **127**: 1692-1701 [PMID: 23536361 DOI: 10.1161/CIRCULATIONAHA.112.001212]
- 38 **Brunet A**, Sweeney LB, Sturgill JF, Chua KF, Greer PL, Lin Y, Tran H, Ross SE, Mostoslavsky R, Cohen HY, Hu LS, Cheng HL, Jedrychowski MP, Gygi SP, Sinclair DA, Alt FW, Greenberg ME. Stress-dependent regulation of FOXO transcription factors by the SIRT1 deacetylase. *Science* 2004; **303**: 2011-2015 [PMID: 14976264 DOI: 10.1126/science.1094637]
- 39 **Babar IA**, Czocho J, Steinmetz A, Weidhaas JB, Glazer PM, Slack FJ. Inhibition of hypoxia-induced miR-155 radiosensitizes hypoxic lung cancer cells. *Cancer Biol Ther* 2011; **12**: 908-914 [PMID: 22027557 DOI: 10.4161/cbt.12.10.17681]
- 40 **Lin H**, Dai T, Xiong H, Zhao X, Chen X, Yu C, Li J, Wang X, Song L. Unregulated miR-96 induces cell proliferation in human breast cancer by downregulating transcriptional factor FOXO3a. *PLoS One* 2010; **5**: e15797 [PMID: 21203424 DOI: 10.1371/journal.pone.0015797]
- 41 **Ling N**, Gu J, Lei Z, Li M, Zhao J, Zhang HT, Li X. microRNA-155 regulates cell proliferation and invasion by targeting FOXO3a in glioma. *Oncol Rep* 2013; **30**: 2111-2118 [PMID: 23970205]
- 42 **Wang K**, Li PF. Foxo3a regulates apoptosis by negatively targeting miR-21. *J Biol Chem* 2010; **285**: 16958-16966 [PMID: 20371612 DOI: 10.1074/jbc.M109.093005]
- 43 **Cai J**, Fang L, Huang Y, Li R, Yuan J, Yang Y, Zhu X, Chen B, Wu J, Li M. miR-205 targets PTEN and PHLPP2 to augment AKT signaling and drive malignant phenotypes in non-small cell lung cancer. *Cancer Res* 2013; **73**: 5402-5415 [PMID: 23856247 DOI: 10.1158/0008-5472.CAN-13-0297]
- 44 **Nowak K**, Killmer K, Gessner C, Lutz W. E2F-1 regulates expression of FOXO1 and FOXO3a. *Biochim Biophys Acta* 2007; **1769**: 244-252 [PMID: 17482685 DOI: 10.1016/j.bbaexp.2007.04.001]
- 45 **Furuyama T**, Yamashita H, Kitayama K, Higami Y, Shimokawa I, Mori N. Effects of aging and caloric restriction on the gene expression of Foxo1, 3, and 4 (FKHR, FKHL1, and AFX) in the rat skeletal muscles. *Microsc Res Tech* 2002; **59**: 331-334 [PMID: 12424797 DOI: 10.1002/jemt.10213]
- 46 **Rathbone CR**, Booth FW, Lees SJ. FoxO3a preferentially induces p27Kip1 expression while impairing muscle precursor cell-cycle progression. *Muscle Nerve* 2008; **37**: 84-89 [PMID: 17894357 DOI: 10.1002/mus.20897]
- 47 **Roy SK**, Chen Q, Fu J, Shankar S, Srivastava RK. Resveratrol inhibits growth of orthotopic pancreatic tumors through activation of FOXO transcription factors. *PLoS One* 2011; **6**: e25166 [PMID: 21980390 DOI: 10.1371/journal.pone.0025166]
- 48 **Gilley J**, Coffey PJ, Ham J. FOXO transcription factors directly activate bim gene expression and promote apoptosis in sympathetic neurons. *J Cell Biol* 2003; **162**: 613-622 [PMID: 12913110 DOI: 10.1083/jcb.200303026]
- 49 **Wilson MK**, McWhirter SM, Amin RH, Huang D, Schlissel MS. Abelson virus transformation prevents TRAIL expression by inhibiting FoxO3a and NF-kappaB. *Mol Cells* 2010; **29**: 333-341 [PMID: 20213318 DOI: 10.1007/s10059-010-0029-8]
- 50 **Ghaffari S**, Jagani Z, Kitidis C, Lodish HF, Khosravi-Far R. Cytokines and BCR-ABL mediate suppression of TRAIL-induced apoptosis through inhibition of forkhead FOXO3a transcription factor. *Proc Natl Acad Sci USA* 2003; **100**: 6523-6528 [PMID: 12750477 DOI: 10.1073/pnas.0731871100]
- 51 **van den Heuvel AP**, Schulze A, Burgering BM. Direct control of caveolin-1 expression by FOXO transcription factors. *Biochem J* 2005; **385**: 795-802 [PMID: 15458387 DOI: 10.1042/BJ20041449]
- 52 **Kops GJ**, Medema RH, Glassford J, Essers MA, Dijkers PF, Coffey PJ, Lam EW, Burgering BM. Control of cell cycle exit and entry by protein kinase B-regulated forkhead transcription factors. *Mol Cell Biol* 2002; **22**: 2025-2036 [PMID: 11884591 DOI: 10.1128/MCB.22.7.2025-2036.2002]
- 53 **Hauck L**, Harms C, Grothe D, An J, Gertz K, Kronenberg G, Dietz R, Endres M, von Harsdorf R. Critical role for FoxO3a-dependent regulation of p21CIP1/WAF1 in response to statin signaling in cardiac myocytes. *Circ Res* 2007; **100**: 50-60 [PMID: 17158337 DOI: 10.1161/01.RES.0000254704.92532.b9]
- 54 **Miyachi H**, Minamino T, Tateno K, Kunieda T, Toko H, Komuro I. Akt negatively regulates the in vitro lifespan of human endothelial cells via a p53/p21-dependent pathway. *EMBO J* 2004; **23**: 212-220 [PMID: 14713953 DOI: 10.1038/sj.emboj.7600045]
- 55 **Chandramohan V**, Jeay S, Pianetti S, Sonenshein GE. Reciprocal control of Forkhead box O 3a and c-Myc via the phosphatidylinositol 3-kinase pathway coordinately regulates p27Kip1 levels. *J Immunol* 2004; **172**: 5522-5527 [PMID: 15100294]
- 56 **Lin L**, Hron JD, Peng SL. Regulation of NF-kappaB, Th activation, and autoinflammation by the forkhead transcription factor Foxo3a. *Immunity* 2004; **21**: 203-213 [PMID: 15308101 DOI: 10.1016/j.immuni.2004.06.016]
- 57 **Hwang JW**, Rajendrasozhan S, Yao H, Chung S, Sundar IK, Huyck HL, Pryhuber GS, Kinnula VL, Rahman I. FOXO3 deficiency leads to increased susceptibility to cigarette smoke-induced inflammation, airspace enlargement, and chronic obstructive pulmonary disease. *J Immunol* 2011; **187**: 987-998 [PMID: 21690325 DOI: 10.4049/jimmunol.1001861]
- 58 **You H**, Pellegrini M, Tsuchihara K, Yamamoto K, Hacker G, Erlacher M, Villunger A, Mak TW. FOXO3a-dependent regulation of Puma in response to cytokine/growth factor withdrawal. *J Exp Med* 2006; **203**: 1657-1663 [PMID: 16801400 DOI: 10.1084/jem.20060353]
- 59 **Obexer P**, Geiger K, Ambros PF, Meister B, Ausserlechner MJ. FKHL1-mediated expression of Noxa and Bim induces apoptosis via the mitochondria in neuroblastoma cells. *Cell Death Differ* 2007; **14**: 534-547 [PMID: 16888645 DOI: 10.1038/sj.cdd.4402017]
- 60 **Sengupta A**, Molkenin JD, Paik JH, DePinho RA, Yutzey KE. FoxO transcription factors promote cardiomyocyte

- survival upon induction of oxidative stress. *J Biol Chem* 2011; **286**: 7468-7478 [PMID: 21159781 DOI: 10.1074/jbc.M110.179242]
- 61 **Liu JW**, Chandra D, Rudd MD, Butler AP, Pallotta V, Brown D, Coffey PJ, Tang DG. Induction of prosurvival molecules by apoptotic stimuli: involvement of FOXO3a and ROS. *Oncogene* 2005; **24**: 2020-2031 [PMID: 15674333 DOI: 10.1038/sj.onc.1208385]
- 62 **Murphy CT**, McCarroll SA, Bargmann CI, Fraser A, Kamath RS, Ahringer J, Li H, Kenyon C. Genes that act downstream of DAF-16 to influence the lifespan of *Caenorhabditis elegans*. *Nature* 2003; **424**: 277-283 [PMID: 12845331 DOI: 10.1038/nature01789]
- 63 **McCormick M**, Chen K, Ramaswamy P, Kenyon C. New genes that extend *Caenorhabditis elegans*' lifespan in response to reproductive signals. *Aging Cell* 2012; **11**: 192-202 [PMID: 22081913 DOI: 10.1111/j.1474-9726.2011.00768.x]
- 64 **Sengupta A**, Molkentin JD, Yutzey KE. FoxO transcription factors promote autophagy in cardiomyocytes. *J Biol Chem* 2009; **284**: 28319-28331 [PMID: 19696026 DOI: 10.1074/jbc.M109.024406]
- 65 **Sundaresan NR**, Gupta M, Kim G, Rajamohan SB, Isbatan A, Gupta MP. Sirt3 blocks the cardiac hypertrophic response by augmenting Foxo3a-dependent antioxidant defense mechanisms in mice. *J Clin Invest* 2009; **119**: 2758-2771 [PMID: 19652361]
- 66 **Huang W**, Li G, Qiu J, Gonzalez P, Challa P. Protective effects of resveratrol in experimental retinal detachment. *PLoS One* 2013; **8**: e75735 [PMID: 24040416 DOI: 10.1371/journal.pone.0075735]
- 67 **Li M**, Chiu JF, Gagne J, Fukagawa NK. Age-related differences in insulin-like growth factor-1 receptor signaling regulates Akt/FOXO3a and ERK/Fos pathways in vascular smooth muscle cells. *J Cell Physiol* 2008; **217**: 377-387 [PMID: 18615585 DOI: 10.1002/jcp.21507]
- 68 **Willcox BJ**, Donlon TA, He Q, Chen R, Grove JS, Yano K, Masaki KH, Willcox DC, Rodriguez B, Curb JD. FOXO3A genotype is strongly associated with human longevity. *Proc Natl Acad Sci USA* 2008; **105**: 13987-13992 [PMID: 18765803 DOI: 10.1073/pnas.0801030105]
- 69 **Flachsbar F**, Caliebe A, Kleindorfer R, Blanché H, von Eller-Eberstein H, Nikolaus S, Schreiber S, Nebel A. Association of FOXO3A variation with human longevity confirmed in German centenarians. *Proc Natl Acad Sci USA* 2009; **106**: 2700-2705 [PMID: 19196970 DOI: 10.1073/pnas.0809594106]
- 70 **Morris BJ**. A forkhead in the road to longevity: the molecular basis of lifespan becomes clearer. *J Hypertens* 2005; **23**: 1285-1309 [PMID: 15942449 DOI: 10.1097/01.hjh.0000173509.45363.dd]
- 71 **Eelen G**, Verlinden L, Meyer MB, Gijssbers R, Pike JW, Bouillon R, Verstuyf A. 1,25-Dihydroxyvitamin D3 and the aging-related forkhead box O and sestrin proteins in osteoblasts. *J Steroid Biochem Mol Biol* 2013; **136**: 112-119 [PMID: 22989484 DOI: 10.1016/j.jsbmb.2012.09.011]
- 72 **Kops GJ**, Dansen TB, Polderman PE, Saarloos I, Wirtz KW, Coffey PJ, Huang TT, Bos JL, Medema RH, Burgering BM. Forkhead transcription factor FOXO3a protects quiescent cells from oxidative stress. *Nature* 2002; **419**: 316-321 [PMID: 12239572 DOI: 10.1038/nature01036]
- 73 **Giordano S**, Darley-Usmar V, Zhang J. Autophagy as an essential cellular antioxidant pathway in neurodegenerative disease. *Redox Biol* 2014; **2**: 82-90 [PMID: 24494187 DOI: 10.1016/j.redox.2013.12.013]
- 74 **El Assar M**, Angulo J, Rodríguez-Mañás L. Oxidative stress and vascular inflammation in aging. *Free Radic Biol Med* 2013; **65**: 380-401 [PMID: 23851032 DOI: 10.1016/j.freeradbiomed.2013.07.003]
- 75 **Liochev SI**. Reactive oxygen species and the free radical theory of aging. *Free Radic Biol Med* 2013; **60**: 1-4 [PMID: 23434764 DOI: 10.1016/j.freeradbiomed.2013.02.011]
- 76 **van der Horst A**, Burgering BM. Stressing the role of FoxO proteins in lifespan and disease. *Nat Rev Mol Cell Biol* 2007; **8**: 440-450 [PMID: 17522590 DOI: 10.1038/nrm2190]
- 77 **Chen IC**, Chiang WF, Chen PF, Chiang HC. STRESS-responsive deacetylase SIRT3 is up-regulated by areca nut extract-induced oxidative stress in human oral keratinocytes. *J Cell Biochem* 2014; **115**: 328-339 [PMID: 24339251 DOI: 10.1002/jcb.24667]
- 78 **Zhang S**, Zhao Y, Xu M, Yu L, Zhao Y, Chen J, Yuan Y, Zheng Q, Niu X. FoxO3a modulates hypoxia stress induced oxidative stress and apoptosis in cardiac microvascular endothelial cells. *PLoS One* 2013; **8**: e80342 [PMID: 24278276 DOI: 10.1371/journal.pone.0080342]
- 79 **Raju I**, Kannan K, Abraham EC. FoxO3a Serves as a Biomarker of Oxidative Stress in Human Lens Epithelial Cells under Conditions of Hyperglycemia. *PLoS One* 2013; **8**: e67126 [PMID: 23805295 DOI: 10.1371/journal.pone.0067126]
- 80 **Chen SJ**, Zhang W, Tong Q, Conrad K, Hirschler-Laszkiewicz I, Bayerl M, Kim JK, Cheung JY, Miller BA. Role of TRPM2 in cell proliferation and susceptibility to oxidative stress. *Am J Physiol Cell Physiol* 2013; **304**: C548-C560 [PMID: 23302782 DOI: 10.1152/ajpcell.00069.2012]
- 81 **Li Z**, Zhang H, Chen Y, Fan L, Fang J. Forkhead transcription factor FOXO3a protein activates nuclear factor  $\kappa$ B through B-cell lymphoma/leukemia 10 (BCL10) protein and promotes tumor cell survival in serum deprivation. *J Biol Chem* 2012; **287**: 17737-17745 [PMID: 22474286 DOI: 10.1074/jbc.M111.291708]
- 82 **Tseng AH**, Shieh SS, Wang DL. SIRT3 deacetylates FOXO3 to protect mitochondria against oxidative damage. *Free Radic Biol Med* 2013; **63**: 222-234 [PMID: 23665396 DOI: 10.1016/j.freeradbiomed.2013.05.002]
- 83 **Yang JY**, Chang CJ, Xia W, Wang Y, Wong KK, Engelman JA, Du Y, Andreeff M, Hortobagyi GN, Hung MC. Activation of FOXO3a is sufficient to reverse mitogen-activated protein/extracellular signal-regulated kinase inhibitor chemoresistance in human cancer. *Cancer Res* 2010; **70**: 4709-4718 [PMID: 20484037 DOI: 10.1158/0008-5472.CAN-09-4524]
- 84 **Chen J**, Gomes AR, Monteiro LJ, Wong SY, Wu LH, Ng TT, Karadedou CT, Millour J, Ip YC, Cheung YN, Sunter A, Chan KY, Lam EW, Khoo US. Constitutively nuclear FOXO3a localization predicts poor survival and promotes Akt phosphorylation in breast cancer. *PLoS One* 2010; **5**: e12293 [PMID: 20808831 DOI: 10.1371/journal.pone.0012293]
- 85 **Lam M**, Carmichael AR, Griffiths HR. An aqueous extract of *Fagonia cretica* induces DNA damage, cell cycle arrest and apoptosis in breast cancer cells via FOXO3a and p53 expression. *PLoS One* 2012; **7**: e40152 [PMID: 22761954 DOI: 10.1371/journal.pone.0040152]
- 86 **Shukla S**, Bhaskaran N, MacLennan GT, Gupta S. Deregulation of FoxO3a accelerates prostate cancer progression in TRAMP mice. *Prostate* 2013; **73**: 1507-1517 [PMID: 23765843 DOI: 10.1002/pros.22698]
- 87 **Shukla S**, Bhaskaran N, Babcook MA, Fu P, MacLennan GT, Gupta S. Apigenin inhibits prostate cancer progression in TRAMP mice via targeting PI3K/Akt/FoxO pathway. *Carcinogenesis* 2014; **35**: 452-460 [PMID: 24067903 DOI: 10.1093/carcin/bgt316]
- 88 **Chen Q**, Ganapathy S, Singh KP, Shankar S, Srivastava RK. Resveratrol induces growth arrest and apoptosis through activation of FOXO transcription factors in prostate cancer cells. *PLoS One* 2010; **5**: e15288 [PMID: 21179458 DOI: 10.1371/journal.pone.0015288]
- 89 **Sunayama J**, Sato A, Matsuda K, Tachibana K, Watanabe E, Seino S, Suzuki K, Narita Y, Shibui S, Sakurada K, Kayama T, Tomiyama A, Kitanaka C. FoxO3a functions as a key integrator of cellular signals that control glioblastoma stem-like cell differentiation and tumorigenicity. *Stem Cells* 2011; **29**:



- 1327-1337 [PMID: 21793107]
- 90 **Ruvolo PP.** The Herculean task of killing cancer cells: suppression of FOXO3A in acute leukemia involves a hydra of multiple survival kinases. *Cell Cycle* 2012; **11**: 2589 [PMID: 22751430 DOI: 10.4161/cc.21233]
- 91 **Wang W,** Li NN, Du Y, Lv FF, Lin GQ. FoxO3a and nilotinib-induced erythroid differentiation of CML-BC cells. *Leuk Res* 2013; **37**: 1309-1314 [PMID: 23915976 DOI: 10.1016/j.leukres.2013.07.001]
- 92 **Yang W,** Dolloff NG, El-Deiry WS. ERK and MDM2 prey on FOXO3a. *Nat Cell Biol* 2008; **10**: 125-126 [PMID: 18246039 DOI: 10.1038/ncb0208-125]
- 93 **Sunters A,** Fernández de Mattos S, Stahl M, Brosens JJ, Zoumpoulidou G, Saunders CA, Coffey PJ, Medema RH, Coombes RC, Lam EW. FoxO3a transcriptional regulation of Bim controls apoptosis in paclitaxel-treated breast cancer cell lines. *J Biol Chem* 2003; **278**: 49795-49805 [PMID: 14527951 DOI: 10.1074/jbc.M309523200]
- 94 **Hui RC,** Francis RE, Guest SK, Costa JR, Gomes AR, Myatt SS, Brosens JJ, Lam EW. Doxorubicin activates FOXO3a to induce the expression of multidrug resistance gene ABCB1 (MDR1) in K562 leukemic cells. *Mol Cancer Ther* 2008; **7**: 670-678 [PMID: 18347152 DOI: 10.1158/1535-7163.MCT-07-0397]
- 95 **Naka K,** Hoshii T, Muraguchi T, Tadokoro Y, Ooshio T, Kondo Y, Nakao S, Motoyama N, Hirao A. TGF-beta-FOXO signalling maintains leukaemia-initiating cells in chronic myeloid leukaemia. *Nature* 2010; **463**: 676-680 [PMID: 20130650 DOI: 10.1038/nature08734]
- 96 **Birkenkamp KU,** Essafi A, van der Vos KE, da Costa M, Hui RC, Holstege F, Koenderman L, Lam EW, Coffey PJ. FOXO3a induces differentiation of Bcr-Abl-transformed cells through transcriptional down-regulation of Id1. *J Biol Chem* 2007; **282**: 2211-2220 [PMID: 17132628 DOI: 10.1074/jbc.M606669200]
- 97 **Buontempo F,** Chiarini F, Bressanin D, Tabellini G, Melchionda F, Pession A, Fini M, Neri LM, McCubrey JA, Martelli AM. Activity of the selective IκB kinase inhibitor BMS-345541 against T-cell acute lymphoblastic leukemia: involvement of FOXO3a. *Cell Cycle* 2012; **11**: 2467-2475 [PMID: 22713244 DOI: 10.4161/cc.20859]
- 98 **Dey P,** Ström A, Gustafsson JÅ. Estrogen receptor β upregulates FOXO3a and causes induction of apoptosis through PUMA in prostate cancer. *Oncogene* 2014; **33**: 4213-4225 [PMID: 24077289 DOI: 10.1038/onc.2013.384]
- 99 **Jia Y,** Mo SJ, Feng QQ, Zhan ML, OuYang LS, Chen JC, Ma YX, Wu JJ, Lei WL. EPO-dependent activation of PI3K/Akt/FoxO3a signalling mediates neuroprotection in in vitro and in vivo models of Parkinson's disease. *J Mol Neurosci* 2014; **53**: 117-124 [PMID: 24390959 DOI: 10.1007/s12031-013-0208-0]
- 100 **Carbajo-Pescador S,** Mauriz JL, García-Palomo A, González-Gallego J. FoxO proteins: regulation and molecular targets in liver cancer. *Curr Med Chem* 2014; **21**: 1231-1246 [PMID: 24372208]
- 101 **Liang C,** Chen W, Zhi X, Ma T, Xia X, Liu H, Zhang Q, Hu Q, Zhang Y, Bai X, Liang T. Serotonin promotes the proliferation of serum-deprived hepatocellular carcinoma cells via upregulation of FOXO3a. *Mol Cancer* 2013; **12**: 14 [PMID: 23418729 DOI: 10.1186/1476-4598-12-14]
- 102 **Ghosh Choudhury G,** Lenin M, Calhaun C, Zhang JH, Aboud HE. PDGF inactivates forkhead family transcription factor by activation of Akt in glomerular mesangial cells. *Cell Signal* 2003; **15**: 161-170 [PMID: 12464387 DOI: 10.1016/S0898-6568(02)00057-8]

**P- Reviewer:** Moreno JJ, Pospelov VA, Yew PR

**S- Editor:** Wen LL **L- Editor:** Roemmele A **E- Editor:** Lu YJ





Published by **Baishideng Publishing Group Inc**  
8226 Regency Drive, Pleasanton, CA 94588, USA  
Telephone: +1-925-223-8242  
Fax: +1-925-223-8243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>  
<http://www.wjgnet.com>

