

# World Journal of *Orthopedics*

*World J Orthop* 2019 March 18; 10(3): 123-175



**MINIREVIEWS**

- 123 Linkage of microbiota and osteoporosis: A mini literature review  
*Yatonsky II D, Pan K, Shendge VB, Liu J, Ebraheim NA*

**ORIGINAL ARTICLE****Retrospective Cohort Study**

- 128 Short-term differences in anterior knee pain and clinical outcomes between rotating and fixed platform posterior stabilized total knee arthroplasty with a new femoral component design  
*Bigoni M, Zanchi N, Turati M, Pirovano G, Zatti G, Munegato D*

**Retrospective Study**

- 137 Comparison of implant related complications amongst patients with opioid use disorder and non-users following total knee arthroplasty  
*Vakharia RM, Sabeh KG, Vakharia AM, Damodar DM, Law TY, Roche MW*

**SYSTEMATIC REVIEW**

- 145 Aetiology of Legg-Calvé-Perthes disease: A systematic review  
*Pavone V, Chisari E, Vescio A, Lizzio C, Sessa G, Testa G*
- 166 Changing trends in the mortality rate at 1-year post hip fracture - a systematic review  
*Downey C, Kelly M, Quinlan JF*

**ABOUT COVER**

Editorial Board of *World Journal of Orthopedics*, Sarah Cartmell, PhD, Professor, School of Materials, The University of Manchester, Manchester M1 7HS, United Kingdom

**AIMS AND SCOPE**

*World Journal of Orthopedics (World J Orthop, WJO*, online ISSN 2218-5836, DOI: 10.5312 ) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

*WJO* covers topics concerning arthroscopy, evidence-based medicine, epidemiology, nursing, sports medicine, therapy of bone and spinal diseases, bone trauma, osteoarthropathy, bone tumors and osteoporosis, etc. Priority publication will be given to articles concerning diagnosis and treatment of orthopedic diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, etc.

We encourage authors to submit their manuscripts to *WJO*.

**INDEXING/ABSTRACTING**

The *WJO* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (CSTJ), and Superstar Journals Database.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Responsible Electronic Editor: *Yun-Xiaojuan Wu* Proofing Editorial Office Director: *Jin-Lei Wang*

**NAME OF JOURNAL**

*World Journal of Orthopedics*

**ISSN**

ISSN 2218-5836 (online)

**LAUNCH DATE**

November 18, 2010

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng

**EDITORIAL BOARD MEMBERS**

<http://www.wjgnet.com/2218-5836/editorialboard.htm>

**EDITORIAL OFFICE**

Jin-Lei Wang, Director

**PUBLICATION DATE**

March 18, 2019

**COPYRIGHT**

© 2019 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Changing trends in the mortality rate at 1-year post hip fracture - a systematic review

Colum Downey, Martin Kelly, John F Quinlan

**ORCID number:** Colum Downey (0000-0001-6594-5732); Martin Kelly (<https://orcid.org/0000-0002-9798-7969>); John Quinlan (0000-0002-9561-7665).

**Author contributions:** Downey C, Kelly M and Quinlan JF contributed equally to this work. Downey C and Quinlan JF designed research. Downey C, Kelly M performed research. Downey C wrote the paper.

**Conflict-of-interest statement:** The authors declare that they have no conflict of interest.

**PRISMA 2009 Checklist statement:** The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Unsolicited manuscript

**Received:** October 10, 2018

**Peer-review started:** October 10,

**Colum Downey, Martin Kelly, John F Quinlan,** Department of Trauma and Orthopaedics, Tallaght University Hospital, Dublin 01, Ireland

**Corresponding author:** Colum Downey, BSc, MBBS, MSc, Doctor, Department of Trauma and Orthopaedics, Tallaght University Hospital, Dublin 01, Ireland. [columdowney@rcsi.ie](mailto:columdowney@rcsi.ie)

**Telephone:** +353-1-4144090

**Fax:** +353-1-4144779

### Abstract

#### BACKGROUND

Traditionally, the mortality rate at 1-year post hip fracture was quoted as approximately 30% of all hip fractures. There have been recent improvements in hip fracture care in the main driven by national hip fracture registries with reductions in 30-d mortality rates reported.

#### AIM

To address recent 1-year post hip fracture mortality rates in the literature.

#### METHODS

Systematic literature review, national hip fracture registries/databases, local studies on hip fracture mortality, 5 years limitation (2013-2017), cohorts > 100, studies in English. Outcome measure: Mortality rate at 1-year post hip fracture.

#### RESULTS

Recent 1-year mortality rates were reviewed using the literature from 8 National Registries and 36 different countries. Recently published 1-year mortality rates appear lower than traditional figures and may represent a downward trend.

#### CONCLUSION

There appears to be a consistent worldwide reduction in mortality at 1-year post hip fracture compared to previously published research. Globally, those which suffer hip fractures may currently be benefiting from the results of approximately 30 years of national registries, rigorous audit processes and international collaboration. The previously quoted mortality rates of 10% at 1-mo and 30% at 1-year may be outdated.

**Key words:** Hip fracture; Mortality; Registries; Databases

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

2018

**First decision:** October 25, 2018**Revised:** January 1, 2019**Accepted:** February 27, 2019**Article in press:** February 27, 2019**Published online:** March 18, 2019

**Core tip:** This review reports a global downward trend in 1-year mortality rate post hip fracture. The results of this study suggest that the previously reported 1-year mortality rate of 30% is out-dated and a figure of 22% is a more accurate, up-to-date estimation of 1-year mortality rate post hip fracture.

**Citation:** Downey C, Kelly M, Quinlan JF. Changing trends in the mortality rate at 1-year post hip fracture - a systematic review. *World J Orthop* 2019; 10(3): 166-175

**URL:** <https://www.wjgnet.com/2218-5836/full/v10/i3/166.htm>

**DOI:** <https://dx.doi.org/10.5312/wjo.v10.i3.166>

## INTRODUCTION

Hip fracture care continues to improve secondary to international audit and optimization of hip fracture management. Hip fractures are exceedingly common presentations to global emergency departments with approximately 3600 in Ireland per year<sup>[1]</sup>, approximately 65000 in the United Kingdom<sup>[2]</sup>, approximately 620000 in the European Union<sup>[3]</sup>, approximately 300000 in the United States<sup>[4]</sup> and roughly 2 million worldwide<sup>[5]</sup>. This global figure is modest projected to increase threefold by 2050<sup>[6]</sup>.

There is significant cost associated with hip fractures regarding: (A) financial cost to the health service providing care; and (B) morbidity and mortality cost to the patient. Recent research estimates the year 1 hospital cost secondary to an index hip fracture to be £14000 (€15900/\$18750) per patient<sup>[7]</sup>. The annual cost of treatment in the European Union and the United States has been recently reported as €32 billion and \$20 billion respectively<sup>[8]</sup>.

The literature describes that after suffering a hip fracture, the patient will experience a reduction in overall functional status, social independence, mobility and increasing scores in depressive symptoms<sup>[9-10]</sup>. Due to the high prevalence of hip fractures and the disadvantage to both the patient and the local health service responsible for care, optimisation and the standardisation of hip fracture management has been targeted to reduce morbidity, mortality and financial expense. This optimisation and standardization, in the form of a national registry, has been shown previously to improve the quality of care and outcomes of hip fracture patients<sup>[11]</sup>.

The earliest hip fracture registry established is "RIKSHÖFT", the Swedish national hip fracture registry, set up in 1988, following on from the successful local implementation of national knee and hip arthroplasty registers in the 1970s<sup>[12]</sup>. In 1993, the Scottish Hip Fracture Audit (SHFA) implemented a registry that reported annually until 2008, after which, the audit was stepped down to capture "snapshot" data only. Following audit and the recognition of deterioration in outcomes post step down<sup>[13]</sup>, the SHFA was re-implemented in May 2016 and year round data collection was re-instituted<sup>[14]</sup>. In 1995, the European Commission funded the initiation of the Standardised Audit of Hip Fracture in Europe due to the triumphs in outcomes secondary to the "RIKSHÖFT" audit in Sweden<sup>[15]</sup>. The Danish orthopaedic community established the Danish Hip Fracture Database in 2003<sup>[5]</sup> whilst the Norwegian community set up their database, the "NOREPOS", in 2005<sup>[16]</sup>.

In 2007, an important milestone was reached regarding the initiation of the largest and fastest growing national hip fracture registry by the United Kingdom. After publishing the "Blue Book", which set quality standards established by the British Orthopaedic Association and the British Geriatrics Society, the National Hip Fracture Database (NFHD) began to collect and publish data annually regarding hip fractures<sup>[3]</sup>. In 2010, a European Union program, named euroHOPE (Healthcare Outcomes, Performance, Efficiency), was created to compare healthcare performance regarding hip fracture treatment across 7 European countries<sup>[17]</sup>. The fragility fracture network was then established in 2011, a global network, with a mission to optimize the multidisciplinary management of patients with fragility fracture globally and prevent secondary fracture<sup>[18]</sup>. This was followed by the foundation of the Irish Hip Fracture Database in 2012<sup>[1]</sup> and more recently the Australian and New Zealand Hip Fracture Registries in 2016<sup>[19]</sup>. Several countries are currently piloting their own national hip registries following the achievements published by the registries described above including Greece, Malta, Netherlands, Slovenia, Spain, Lebanon and Argentina.

Traditionally, orthopaedic trainees are taught that the mortality rates in this fragile

population were approximately 10% at 30 d and approximately 30% at 1 year<sup>[20]</sup>. These figures are important at: (A) a patient level with regards to family discussions surrounding realistic expectations and prognosis following hip fracture; and (B) at a hospital and national level with regards to the optimisation of care and distribution of finite resources.

In 2015, following a review of 471590 patients which suffered a hip fracture in the United Kingdom between 2003-2011, it was concluded that “the launch of a national clinician-led audit initiative was associated with substantial improvements in care and survival of older people”<sup>[21]</sup>. This group reported significant reductions in 30-d mortality.

Following review of the recent literature emanating from the national registries, it was noted that whilst 87.5% of the national registries publish annual data on in-patient mortality and 62.5% publish annually on 30-d mortality, only 12.5% (1 registry) published mortality rate at 1-year post hip fracture. In order to explore whether there have been any recent global reductions in 1-year mortality post fracture, a systematic literature review was undertaken.

---

## MATERIALS AND METHODS

---

A review of the available literature on (A) national hip fracture registries/databases and (B) local hip fracture studies which included 1-year mortality data was conducted using Pubmed and Google Scholar. The following keywords were used: “hip fracture”, “mortality”, “year”, “death”, “outcomes”, “registry”, “registries”, “database” and “databases”. The inclusion criteria were those studies published within the last 5 years (2013-2017), studies including a cohort > 100 and those in the English language. The most recent national registry reports were obtained and personal contact to authors of the Swedish and Danish reports *via* e-mail was required as these are reported in local language only.

---

## RESULTS

---

### **National Registries**

**Table 1** lists the countries, registry names, foundation year, age inclusion criteria, recent annual reports with reference and the mortality data published by each of the 8 national databases. 87.5% of the registries have an age inclusion criteria of 50 years or more. The table shows a dearth in published data for 1-year mortality rates post hip fracture.

### **Europe**

Recent (previous 5 years) 1-year mortality research data from the European continent was extracted from the literature and is summarised in **Table 2**. There were 22 articles reviewed with a mean 1-year mortality of 23.3% [standard deviation (SD) 6.3%, median 23.4%] reported. The lowest figure was 9.7% with the highest 34.8%. 19 of the 22 articles (86%) reported data regarding cohort numbers ranging from 124 to 31668 patients included (median 444, mean 6902, SD 11804). 91% reported age related inclusion criteria, with 55% including > 60 years only and the remaining 45% including patients < 60 years old. 82% of the articles were published within the period 2015-2017.

### **Asia**

One year mortality data (2013-2017) from the Asian continent was extracted from the literature and is summarised in **Table 3**. There were 10 articles reviewed with a mean 1-year mortality of 17.89% (SD 8.7%, median 18.9%) reported in studies selected. The lowest figure was 2.4% with the highest 30.4%. All of the articles reported data regarding cohort numbers, which ranged from 110 to 43830 patients included (median 400, mean 5621, SD 13566). 90% reported age related inclusion criteria, with 67% including > 60 years only and the remaining 33% including patients < 60 years old. 70% of the articles were published in the year 2017 with 90% published within 2015-2017.

### **Oceania**

**Table 4** lists the results from a large study undertaken in New South Wales, Australia which published a 1-year mortality rate post hip fracture of 24.9%. This study included 27888 patients which were aged > 60 years. To the best of the author's knowledge, there were no recent local studies emanating from New Zealand

**Table 1** Data from the National Registries

Country	Registry	Setup	Age	Report	Inpatient (%)	30-d (%)	1-Yr (%)
Sweden	RIKSHÖFT	1988	> 50	2016 <sup>[22]</sup>	3.39	-	-
Scotland	SHFA	1993	> 50	201 <sup>[23]</sup>	5	-	-
Denmark	DHFD	2003	> 65	2016 <sup>[24]</sup>	3	10	-
Norway	NHFR	2005	> 18	2016 <sup>[25]</sup>	-	7.7	24
UK*	NHFD	2007	> 60	2016 <sup>[2]</sup>	7	6.7	-
Ireland	IHFD	2012	> 60	2016 <sup>[1]</sup>	5	-	-
Australia	ANZHFR	2016	> 50	2016 <sup>[19]</sup>	5	5	-
New Zealand	ANZHFR	2016	> 50	2016 <sup>[19]</sup>	4	3	-

\*England, Wales & Northern Ireland. SHFA: Scottish Hip Fracture Audit; DHFD: Danish Hip Fracture Database; NHFD: National Hip Fracture Database; ANZHFR: Australian and New Zealand Guideline for Hip Fracture Care.

regarding mortality post hip fracture.

### North America

Table 5 lists the results from a large study undertaken in the United States using the Kaiser Permanente database. This recent publication describes a 1-year mortality rate post hip fracture of 21%. This study included 14294 patients which were aged > 60 years. To the best of the author's knowledge, there were no recent local studies published from Canada regarding 1-year mortality post hip fracture.

### South America

One year mortality data from the South American continent was extracted from the literature and is summarised in Table 6. There were 2 articles reviewed with a mean 1-year mortality of 26.8% (SD 4.5%). The articles included small cohorts with 100 and 213 patients respectively. Both of the studies included patients > 65 only.

### Africa

There has been no research regarding hip fracture mortality published throughout the African continent during the past 5 years to the best of the authors' knowledge.

### Global overview

Overall, local hip fracture related studies from 36 different countries were reviewed with regards to 1-year mortality rates post hip fracture. The mean overall 1-year mortality rate was 22.0% (SD 7.2%, median 22.8%) with a range from 2.4%-34.8%. A combined total of 229851 patients were included, with a range of 100-43830 patients from the smallest and largest cohort respectively. The median cohort was 402 patients. 61% of the studies involved a cohort between 100-1000 patients, with 18% between 1001-10000 and the remaining 21% including more than 10001 patients. 58.3% of the studies included patients > 60 years only, with 33.3% including patients < 60 also with a small proportion not declaring age inclusion/exclusion criteria. Using studies which included patients > 60 years only ( $n = 21/36$ ) may result in a more accurate representation of fragility hip fracture cohorts, however the results were quite similar with a mean rate of 21.8% (SD 7.6%, median 23.6%) mortality 1-year post hip fracture. 83.3% of the local studies research included in this study was published in the years 2015-2017.

## DISCUSSION

The mortality at 1 year post hip fracture has traditionally been reported as 30%<sup>[20]</sup> and more recently described as "around a third within a year"<sup>[60]</sup>, whilst the 2017 NFHD annual report suggests "up to a third die within a year following a hip fracture"<sup>[2]</sup>. There have been significant developments with regards to data collection in the areas of casemix, surgical/anaesthetic practice, process and outcomes of hip fracture care by the national registries. However, only 1 of the 8 national registries is publishing data regarding mortality at 1-year post hip fracture. There is therefore a void in the published literature on this topic.

After reviewing the recent hip fracture related mortality literature, it was found that the mean 1-year mortality rate in Europe was 23.3% (22 studies), whilst in Asia

**Table 2** Data from local hip fracture studies in Europe

Country	Origin	Published	n	Age	1-Yr (%)
Ireland	Limerick <sup>[26]</sup>	2017	206	> 60	9.7
Sweden	Lund <sup>[27]</sup>	2016	373	> 50	9.91
Denmark	Copenhagen <sup>[28]</sup>	2017	444	> 60	16
Italy	euroHOPE Registry <sup>[17]</sup>	2015	-	-	19.1
Switzerland	National Medical Statistic <sup>[29]</sup>	2014	24,678	> 18	20
Austria	Austrian Social Insur. Data <sup>[30]</sup>	2014	31,668	> 50	20.2
Portugal	Nat. Hosp. Discharge Reg <sup>[31]</sup>	2015	186	> 60	20.4
UK	Clin. Practice Res. Datalink <sup>[32]</sup>	2014	31,495	> 18	20.5
Romania	Timisoara <sup>[33]</sup>	2014	1,866	> 55	21.1
Nor. Ireland	FORD Database <sup>[34]</sup>	2017	27,055	> 18	21.7
Netherlands	National Discharge Register <sup>[35]</sup>	2017	850	> 70	23.2
Finland	National Discharge Register <sup>[17]</sup>	2015	-	> 60	23.6
Malta	National Mortality Register <sup>[36]</sup>	2016	281	> 60	25.6
Norway	Trondheim <sup>[37]</sup>	2017	1,820	> 65	25.9
Iceland	Reykjavik <sup>[38]</sup>	2016	255	> 60	27
Spain	Igualada <sup>[39]</sup>	2017	792	> 69	27.3
Estonia	Health Insurance Fund <sup>[40]</sup>	2017	8,298	>50	28.3
Croatia	Zagreb <sup>[41]</sup>	2017	236	> 18	28.4
Belgium	Brussels <sup>[42]</sup>	2016	286	> 64	28.7
Turkey	Ankara <sup>[43]</sup>	2016	124	> 23	29.2
Germany	Trauma Registries <sup>[44]</sup>	2017	231	> 70	31.4
Hungary	euroHOPE Registry <sup>[17]</sup>	2015	-	-	34.8
Greece	Pilot Program	-	-	-	-
Slovenia	Pilot Program	-	-	-	-

was 17.9% (10 studies). The figures for the United States and Australia were 21% and 24.9% respectively. The figure in South America was 26.8%, however, this originated from small studies. On analysis of data from all 36 studies ( $n = 229851$  patients), the mean 1-year global mortality rate was 22% (SD 7.2%, median 22.8%).

These results are at odds with those reported previously and may suggest a consistent global improvement in longer-term survival following hip fracture. Best practice guidelines have been developed by way of continuous optimization through the auditing process of the national registries and continue to be developed, refined or discarded based on continuous audit cycle. The improvement in longer term survival post hip fracture may be secondary to worldwide adoption of current best practice in the care and secondary prevention of hip fracture.

This is the first study that has reviewed recent worldwide-published data on 1-year mortality rates including data collected from 229851 patients. Although the literature review included studies within the last 5 years, it is worth noting that the vast majority (83%) of the local studies (30/36) were published within the last 3 years (2015-2017). It is also worth noting that the most recent papers on this topic (published in 2018) confirm that the changing trend in 1-year mortality continues to be observed<sup>[61-63]</sup>. Cancio *et al*<sup>[64]</sup> this year published a 1-year mortality rate of 22% for a large hip fracture cohort ( $n = 30552$ ), which is similar to our findings described above.

Whilst this article reports a decreasing trend in mortality at 1-year post hip fracture, it is important to recognize limitations of this study. A potential source of error is the use of data from all 36 studies to ascertain a mean mortality rate as some of the studies included patients < 60 years of age. However, limiting the inclusion criteria to those studies that include patients > 60 years only, the mortality rate is 21.8% (SD 7.6%, median 23.6%) which is a similar finding. Another limitation is equal weighting to mortality rates despite wide variation in cohort numbers. Limiting the inclusion criteria to those studies which include > 10001 patients only ( $n = 7$ ), the mean mortality rate is 20.7% (SD 2.4%, median 20.5%) which also is a similar finding.

It is also worth noting that as an index hip fracture outcome measure, mortality is an imperfect quality indicator and is affected by both patient age and casemix (multiple comorbidities typically). Therefore, it is the trend in mortality rate data over time that is more useful as a quality indicator.

**Table 3 Data from local hip fracture studies in Asia**

Country	Origin	Published	n	Age	1-Yr (%)
Japan	Tsukuba <sup>[45]</sup>	2017	110	> 60	2.4
Thailand	Bangkok <sup>[46]</sup>	2015	120	> 50	9.2
South Korea	Daejeon <sup>[47]</sup>	2017	402	> 60	9.5
Hong Kong	Laichikok/Shamshuipo <sup>[48]</sup>	2017	43830	> 65	16.8
Taiwan	Kaohsiun <sup>[49]</sup>	2017	5442	> 60	16.8
Kazachstan	Almaty <sup>[50]</sup>	2017	398	> 50	21
Iran	Tehran <sup>[51]</sup>	2017	1015	> 50	22.4
China	Beijing <sup>[52]</sup>	2016	4504	> 60	23.4
Saudi Arabia	Al Khobar/Dammam <sup>[53]</sup>	2017	203	-	27
India	New Delhi <sup>[54]</sup>	2013	188	> 60	30.4
Lebanon	Pilot Program	-	-	-	-

Osteoporosis currently causes approximately 9 million fractures annually around the world<sup>[65]</sup>. As alluded to in the introduction, hip fractures are increasing with excess of 6 million per year predicted in 2050. The greatest increases are expected in Asia<sup>[66]</sup>. The World Health Organisation predicts that between 2000 and 2050, the proportion of those aged > 60 will double in numbers from 11 to 22% or 605 million to 2 billion people<sup>[67]</sup>. It is important that this expected rise is planned for and this literature review suggests that as a global orthopaedic and orthogeriatric community, vast strides are being made in the optimisation of care with hip fracture patients to mitigate for the impending surge in hip fracture incidence. The reduction in 1-year mortality provides feedback that following adoption of hip fracture management standards of care, it is not simply improvements in early mortality with deferral of mortality outside of the acute window (30 d) but longer-term survival at 1-year post hip fracture. The United Kingdom National Hip Fracture Database website publishes monthly on overall performance measures including national average 30-d mortality. The most recent figure was 5.7% (July, 2018)<sup>[68]</sup> with almost month-to-month reductions since initially published in April 2012. These findings are very positive. Another recent positive development in hip fracture care is the increasing involvement of orthogeriatricians in the care of these fragile patients and their corresponding impact on mortality reduction<sup>[69]</sup>.

The data collected by the national registries allow us as a community to tackle a complex international problem to minimise morbidity, mortality and costs associated with hip fractures. These registries continue to be optimised and it has been recently suggested that (1) simplicity; (2) continuous collection of all cases as routine practice; (3) definition of variables collected; and (4) a common set of variables collected will fulfil the criteria of a model registry which should allow direct national and international comparison and further improvements in care delivered<sup>[3]</sup>.

In conclusion, globally, those which suffer hip fractures may currently be benefiting from 30 years of national registries, rigorous audit processes and international collaboration. This literature review suggests a decrease in trends in mortality at 1-year post hip fracture compared to previous research on this topic. Further improvements in hip fracture care quality indicators are expected by the authors in light of increased involvement of the orthogeriatric subspeciality and the continuous refinement process of the national registries. The results of 7 novel national pilot registries are anticipated.

**Table 4** Data from local hip fracture studies in Oceania

Country	Origin	Published	n	Age	1-Yr (%)
Australia	New South Wales <sup>[55]</sup>	2017	27888	> 65	24.9
New Zealand	-	-	-	-	-

**Table 5** Data from local hip fracture studies in North America

Country	Origin	Published	n	Age	1-Yr (%)
United States	Kaiser Permanente <sup>[56]</sup>	2017	14294	> 60	21
Canada	-	-	-	-	-

**Table 6** Data from local hip fracture studies in South America

Country	Origin	Published	n	Age	1-Yr (%)
Brazil	Canoas <sup>[57]</sup>	2017	213	> 65	23.6
Mexico	Toluca <sup>[58]</sup>	2014	100	> 65	30
Argentina	Pilot Program <sup>[59]</sup>	-	-	-	-

## ARTICLE HIGHLIGHTS

### Research background

Hip fracture audit is increasing globally with 8 countries producing national reports annually and several countries currently piloting hip fracture audit schemes. Hip fracture audit has been shown to result in improvements in care and survival of older people. Short-term mortality rates (in-patient and 30-d) are published by the majority of national registries and there is evidence that short term mortality rates are decreasing. The 1-year mortality rate post hip fracture is currently recognized as approximately 30%. Recent longer-term (1-year) mortality information is lacking in the literature.

### Research motivation

Our research group believes there may also be a changing trend (decreasing) in 1-year mortality post hip fracture. Knowledge of a more current, up-to-date figure will lead to a more informed discussion with family of those affected and optimization of care of this fragile cohort at hospital and national levels.

### Research objectives

To determine the current 1-year mortality rate post hip fracture.

### Research methods

Systematic literature review. The following keywords were used: "hip fracture", "mortality", "year", "death", "outcomes", "registry", "registries", "database" and "databases" (PubMed and Google Scholar). The inclusion criteria were those studies published within the last 5 years (2013-2017), studies including a cohort >100 and those in the English language.

### Research results

Data from 8 national registries and 36 countries (6 continents) were reviewed. Recently published 1-year mortality rates appear lower than traditional figures and suggest a downward trend.

### Research conclusions

The previously quoted 1-year mortality rate of approximately 30% appears to be outdated. A figure of 22% is a more accurate, up-to-date figure for 1-year mortality rate post hip fracture.

### Research perspectives

This research provides a more current, up-to-date figure for 1-year mortality rate post hip fracture.

## REFERENCES

- 1 Irish Hip Fracture Database National Report 2016 [Internet]. Available from: <https://www.nhfd.co.uk/files/2017ReportFiles/NHFD-AnnualReport2017.pdf>
- 2 National Hip Fracture Database Annual Report 2017. Available from: <https://www.nhfd.co.uk/files/2017ReportFiles/NHFD-AnnualReport2017.pdf>
- 3 Sáez-López P, Brañas F, Sánchez-Hernández N, Alonso-García N, González-Montalvo JI. Hip fracture registries: utility, description, and comparison. *Osteoporos Int* 2017; **28**: 1157-1166 [PMID: 27872956 DOI: 10.1007/s00198-016-3834-x]
- 4 Important Facts about Falls 2017. Centre for Disease Control and Prevention. Available from: <https://www.cdc.gov/homeandrecreationalafety/falls/adultfalls.html>
- 5 Johansen A, Golding D, Brent L, Close J, Gjertsen JE, Holt G, Hommel A, Pedersen AB, Röck ND, Thorngren KG. Using national hip fracture registries and audit databases to develop an international perspective. *Injury* 2017; **48**: 2174-2179 [PMID: 28803651 DOI: 10.1016/j.injury.2017.08.001]
- 6 Gullberg B, Johnell O, Kanis JA. World-wide projections for hip fracture. *Osteoporos Int* 1997; **7**: 407-413 [PMID: 9425497]
- 7 Leal J, Gray AM, Prieto-Alhambra D, Arden NK, Cooper C, Javaid MK, Judge A; REFRESH study group. Impact of hip fracture on hospital care costs: a population-based study. *Osteoporos Int* 2016; **27**: 549-558 [PMID: 26286626 DOI: 10.1007/s00198-015-3277-9]
- 8 International Osteoporosis Foundation. Available from: <http://www.capturethefracture.org/health-economics>
- 9 Bentler SE, Liu L, Obrizan M, Cook EA, Wright KB, Geweke JF, Chrischilles EA, Pavlik CE, Wallace RB, Ohsfeldt RL, Jones MP, Rosenthal GE, Wolinsky FD. The aftermath of hip fracture: discharge placement, functional status change, and mortality. *Am J Epidemiol* 2009; **170**: 1290-1299 [PMID: 19808632 DOI: 10.1093/aje/kwp266]
- 10 Moore DP, Quinlan W. Mortality and morbidity associated with hip fractures. *Ir J Med Sci* 1989; **158**: 40-42 [PMID: 2745031]
- 11 Patel NK, Sarraf KM, Joseph S, Lee C, Middleton FR. Implementing the National Hip Fracture Database: An audit of care. *Injury* 2013; **44**: 1934-1939 [PMID: 23680283 DOI: 10.1016/j.injury.2013.04.012]
- 12 Thorngren KG. National registration of hip fractures. In: European Instructional Lectures. 10th EFORT Congress; Vienna, Austria 2009; 11-18
- 13 Ferguson KB, Halai M, Winter A, Elswood T, Smith R, Hutchison JD, Holt G. National audits of hip fractures: Are yearly audits required? *Injury* 2016; **47**: 439-443 [PMID: 26657888 DOI: 10.1016/j.injury.2015.11.018]
- 14 National Health Services Scotland TSHFA. National Services Scotland 2017. Available from: <http://www.shfa.scot.nhs.uk/About/index.html>
- 15 Heikkinen T, Partanen J, Ristiniemi J, Jalovaara P. Evaluation of 238 consecutive patients with the extended data set of the Standardised Audit for Hip Fractures in Europe (SAHFE). *Disabil Rehabil* 2005; **27**: 1107-1115 [PMID: 16278179 DOI: 10.1080/09638280500061220]
- 16 Gjertsen JE, Engesaeter LB, Furnes O, Havelin LI, Steindal K, Vinje T, Fevang JM. The Norwegian Hip Fracture Register: experiences after the first 2 years and 15,576 reported operations. *Acta Orthop* 2008; **79**: 583-593 [PMID: 18839363 DOI: 10.1080/17453670810016588]
- 17 Medin E, Goude F, Melberg HO, Tediosi F, Belicza E, Peltola M; EuroHOPE study group. European Regional Differences in All-Cause Mortality and Length of Stay for Patients with Hip Fracture. *Health Econ* 2015; **24** Suppl 2: 53-64 [PMID: 26633868 DOI: 10.1002/hec.3278]
- 18 Fragility Fracture Network of the Bone and Joint Decade. Fragility Fracture Network 2017. Available from: <http://www.fragilityfracturenetwork.org/about-2/other-leading-organisations/international/bone-and-joint-decade/>
- 19 Annual Report 2017 [Internet]. ANZHFR 2017. Available from: <https://anzhfr.org/2017-annual-report/>
- 20 Moran CG, Wenn RT, Sikand M, Taylor AM. Early mortality after hip fracture: is delay before surgery important? *J Bone Joint Surg Am* 2005; **87**: 483-489 [PMID: 15741611 DOI: 10.2106/JBJS.D.01796]
- 21 Neuburger J, Currie C, Wakeman R, Tsang C, Plant F, De Stavola B, Cromwell DA, van der Meulen J. The impact of a national clinician-led audit initiative on care and mortality after hip fracture in England: an external evaluation using time trends in non-audit data. *Med Care* 2015; **53**: 686-691 [PMID: 26172938 DOI: 10.1097/MLR.0000000000000383]
- 22 RIKSHOFT / ARSRAPPORT 2016. Available from: [http://rikshoft.se/wp-content/uploads/2013/07/rikshoft\\_rapport2016.pdf](http://rikshoft.se/wp-content/uploads/2013/07/rikshoft_rapport2016.pdf)
- 23 The Scottish Hip Fracture Audit. Hip Fracture Monthly Report 2016. Available from: <http://www.shfa.scot.nhs.uk/>
- 24 Registry DMHF. Annual Report 2016. Available from: [https://www.sundhed.dk/content/cms/62/4662\\_hofte-fraktur\\_%C3%A5rsrapport-2016.pdf](https://www.sundhed.dk/content/cms/62/4662_hofte-fraktur_%C3%A5rsrapport-2016.pdf)
- 25 Norwegian National Advisory Unit on Arthroplasty and Hip Fractures. Annual Report 2016. Available from: [http://nrlweb.ihelse.net/eng/Rapporter/Report2016\\_english.pdf](http://nrlweb.ihelse.net/eng/Rapporter/Report2016_english.pdf)
- 26 Henderson CY, Shanahan E, Butler A, Lenehan B, O'Connor M, Lyons D, Ryan JP. Dedicated orthogeriatric service reduces hip fracture mortality. *Ir J Med Sci* 2017; **186**: 179-184 [PMID: 27059996 DOI: 10.1007/s11845-016-1453-3]
- 27 Hommel AAP, Dept. of Health Sciences, Lund University, Lund, Sweden. Local 1-Year Mortality Information Request. 2017
- 28 Kristensen MT, Kehlet H. The basic mobility status upon acute hospital discharge is an independent risk factor for mortality up to 5 years after hip fracture surgery. *Acta Orthop* 2018; **89**: 47-52 [PMID: 28946781 DOI: 10.1080/17453674.2017.1382038]
- 29 Suhm N, Kaelin R, Studer P, Wang Q, Kressig RW, Rikli D, Jakob M, Pretto M. Orthogeriatric care pathway: a prospective survey of impact on length of stay, mortality and institutionalisation. *Arch Orthop Trauma Surg* 2014; **134**: 1261-1269 [PMID: 25060921 DOI: 10.1007/s00402-014-2057-x]
- 30 Brozek W, Reichardt B, Kimberger O, Zwerina J, Dimai HP, Kritsch D, Klaushofer K, Zwettler E. Mortality after hip fracture in Austria 2008-2011. *Calcif Tissue Int* 2014; **95**: 257-266 [PMID: 24989776 DOI: 10.1007/s00223-014-9889-9]
- 31 Marques A, Lourenço Ó, da Silva JA. Portuguese Working Group for the Study of the Burden of Hip Fractures in Portugal. The burden of osteoporotic hip fractures in Portugal: costs, health related quality of life and mortality. *Osteoporos Int* 2015; **26**: 2623-2630 [PMID: 25986386 DOI: 10.1007/s00198-015-3171-5]
- 32 Klop C, Welsing PM, Cooper C, Harvey NC, Elders PJ, Bijlsma JW, Leufkens HG, de Vries F. Mortality

- in British hip fracture patients, 2000-2010: a population-based retrospective cohort study. *Bone* 2014; **66**: 171-177 [PMID: 24933345 DOI: 10.1016/j.bone.2014.06.011]
- 33 **Poenaru DV**, Prejbeanu R, Iulian P, Haragus H, Popovici E, Golet I, Vermesan D. Epidemiology of osteoporotic hip fractures in Western Romania. *Int Orthop* 2014; **38**: 2329-2334 [PMID: 25069426 DOI: 10.1007/s00264-014-2407-x]
- 34 **Tucker A**, Donnelly KJ, McDonald S, Craig J, Foster AP, Acton JD. The changing face of fractures of the hip in Northern Ireland: a 15-year review. *Bone Joint J* 2017; **99-B**: 1223-1231 [PMID: 28860404 DOI: 10.1302/0301-620X.99B9.BJJ-2016-1284.R1]
- 35 **Folbert EC**, Hegeman JH, Vermeer M, Regtuijt EM, van der Velde D, Ten Duis HJ, Slaets JP. Improved 1-year mortality in elderly patients with a hip fracture following integrated orthogeriatric treatment. *Osteoporos Int* 2017; **28**: 269-277 [PMID: 27443570 DOI: 10.1007/s00198-016-3711-7]
- 36 **Grech S CS**. Osteoporotic hip fractures - Three year follow up mortality rate in Malta. *Malta Medical Journal* 2016; **28**: 5-10
- 37 **Haugan K**, Johnsen LG, Basso T, Foss OA. Mortality and readmission following hip fracture surgery: a retrospective study comparing conventional and fast-track care. *BMJ Open* 2017; **7**: e015574 [PMID: 28851773 DOI: 10.1136/bmjopen-2016-015574]
- 38 **Magnusson KA**, Gunnarsson B, Sigurdsson GH, Mogensen B, Olafsson Y, Karason S. [Treatment and outcome of patients with hip fracture]. *Laeknabladid* 2016; **102**: 119-125 [PMID: 26985589 DOI: 10.17992/ibl.2016.03.69]
- 39 **Duaso E**, Formiga F, Marimón P, Sandiumenge M, Salgado MT, Murga V, Gamboa A, Rodriguez C, Castellà J, Escalante E, Lumbreras C, Tarrida A, Sellarès R. Advantages of care for patients with hip fractures in the acute geriatric unit: Hip study Anoa. *Geriatr Gerontol Int* 2018; **18**: 407-414 [PMID: 29139194 DOI: 10.1111/ggi.13191]
- 40 **Jürisson M**, Raag M, Kallikorm R, Lember M, Uusküla A. The impact of comorbidities on hip fracture mortality: a retrospective population-based cohort study. *Arch Osteoporos* 2017; **12**: 76 [PMID: 28849347 DOI: 10.1007/s11657-017-0370-z]
- 41 **Gulin T**, Kruljac I, Kirigin Biloš LS, Gulin M, Grgurević M, Borojević M. The role of adipokines as prognostic factors of one-year mortality in hip fracture patients. *Osteoporos Int* 2017; **28**: 2475-2483 [PMID: 28501890 DOI: 10.1007/s00198-017-4068-2]
- 42 **Forget P**, Dillien P, Engel H, Cornu O, De Kock M, Yombi JC. Use of the neutrophil-to-lymphocyte ratio as a component of a score to predict postoperative mortality after surgery for hip fracture in elderly subjects. *BMC Res Notes* 2016; **9**: 284 [PMID: 27230508 DOI: 10.1186/s13104-016-2089-0]
- 43 **Kilci O**, Un C, Sacan O, Gamli M, Baskan S, Baydar M, Ozkurt B. Postoperative Mortality after Hip Fracture Surgery: A 3 Years Follow Up. *PLoS One* 2016; **11**: e0162097 [PMID: 27788137 DOI: 10.1371/journal.pone.0162097]
- 44 **Förch S**, Kretschmer R, Haufe T, Plath J, Mayr E. Orthogeriatric Combined Management of Elderly Patients With Proximal Femoral Fracture: Results of a 1-Year Follow-Up. *Geriatr Orthop Surg Rehabil* 2017; **8**: 109-114 [PMID: 28540117 DOI: 10.1177/2151458517698536]
- 45 **Yoshizawa T**, Nishino T, Mishima H, Ainoya T, Yamazaki M. Rehabilitation in a convalescent rehabilitation ward following an acute ward improves functional recovery and mortality for hip fracture patients: a sequence in a single hospital. *J Phys Ther Sci* 2017; **29**: 1102-1107 [PMID: 28626336 DOI: 10.1589/jpts.29.1102]
- 46 **Amphansap T**, Nitiwarangkul L. One year mortality after osteoporotic hip fractures and associated risk factors in Police General Hospital. *Osteoporosis and Sarcopenia* 2015; **1**: 75-9
- 47 **Cha YH**, Ha YC, Yoo JI, Min YS, Lee YK, Koo KH. Effect of causes of surgical delay on early and late mortality in patients with proximal hip fracture. *Arch Orthop Trauma Surg* 2017; **137**: 625-630 [PMID: 28321571 DOI: 10.1007/s00402-017-2674-2]
- 48 **Liu SK**, Ho AW, Wong SH. Early surgery for Hong Kong Chinese elderly patients with hip fracture reduces short-term and long-term mortality. *Hong Kong Med J* 2017; **23**: 374-380 [PMID: 28655864 DOI: 10.12809/hkmj165005]
- 49 **Lee TC**, Ho PS, Lin HT, Ho ML, Huang HT, Chang JK. One-Year Readmission Risk and Mortality after Hip Fracture Surgery: A National Population-Based Study in Taiwan. *Aging Dis* 2017; **8**: 402-409 [PMID: 28840055 DOI: 10.14336/AD.2016.1228]
- 50 **Amrayev S**, AbuJazar U, Stucinskas J, Smailys A, Tarasevicius S. Outcomes and mortality after hip fractures treated in Kazakhstan. *Hip Int* 2018; **28**: 205-209 [PMID: 29890912 DOI: 10.1177/1120700018773395]
- 51 **Vosoughi AR**, Emami MJ, Pourabbas B, Mahdaviyazad H. Factors increasing mortality of the elderly following hip fracture surgery: role of body mass index, age, and smoking. *Musculoskelet Surg* 2017; **101**: 25-29 [PMID: 27766497 DOI: 10.1007/s12306-016-0432-1]
- 52 **Li S**, Sun T, Liu Z. Excess mortality of 1 year in elderly hip fracture patients compared with the general population in Beijing, China. *Arch Osteoporos* 2016; **11**: 35 [PMID: 27785743 DOI: 10.1007/s11657-016-0289-9]
- 53 **Sadat-Ali M**, Alfaraidy M, AlHawas A, Al-Othman AA, Al-Dakheel DA, Tayara BK. Morbidity and mortality after fragility hip fracture in a Saudi Arabian population: Report from a single center. *J Int Med Res* 2017; **45**: 1175-1180 [PMID: 28480812 DOI: 10.1177/0300060517706283]
- 54 **Dhanwal DK**. Mortality and Functional Impairment after Hip Fracture - A Prospective Study in Indian Population. In: ESCE013-IOF Congress; April 18 2013; Rome, Italy 2013
- 55 **Mitchell R**, Harvey L, Brodaty H, Draper B, Close J. One-year mortality after hip fracture in older individuals: the effects of delirium and dementia. *Arch Gerontol Geriatr* 2017; **72**: 135-141 [PMID: 28628893 DOI: 10.1016/j.archger.2017.06.006]
- 56 **Okike K**, Chan PH, Paxton EW. Effect of Surgeon and Hospital Volume on Morbidity and Mortality After Hip Fracture. *J Bone Joint Surg Am* 2017; **99**: 1547-1553 [PMID: 28926384 DOI: 10.2106/JBJS.16.01133]
- 57 **Guerra MT**, Viana RD, Feil L, Feron ET, Maboni J, Vargas AS. One-year mortality of elderly patients with hip fracture surgically treated at a hospital in Southern Brazil. *Rev Bras Ortop* 2016; **52**: 17-23 [PMID: 28194376 DOI: 10.1016/j.rboe.2016.11.006]
- 58 **Negrete-Corona J**, Alvarado-Soriano JC, Reyes-Santiago LA. Hip fracture as risk factor for mortality in patients over 65 years of age. Case-control study. *Acta Ortop Mex* 2014; **28**: 352-362 [PMID: 26016287]
- 59 **Benchimol J**, Fiorentini F, Elizondo CM, Boietti BR, Carabelli G, Barla J, Sancineto C, Waisman GD, Giunta DH. Institutional Registry of Elderly Patients With Hip Fracture in a Community-Based Tertiary Care Hospital in Argentina (RIAFC). *Geriatr Orthop Surg Rehabil* 2016; **7**: 121-125 [PMID: 27551569]

- DOI: [10.1177/2151458516651309](https://doi.org/10.1177/2151458516651309)]
- 60 **Neuburger J**, Currie C, Wakeman R, Johansen A, Tsang C, Plant F, Wilson H, Cromwell DA, van der Meulen J, De Stavola B. Increased orthogeriatrician involvement in hip fracture care and its impact on mortality in England. *Age Ageing* 2017; **46**: 187-192 [PMID: [27915229](https://pubmed.ncbi.nlm.nih.gov/27915229/) DOI: [10.1093/ageing/afw201](https://doi.org/10.1093/ageing/afw201)]
- 61 **Reguant F**, Arnau A, Lorente JV, Maestro L, Bosch J. Efficacy of a multidisciplinary approach on postoperative morbidity and mortality of elderly patients with hip fracture. *J Clin Anesth* 2018; **53**: 11-19 [PMID: [30286380](https://pubmed.ncbi.nlm.nih.gov/30286380/) DOI: [10.1016/j.jclinane.2018.09.029](https://doi.org/10.1016/j.jclinane.2018.09.029)]
- 62 **Zanetti M**, Gortan Cappellari G, Ratti C, Ceschia G, Murena L, De Colle P, Barazzoni R. Poor nutritional status but not cognitive or functional impairment per se independently predict 1 year mortality in elderly patients with hip-fracture. *Clin Nutr* 2018 [PMID: [30217470](https://pubmed.ncbi.nlm.nih.gov/30217470/) DOI: [10.1016/j.clnu.2018.08.030](https://doi.org/10.1016/j.clnu.2018.08.030)]
- 63 **Chow SK**, Qin JH, Wong RM, Yuen WF, Ngai WK, Tang N, Lam CY, Lau TW, Lee KB, Siu KM, Wong SH, Zhu TY, Cheung WH, Leung KS. One-year mortality in displaced intracapsular hip fractures and associated risk: a report of Chinese-based fragility fracture registry. *J Orthop Surg Res* 2018; **13**: 235 [PMID: [30217215](https://pubmed.ncbi.nlm.nih.gov/30217215/) DOI: [10.1186/s13018-018-0936-5](https://doi.org/10.1186/s13018-018-0936-5)]
- 64 **Cancio JM**, Vela E, Santaugènia S, Clèries M, Inzitari M, Ruiz D. Influence of demographic and clinical characteristics of elderly patients with a hip fracture on mortality: A retrospective, total cohort study in North-East Spain. *Bone* 2018; **117**: 123-129 [PMID: [30223133](https://pubmed.ncbi.nlm.nih.gov/30223133/) DOI: [10.1016/j.bone.2018.09.002](https://doi.org/10.1016/j.bone.2018.09.002)]
- 65 **Johnell O**, Kanis JA. An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporos Int* 2006; **17**: 1726-1733 [PMID: [16983459](https://pubmed.ncbi.nlm.nih.gov/16983459/) DOI: [10.1007/s00198-006-0172-4](https://doi.org/10.1007/s00198-006-0172-4)]
- 66 **Cooper C**, Campion G, Melton LJ. Hip fractures in the elderly: a world-wide projection. *Osteoporos Int* 1992; **2**: 285-289 [PMID: [1421796](https://pubmed.ncbi.nlm.nih.gov/1421796/)]
- 67 **World Health Organisation**. Facts on Ageing 2017. Available from: <http://www.who.int/ageing/about/facts/en/>
- 68 **National Hip Fracture Database**. Charts and Reports 2017. Available from: <https://www.nhfd.co.uk/20/nhfdcharts.nsf/vwCharts/OverallPerformance>

**P- Reviewer:** Regis D; Surace MF; Ünver B

**S- Editor:** Dou Y **L- Editor:** A **E- Editor:** Wu YXJ





Published By Baishideng Publishing Group Inc  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

