

Match Overview

Rank	Source	Words	Similarity
1	Crossref	47 words	2%
	Rosalie A. Carr, Michele T. Yip-Schneider, Rachel E. Simpson, Scott Dolejs et al. "Pancreatic cyst fluid glucose: rapid ..."		
2	Internet	27 words crawled on 02-Feb-2017 pdfs.semanticscholar.org	1%
3	Crossref	25 words	1%
	"Pancreatic Masses", Springer Nature America, Inc, 2016		
4	Crossref	23 words	1%
	Tang, Raymond S., and Thomas J. Savides. "Pancreatic Cystic Lesions", <i>Gastrointestinal Endoscopy in the Cancer Pa</i>		
5	Crossref	18 words	1%
	Matthew Kulzer, Aatur D. Singhi, Alessandro Furlan, Matthew T. Heller et al. "Current concepts in molecular genetics a		
6	Crossref	13 words	1%
	Atsuhiko Masuda, Yoshifumi Arisaka, Shigeo Hara, Ippei Matsumoto et al. "MUC2 expression and prevalence of high-		
7	Internet	12 words crawled on 22-Mar-2016 www.europeanreview.org	1%
8	Crossref	11 words	<1%
	Walter Gwang-Up Park. "Diagnostic Performance of Cyst Fluid Carcinoembryonic Antigen and Amylase in Histological		
9	Crossref	10 words	<1%
	Barresi, Luca. "Pancreatic cystic lesions: How endoscopic ultrasound morphology and endoscopic ultrasound fine r ..."		

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 45597

Manuscript type: OPINION REVIEW

Cyst fluid glucose: An alternative to carcinoembryonic antigen for pancreatic mucinous cysts

Lopes CV. Cyst fluid glucose

César Vivian Lopes

Abstract

Pancreatic cystic lesions (PCLs) have been increasingly recognized in clinical practice. Although inflammatory cysts (pseudocysts) are the most common

找到约 3,440 条结果 (用时 0.72 秒)

Pancreatic cyst fluid glucose: rapid, inexpensive, and accurate ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/29241991> - 翻译此页

作者: RA Carr - 2018 - 被引用次数: 2 - 相关文章

2017年12月11日 - **Pancreatic cyst fluid glucose**: rapid, inexpensive, and accurate **diagnosis of mucinous pancreatic cysts**. ... differentiation of **mucinous** from benign, **nonmucinous pancreatic cysts** is **cyst fluid carcinoembryonic antigen**. However ... **cyst fluid glucose** may be an **attractive alternative to carcinoembryonic antigen**.

缺少字词: differential

Pancreatic cyst fluid glucose: Rapid, inexpensive, and ... - ResearchGate

https://www.researchgate.net/.../321737484_Pancreatic_cyst_fluid_glucose_... - 翻译此页

2018年10月19日 - Request PDF on ResearchGate | **Pancreatic cyst fluid glucose**: Rapid, inexpensive, and accurate **diagnosis of mucinous pancreatic cysts** ... of **mucinous** from benign, **nonmucinous pancreatic cysts** is **cyst fluid carcinoembryonic antigen**. ... may be an **attractive alternative to carcinoembryonic antigen**.

P.08.8 PANCREATIC CYSTIC FLUID GLUCOSE AND CEA ...

https://www.researchgate.net/.../323865962_P088_PANCREATIC_CYSTIC_... - 翻译此页

... and others published **P.08.8 PANCREATIC CYSTIC FLUID GLUCOSE AND CEA ... NON-MUCINOUS FROM PRE-MALIGNANT MUCINOUS PANCREATIC CYSTS**. ... with cytology and viscosity for the **differential diagnosis** of **pancreatic cysts**. Methods: ... that **pancreatic cyst fluid glucose** may be an **attractive alternative to** ...

Pancreatic cyst fluid glucose: rapid, inexpensive, and accurate ...

<https://scholarworks.iupui.edu/handle/1805/17337> ▾ 翻译此页

作者: RA Carr - 2018 - 被引用次数: 2 - 相关文章

Keywords: **cyst fluid glucose** ; **mucinous pancreatic cysts** ; **diagnostic testing** ... differentiation of **mucinous** from benign, **nonmucinous pancreatic cysts** is **cyst fluid** ... **cyst fluid glucose** may be an



All

Images

Videos

翻译成中文

关闭取词

1,630 Results

Any time ▾

Pancreatic cyst fluid glucose: rapid, inexpensive, and ...

<https://www.sciencedirect.com/science/article/pii/S0039606017307043>

The most widely accepted biochemical test for preoperative differentiation of **mucinous** from benign, nonmucinous **pancreatic cysts** is **cyst fluid carcinoembryonic antigen**. However, the diagnostic accuracy of **carcinoembryonic antigen** ranges from 70% to 86%.

Cited by: 1

Author: Rosalie A. Carr, Rosalie A. Carr, Michele ...

Publish Year: 2017

Cyst fluid analysis in the differential diagnosis of ...

https://www.researchgate.net/publication/7648689_Cyst_fluid...

Conclusion: **Cyst fluid glucose** has significant advantages over **carcinoembryonic antigen** and should be considered for use as a routine diagnostic test for **pancreatic mucinous cysts**. View Show abstract

Cytopathological Analysis of Cyst Fluid Enhances ...

www.ncbi.nlm.nih.gov › ... › [Medicine \(Baltimore\)](#) › v.94(24); 2015 Jun

Pancreatic Cyst Fluid . After EUS-FNA of a **pancreatic cyst**, **carcinoembryonic antigen** (CEA) levels were measured in **pancreatic cyst fluid**. Furthermore, cytopathology was evaluated for the presence of neoplastic epithelial cells and **mucinous** background.

Cited by: 1

Author: Wesley K. Utomo, Henri Braat, Marco J. ...

Publish Year: 2015

The role of cytopathology and cyst fluid analysis in the ...

onlinelibrary.wiley.com › [Oncology & Radiotherapy](#)

In addition, cytology alone is inferior to **carcinoembryonic antigen** (CEA) analysis of **cyst fluid** for the distinction of **mucinous** and nonmucinous **cyst** type. [5, 6] Treatment options for **pancreatic cysts** vary depending on the **cyst** type and the risk of malignancy.

Published in: [Cancer Cytopathology](#) - 2014

Authors: [Ivan Chebib](#) · [Kurt Yaeger](#) · [Mari Minokenudson](#) · [Martha B Pitman](#)

Affiliation: [Harvard University](#)

About: [Cyst](#) · [Endoscopic ultrasound](#) · [Pancreas](#) · [Neoplasm](#) · [Cytology](#)

Cyst Fluid Glucose is Rapidly Feasible and Accurate in ...

https://www.researchgate.net/publication/277088068_Cyst_Fluid...

With a threshold of ≤ 50 mg/dL, **cyst fluid glucose** was 92% sensitive, 87% specific, and 90% accurate in diagnosing **mucinous pancreatic cysts**. In comparison, **cyst fluid carcinoembryonic antigen**

找到约 51,500 条结果 (用时 0.45 秒)

[Pancreatic cyst fluid glucose: rapid, inexpensive, and accurate ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pubmed/29241991> - [翻译此页](#)

作者: RA Carr - 2018 - 被引用次数: 2 - [相关文章](#)

2017年12月11日 - **Pancreatic cyst fluid glucose**: rapid, inexpensive, and accurate diagnosis of **mucinous pancreatic cysts**. ... work, we hypothesize that **pancreatic cyst fluid glucose** may be an attractive **alternative to carcinoembryonic antigen**.

[Update on pancreatic cyst fluid analysis - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3959935/> - [翻译此页](#)

作者: M Rockacy - 2013 - 被引用次数: 25 - [相关文章](#)

These represent a wide variety of lesions including **mucinous cysts** [intraductal ... **Pancreatic cyst CEA** level is considered the most accurate tumor marker for ... A **CEA**<5 had **similar** performance characteristics (sensitivity 50%, specificity 95%, ...

[Abstract](#) · [Introduction](#) · [Tumor markers](#) · [Concluding remarks](#)

[Analysis of pancreatic cyst fluid - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5145907/> - [翻译此页](#)

作者: S Ngamruengphong - 2016 - 被引用次数: 9 - [相关文章](#)

2017年12月1日 - **Pancreatic cysts** are extremely common, and are identified in between 2% to ...
Keywords: molecular markers, **pancreatic cyst**, **cyst fluid**, intraductal papillary **mucinous** neoplasm, serous ... However, other groups have proposed **alternative** cutoffs. ... Some studies have suggested that a high **cyst fluid CEA** is ...

[\[PDF\] Cyst Fluid Glucose is Rapidly Feasible and Accurate in ... - acgblog](#)

acgblog.org/wp-content/uploads/2015/06/Pancreatic-Cysts.pdf - [翻译此页](#)

2015 - 被引用次数: 16 - [相关文章](#)

2015年5月19日 - nous **cystic** neoplasms and intraductal papillary **mucinous** neoplasms (IPMNs). ... analysis of **pancreatic cyst fluid**, **glucose** appeared to accurately differentiate All three **glucose** assays and **CEA** were **similar** in differentiat-.

[Pancreatic cyst fluid glucose: rapid, inexpensive, and ... - Surgery](#)