



ADAMTS13 and von Willebrand factor are useful biomarkers for sorafenib t



网页 图片 视频 学术 词典 地图

检测到您输入了英文，试试切换到国际版？搜英文结果更丰富更准确 >

82 条结果 时间不限

### Mutations in Coagulation Factor VIII Are Associated ... 翻译此页

2017-10-28 · Coagulation factor VIII (FVIII), von Willebrand factor (VWF), and ADAMTS13 (a disintegrin and metalloprotease with thrombospondin type 1 repeats 13) play an important role in the regulation of ... [https://www.researchgate.net/publication/320573335\\_Mutations\\_in...](https://www.researchgate.net/publication/320573335_Mutations_in...)

### The role of von Willebrand factor as a biomarker of ... 翻译此页

The role of von Willebrand factor as a biomarker of tumor development in hepatitis B virus-associated human hepatocellular carcinoma: A quantitative proteomic based study ... despite sorafenib ... [https://www.researchgate.net/publication/261920137\\_The\\_role\\_of\\_von...](https://www.researchgate.net/publication/261920137_The_role_of_von...)

### β-gal Publications | PubFacts 翻译此页

Von Willebrand factor (vWF) is a glycoprotein with a crucial role in the formation of platelet thrombi, and ADAMTS13 is the main enzyme responsible for vWF cleavage. Both are important in the relationship between diabetic nephropathy, hypercoagulability, and cardiovascular disease. <https://www.pubfacts.com/search/β-gal/1>

### guias EVW 2008. | Coagulation | Bleeding - Scribd 翻译此页

guias EVW 2008. - Download as PDF File (.pdf), Text File (.txt) or read online. . Buscar Buscar. ... needed are better knowledge of and treatment options for management of VWD and bleeding or bleeding risk. As documented in this VWD guidelines ... VWF:Ag, von Willebrand factor antigen. Modified from Gill et al. [45]. Used with permission. In this <https://es.scribd.com/document/258843741/guias-EVW-2008>

### JoVE | Peer Reviewed Scientific Video Journal - ... 翻译此页

**Name of Journal:** *World Journal of Gastrointestinal Oncology*

**Manuscript NO:** 45968

**Manuscript Type:** ORIGINAL ARTICLE

*Observational Study*

**ADAMTS13 and von Willebrand factor are useful biomarkers for sorafenib treatment efficiency in patients with hepatocellular carcinoma**

Hiroaki Takaya, Tadashi Namisaki, Naotaka Shimozato, Kosuke Kaji, Mitsuteru Kitade, Kei Moriya, Shinya Sato, Hideto Kawaratani, Takemi Akahane, Masanori Matsumoto, Hitoshi Yoshii

## Match Overview

1	<b>Crossref</b> 123 words Hiroaki Takaya, Hideto Kawaratani, Takuya Kuba, Kenichiro Saki et al. "Diagnosis of Hepatocellular Carcinoma"	4%
2	<b>Internet</b> 72 words crawled on 08-Apr-2018 <a href="http://www.spandidos-publications.com">www.spandidos-publications.com</a>	2%
3	<b>Internet</b> 44 words crawled on 02-Nov-2018 <a href="http://journals.sagepub.com">journals.sagepub.com</a>	1%
4	<b>Crossref</b> 25 words Chenchen Wang, Ying Tong, Yankai Wen, Jiwei Cai et al. "Hepatocellular Carcinoma: A Review"	1%
5	<b>Internet</b> 21 words crawled on 08-Mar-2019 <a href="http://www.jcancer.org">www.jcancer.org</a>	1%
6	<b>Crossref</b> 19 words Nishigori, Naoto, Masanori Matsumoto, Fumiko Koyama, Masaki Hayakawa, Kenta Ueta	1%
7	<b>Crossref</b> 18 words "Posters (Abstracts 301-2389)", Hepatology, 2019	1%
8	<b>Crossref</b> 17 words Shiozawa, Kazuo. "Comparison of percutaneous radiofrequency ablation and CyberKnife"	1%



国内版

国际版

ADAMTS13 and von Willebrand factor are useful biomarkers for sorafenib treatme



All

Images

Videos

翻译成中文

关闭取词

12,300 Results

Any time ▾

### The role of von Willebrand factor as a biomarker of tumor ...

[https://www.researchgate.net/publication/261920137\\_The\\_role\\_of\\_von...](https://www.researchgate.net/publication/261920137_The_role_of_von...)

The role of **von Willebrand factor** as a **biomarker** of tumor development in hepatitis B virus-associated human **hepatocellular carcinoma**: A quantitative proteomic based study

### von Willebrand factor is a useful biomarker for liver ...

[https://www.researchgate.net/publication/325369346\\_von\\_Willebrand...](https://www.researchgate.net/publication/325369346_von_Willebrand...)

Background: Several noninvasive **biomarkers** are available for diagnosing liver fibrosis stage and predicting **hepatocellular carcinoma (HCC)** development in **patients** with chronic hep

### von Willebrand factor is a useful biomarker for liver ...

<https://journals.sagepub.com/doi/abs/10.1177/2050640618779660>

BackgroundSeveral **noninvasive biomarkers** are available for diagnosing **liver fibrosis** stage and predicting hepatocellular carcinoma (HCC) development in **patients** with **chronic hepatitis** and **liver cirrhosis**. However, these **biomarkers** are not sufficiently accurate. Recently, **von Willebrand factor (VWF)** has been related to **angiogenesis** and apoptosis.

**Author:** Hiroaki Takaya, Hideto Kawaratani, Yu... **Publish Year:** 2018

### von Willebrand factor is a useful biomarker for liver ...

[journals.sagepub.com/doi/full/10.1177/2050640618779660](https://journals.sagepub.com/doi/full/10.1177/2050640618779660)

**von Willebrand factor** is a **useful biomarker** for liver fibrosis and prediction of **hepatocellular carcinoma** development in **patients** with hepatitis B and C ... during the observation period was higher than that of **patients** without **hepatocellular carcinoma** at the time of liver biopsy ((a) and (b)). ... Ag may become a **biomarker** of **treatment effects** ...

**Author:** Hiroaki Takaya, Hideto Kawaratani, Yu... **Publish Year:** 2018



国内版

国际版

ADAMTS13 and von Willebrand factor are useful biomarkers for sorafenib treatme



Chat with Bing

Sign in



All

Images

Videos

翻译成中文

关闭取词

Make Bing your search engine >

12,300 Results

Any time ▾

### von Willebrand factor is a useful biomarker for liver ...

[https://www.researchgate.net/publication/325369346\\_von\\_Willebrand...](https://www.researchgate.net/publication/325369346_von_Willebrand...)

Background: Several noninvasive **biomarkers** are available for diagnosing liver fibrosis stage and predicting **hepatocellular carcinoma** (HCC) development in **patients** with chronic hep

### von Willebrand factor is a useful biomarker for liver ...

[journals.sagepub.com/doi/full/10.1177/2050640618779660](https://journals.sagepub.com/doi/full/10.1177/2050640618779660)

Relationship between **von Willebrand factor antigen** levels and hepatocellular carcinoma development. The **von Willebrand factor antigen** levels of **patients** who developed hepatocellular carcinoma with absent to moderate liver **fibrosis** (F0–2) and severe **fibrosis** (F3 and F4) during the observation period was higher than that of **patients** without hepatocellular carcinoma at the time of liver biopsy ((a) and (b)).

**Author:** Hiroaki Takaya, Hideto Kawaratani, Yu... **Publish Year:** 2018

### Sorafenib Inhibits Hypoxia-Inducible Factor-1 $\alpha$ Synthesis ...

[clincancerres.aacrjournals.org/content/18/20/5662](https://clincancerres.aacrjournals.org/content/18/20/5662) ▾

Oct 15, 2012 · Purpose: The overexpression of hypoxia-inducible factor 1 $\alpha$  (HIF-1 $\alpha$ ) is a common finding in hepatocellular carcinoma (HCC), and it leads to angiogenesis and poor prognosis. Sorafenib, a multikinase inhibitor, has shown significant improvement in ...

Published in: **Clinical Cancer Research** · 2012

Authors: Liping Liu · Rocky L K Ho · George G Chen · Paul B S Lai

Affiliation: The Chinese University of Hong Kong

### The role of von Willebrand factor as a biomarker of tumor ...

[https://www.researchgate.net/publication/261920137\\_The\\_role\\_of\\_von...](https://www.researchgate.net/publication/261920137_The_role_of_von...)

The role of **von Willebrand factor** as a **biomarker** of tumor development in hepatitis B virus-associated human **hepatocellular carcinoma**: A quantitative proteomic based study

### Cancer Awareness in Atypical Thrombotic Microangiopathies