

Factors associated with complicated erosive esophagitis: A Japanese multicenter

全部 图片 视频 新闻 地图 图书

找到约 21,800 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Factors associated with complicated erosive esophagitis: A Japanese multicenter, prospective, cross-sectional study](#)

[Epidemiology of gastroesophageal reflux disease in ...](#) - Jung - 被引用次数: 110

[Racial and geographic issues in gastroesophageal ...](#) - Sharma - 被引用次数: 67

[... risk factors for Barrett's esophagus and associated ...](#) - Wong - 被引用次数: 135

[Risk factors for osteoporosis in Japan: is it associated with ...](#) - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4358368/>

6 Mar 2015 ... The aim of this study was to investigate the risk factors for osteoporosis This cross-sectional study investigated the risk of osteoporosis area, no case was complicated with vertebral fracture or kyphosis. Proportion of reflux esophagitis in 6,010 Japanese adults: prospective evaluation by endoscopy.

[Characteristics of Colonic Diverticulitis and Factors Associated With ...](#)

www.ncbi.nlm.nih.gov/pubmed/26544815

Characteristics of Colonic Diverticulitis and Factors Associated With Complications: A Japanese Multicenter, Retrospective, Cross-Sectional Study. Manabe N(1) ...

[References in Environmental – Lifestyle related factors - Best ...](#)

[www.bpgastro.com/article/S1521-6918\(10\)00133-2/references](http://www.bpgastro.com/article/S1521-6918(10)00133-2/references)

MD (Associate Professor) correspondence ... M., Sakamoto, C. Mechanisms of acid gastroesophageal reflux in the Japanese population. ... luminal release of esophageal epidermal growth factor in patients with reflux esophagitis. and non-erosive reflux disease: a nationwide multicentre prospective study in Korea.

[Background Factors of Reflux Esophagitis and Non-Erosive Reflux ...](#)

journals.plos.org/plosone/article?id=10.1371/journal.pone...

26 Jul 2013 ... Conclusions Significantly associated factors of NERD are ... Reflux Disease: A ...

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 30221

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Factors associated with complicated erosive esophagitis: A Japanese multicenter, prospective, cross-sectional study

Sakaguchi M *et al.* Complications of erosive esophagitis

Masahiro Sakaguchi, Noriaki Manabe, Nobuo Ueki, Jun Miwa, Tomoki Inaba, Norimasa Yoshida, Kouichi Sakurai, Masahiro Nakagawa, Hajime Yamada, Michiya Saito, Koji Nakada, Katsuhiko Iwakiri, Takashi Joh, Ken Haruma

Masahiro Sakaguchi, Department of Gastroenterology, Moriguchi Keijinkai Hospital, Osaka 570-0021, Japan

Match Overview

1	Internet 147 words crawled on 29-Sep-2016 www.wjgnet.com	3%
2	Crossref 32 words N. Manabe. "Efficacy of adding sodium alginate to omeprazole in patients with nonerosive reflux disease: a randomiz ...	1%
3	Internet 24 words crawled on 27-Feb-2008 www.medworm.com	1%
4	Crossref 24 words Bruley des Varannes, S.. "Short and long-term PPI treatment for GERD. Do we need more-potent anti-secretory drugs ...	1%
5	Crossref 22 words Shimatani, Tomohiko, Mitsushige Sugimoto, Masafumi Nishino, Kyoichi Adachi, Kenji Furuta, Masanori Ito, Susumu Kur	<1%
6	Crossref 21 words Noriaki Manabe, Ken Haruma. "Optimal acid suppressive treatment for adequate symptom relief and prevention of the	<1%

学术搜索

找到约 537 条结果 (用时0.06秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

☐ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[HTML] Epidemiology of gastroesophageal reflux disease in Asia: a systematic review

HK Jung - 2011 - jnmjournal.org

... H, Iwasaki E, Yokoyama H, Sugino Y, Hibi T. Serum lipid levels are positively associated with non ... Obesity as a risk factor for GERD in Japan. ... Risk factors for gastroesophageal reflux disease, reflux esophagitis and non-erosive reflux disease among Chinese patients undergoing ...

被引用次数: 110 相关文章 所有 12 个版本 引用 保存 更多

[HTML] jnmjournal.org

Racial and geographic issues in gastroesophageal reflux disease

P Sharma, S Wani, Y Romero, D Johnson... - The American journal of ..., 2008 - nature.com

... has increased markedly in recent years (100), appears to be an important risk factor for GERD ... the weight gain (>5 kg, adjusted OR 3.0, 95% CI 1.6-6) was associated with new ... Prevalence of and risk factors for Helicobacter pylori infection in a multi-racial dyspeptic Malaysian ...

被引用次数: 67 相关文章 所有 7 个版本 引用 保存

[HTML] nature.com

Epidemiologic risk factors for Barrett's esophagus and associated adenocarcinoma

A Wong, RC Fitzgerald - Clinical Gastroenterology and Hepatology, 2005 - Elsevier

... In addition, the altered body composition associated with obesity affects the pharmacokinetics of drugs. ... With regard to specific dietary factors, although there have been numerous studies assessing ... They showed that a complex risk factor score (including tobacco use, alcohol, fat ...

被引用次数: 134 相关文章 所有 6 个版本 引用 保存

[HTML] cghjournal.org

[HTML] Geographical and ethnic differences in gastro - oesophageal reflux disease

JY Kang - Alimentary pharmacology & therapeutics, 2004 - Wiley Online Library

... Details of published studies satisfying the inclusion criteria on the prevalence of symptoms associated with GERD ... Although body mass index, age and sex were risk factors for oesophagitis and hiatus hernia ... from 1.02 to 1.11, race was by far the most important risk factor (4.04 for ...

被引用次数: 105 相关文章 所有 4 个版本 引用 保存

[HTML] wiley.com

Environmental-Lifestyle related factors

S Roman, JE Pandolfino, Best Practice & Research Clinical ..., 2010, Elsevier

[HTML] nih.gov

学术搜索

找到约 524 条结果（用时0.08秒）

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序


搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

 创建快讯

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

[HTML] Epidemiology of gastroesophageal reflux disease in Asia: a systematic review

[HTML] jnmjournal.org

HK Jung - 2011 - jnmjournal.org

... H, Iwasaki E, Yokoyama H, Sugino Y, Hibi T. Serum lipid levels are positively associated with non ... Obesity as a risk factor for GERD in Japan. ... Risk factors for gastroesophageal reflux disease, reflux esophagitis and non-erosive reflux disease among Chinese patients undergoing ...

被引用次数: 109 相关文章 所有 12 个版本 引用 保存 更多

Racial and geographic issues in gastroesophageal reflux disease

[HTML] nature.com

P Sharma, S Wani, Y Romero, D Johnson... - The American journal of ..., 2008 - nature.com

... has increased markedly in recent years (100), appears to be an important risk factor for GERD ... the weight gain (>5 kg, adjusted OR 3.0, 95% CI 1.6–6) was associated with new ... Prevalence of and risk factors for Helicobacter pylori infection in a multi-racial dyspeptic Malaysian ...

被引用次数: 67 相关文章 所有 7 个版本 引用 保存

Epidemiologic risk factors for Barrett's esophagus and associated adenocarcinoma

[HTML] cghjournal.org

A Wong, RC Fitzgerald - Clinical Gastroenterology and Hepatology, 2005 - Elsevier

... In addition, the altered body composition associated with obesity affects the pharmacokinetics of drugs. ... With regard to specific dietary factors, although there have been numerous studies assessing ... They showed that a complex risk factor score (including tobacco use, alcohol, fat ...

被引用次数: 132 相关文章 所有 6 个版本 引用 保存

[HTML] Geographical and ethnic differences in gastro - oesophageal reflux disease

[HTML] wiley.com

JY Kang - Alimentary pharmacology & therapeutics, 2004 - Wiley Online Library

... Details of published studies satisfying the inclusion criteria on the prevalence of symptoms associated with GERD ... Although body mass index, age and sex were risk factors for oesophagitis and hiatus hernia ... from 1.02 to 1.11, race was by far the most important risk factor (4.04 for ...

被引用次数: 104 相关文章 所有 4 个版本 引用 保存

Environmental–Lifestyle related factors

[HTML] nih.gov

S Roman, JE Pandolfino - Best Practice & Research Clinical ..., 2010 - Elsevier

Fig. 1. The environmental factors interact with the different mechanisms involved in GORD